

# Spectrum Generating Symmetries of Magical and Maximal Supergravities and Their Unitary Representations

Murat Günaydin

Paul Fest, July 2-3, 2012  
Cambridge University

- ▶ Review of magical supergravities in various dimensions.
- ▶ Orbits of extremal BPS and non-BPS black hole solutions in five dimensions and conformal extensions of  $5d$  U-duality groups as spectrum generating symmetry group.
- ▶ Orbits of extremal BPS and non-BPS black hole solutions in four dimensions and quasiconformal extensions of  $4d$  U-duality groups as spectrum generating symmetry group.
- ▶ Quantum spectra of extremal  $4d$  black holes and unitary representations of their  $3d$  U-duality groups
- ▶ Quantization of quasiconformal group actions, minimal unitary realizations and harmonic superspace
- ▶ Minimal unitary representations of noncompact groups and supergroups and *AdS/CFT* dualities
- ▶ Quantum spectra of magical and maximal sugras and the minimal unitary realizations of their  $3d$  U-duality groups.
- ▶ Open problems

- ▶ Bosonic part of the 5D Maxwell-Einstein supergravity Lagrangian (MESGT)

MG, Sierra and Townsend (1983)

$$e^{-1}\mathcal{L} = -\frac{1}{2}R - \frac{1}{4}\overset{\circ}{a}_{IJ}F_{\mu\nu}^I F^{J\mu\nu} - \frac{1}{2}g_{xy}(\partial_\mu\varphi^x)(\partial^\mu\varphi^y) + \frac{e^{-1}}{6\sqrt{6}}C_{IJK}\varepsilon^{\mu\nu\rho\sigma\lambda}F_{\mu\nu}^I F_{\rho\sigma}^J A_\lambda^K$$

coupling of  $(n_V - 1)$  vector multiplets  $(A_\mu^a, \lambda^{ai}, \varphi^a)$  to  $N = 2$  supergravity  $(g_{\mu\nu}, \psi_\mu^i, A_\mu)$  ( $I, J, K = 1, \dots, n_V, i=1,2, x, a = 1, \dots, (n_V - 1)$ )

- ▶ 5D,  $N = 2$  MESGT is uniquely determined by the constant symmetric tensor  $C_{IJK}$ .
- ▶ 5D MESGTs with symmetric scalar manifolds  $G/H$  such that  $G$  is a symmetry of the Lagrangian  $\iff C_{IJK}$  is given by the norm (determinant)  $\mathcal{N}_3$  of a Euclidean Jordan algebra  $J$  of degree 3.

$$\mathcal{N}_3(J) = C_{IJK}h^I h^J h^K$$

Euclidean  $J : \iff X^2 + Y^2 = 0 \implies X = Y = 0 \forall X, Y \in J$

# Symmetry Groups of Simple Jordan algebras

$n \times n$  Hermitian matrices over the division algebra  $\mathbb{A}$  form a Jordan algebra  $J_n^{\mathbb{A}}$  under the symmetric product  $A \cdot B \equiv 1/2(AB + BA)$ .

$J$	$Rot(J)$	$Lor(J)$	$Conf(J)$
$J_2^{\mathbb{C}}$	$SU(2)$	$SL(2, \mathbb{C})$	$SU(2, 2)$
$J_n^{\mathbb{R}}$	$SO(n)$	$SL(n, \mathbb{R})$	$Sp(2n, \mathbb{R})$
$J_n^{\mathbb{C}}$	$SU(n)$	$SL(n, \mathbb{C})$	$SU(n, n)$
$J_n^{\mathbb{H}}$	$USp(2n)$	$SU^*(2n)$	$SO^*(4n)$
$J_3^{\mathbb{O}}$	$F_4$	$E_{6(-26)}$	$E_{7(-25)}$
$\Gamma_{(1,d)}$	$SO(d)$	$SO(d, 1)$	$SO(d, 2)$

**Table:** The complete list of simple Euclidean Jordan algebras and their rotation ( automorphism), "Lorentz" ( reduced structure) and "Conformal" ( linear fractional) groups. The symbols  $\mathbb{R}$ ,  $\mathbb{C}$ ,  $\mathbb{H}$ ,  $\mathbb{O}$  represent the four division algebras.  $J_n^{\mathbb{A}}$  denotes a Jordan algebra of  $n \times n$  hermitian matrices over  $\mathbb{A}$ .  $\Gamma_{(1,d)}$  denotes the Jordan algebra of Dirac gamma matrices.

- ▶ Unified  $N = 2$  Maxwell-Einstein Supergravity theories in  $5d \Leftrightarrow$  all the vectors fields including the graviphoton transform in an irreducible representation of a simple U-duality group of the action.
- ▶ There exist only four unified MESGTs in  $d = 5$  with symmetric target spaces . They are defined by the four simple Euclidean Jordan algebras  $J_3^{\mathbb{A}}$  of  $3 \times 3$  Hermitian matrices over  $\mathbb{R}, \mathbb{C}, \mathbb{H}$  and  $\mathbb{O}$  and describe the coupling of 5, 8, 14 and 26 vector multiplets to supergravity. Their symmetries in 5,4 and 3 dimensions give the groups of the Magic Square of Freudenthal, Rozenfeld and Tits  $\implies$  *Magical supergravity theories* ( GST 1983)
- ▶ Scalar manifolds of five dimensional magical sugras are the irreducible symmetric spaces

$$\begin{array}{rcccccc}
 J & = & J_3^{\mathbb{R}} & & J_3^{\mathbb{C}} & & J_3^{\mathbb{H}} & & J_3^{\mathbb{O}} \\
 \mathcal{M}_5 & = & SL(3, \mathbb{R})/SO(3) & & SL(3, \mathbb{C})/SU(3) & & SU^*(6)/USp(6) & & E_{6(-26)}/F_4
 \end{array}$$

- ▶ The generic Jordan family defined by non-simple Jordan algebras  $J = \Gamma_{(1, n-1)} \oplus \mathbb{R}$  has the scalar manifold  $(SO(n-1, 1) \times SO(1, 1)) / SO(n-1)$
- ▶ In addition there exist three infinite families of unified MESGT's in 5d whose scalar manifolds are neither symmetric nor homogeneous! They are defined by Lorentzian Jordan of degree  $n \geq 3$  generated by  $n \times n$  matrices over  $\mathbb{R}, \mathbb{C}, \mathbb{H}$  Hermitian with respect to the Lorentzian metric  $\eta$  (MG, Zagermann 2003)

$$(A\eta)^\dagger = A\eta$$

- ▶ Under dimensional reduction of 5d MESGTs defined by Euclidean Jordan algebras to four dimensions one has

$$\mathcal{M}_5 = \frac{\text{Lor}(J)}{\text{Rot}(J)} \Rightarrow \mathcal{M}_4 = \frac{\text{Conf}(J)}{\widetilde{\text{Lor}}(J) \times U(1)}$$

where  $\widetilde{\text{Lor}}(J)$  is the compact real form of the Lorentz group of  $J$ .

- ▶ The bosonic sector of dimensionally reduced Lagrangian is

$$\mathcal{L}^{(4)} = -\frac{1}{2}R - g_{IJ}(\partial_\mu z^I)(\partial^\mu \bar{z}^J) + \frac{1}{4}\text{Im}(\mathcal{N}_{AB})F_{\mu\nu}^A F^{\mu\nu B} - \frac{1}{8}\text{Re}(\mathcal{N}_{AB})\epsilon^{\mu\nu\rho\sigma} F_{\mu\nu}^A F_{\rho\sigma}^B$$

In 5D: Vector fields  $A_\mu^I \Leftrightarrow$  Elements of Jordan algebra  $J$

In 4D:  $F_{\mu\nu}^A \oplus \tilde{F}_{\mu\nu}^A \Leftrightarrow$  Freudenthal triple system (FTS)  $\mathcal{F}(J)$  :

$$\mathcal{F}(J) \ni X = \left| \begin{array}{cc} \mathbb{R} & \mathbf{J} \\ \tilde{\mathbf{J}} & \mathbb{R} \end{array} \right| \Leftrightarrow \left| \begin{array}{cc} F_{\mu\nu}^0 & F_{\mu\nu}^I \\ \tilde{F}_{\mu\nu}^I & \tilde{F}_{\mu\nu}^0 \end{array} \right|$$

Automorphism group of  $\mathcal{F}(J) \cong$  Conformal group of  $J$

$N = 2$  MESGTs reduce to  $N = 4$  supersymmetric sigma models coupled to gravity in  $d = 3$ . The target manifolds of magical supergravity theories in  $d = 3$  are the exceptional quaternionic symmetric spaces:

- ▶  $\frac{F_{4(4)}}{USp(6) \times USp(2)}$ ,  $\frac{E_{6(2)}}{SU(6) \times SU(2)}$ ,  $\frac{E_{7(5)}}{SO(12) \times SU(2)}$ ,  $\frac{E_{8(-24)}}{E_7 \times SU(2)}$
- ▶ The generic Jordan family of MESGT reduced to  $d = 3$  have the target spaces:  $\frac{SO(n+2,4)}{SO(n+2) \times SO(4)}$
- ▶ Pure  $N = 2$  supergravity reduces to  $N = 4$  sigma model with target space  $\frac{G_{2(2)}}{SO(4)}$ .

### Exceptional $N = 2$ versus Maximal $N = 8$ Supergravity:

- ▶ The exceptional  $N = 2$  supergravity is defined by the exceptional Jordan algebra  $J_3^{\mathbb{O}}$  of  $3 \times 3$  Hermitian matrices over real octonions  $\mathbb{O}$ . Its global invariance group in  $5D$  is  $E_{6(-26)}$  with maximal compact subgroup  $F_4$ .
- ▶ The C-tensor  $C_{IJK}$  of  $N = 8$  supergravity in five dimensions can be identified with the symmetric tensor given by the cubic norm of the split exceptional Jordan algebra  $J_3^{\mathbb{O}_s}$  defined over split octonions  $\mathbb{O}_s$ . Its global invariance group in  $5D$  is  $E_{6(6)}$  with maximal compact subgroup  $USp(8)$ .
- ▶ In  $D = 4$  and  $D = 3$  the exceptional supergravity has  $E_{7(-25)}$  and  $E_{8(-24)}$  as its U-duality group while the maximal  $N = 8$  supergravity has  $E_{7(7)}$  and  $E_{8(8)}$ , respectively.

$D = 6$		$D = 5$		$D = 4$		$D = 3$
$\frac{SO(9,1)}{SO(9)}$	$\longrightarrow$	$\frac{E_{6(-26)}}{F_4}$	$\longrightarrow$	$\frac{E_{7(-25)}}{E_6 \times SO(2)}$	$\longrightarrow$	$\frac{E_{8(-24)}}{E_7 \times SU(2)}$
$\frac{SO(5,1)}{SO(5)}$	$\longrightarrow$	$\frac{SU^*(6)}{USp(6)}$	$\longrightarrow$	$\frac{SO^*(12)}{U(6)}$	$\longrightarrow$	$\frac{E_{7(-5)}}{SO(12) \times SU(2)}$
$\frac{SO(3,1)}{SO(3)}$	$\longrightarrow$	$\frac{SL(3, \mathbb{C})}{SO(3)}$	$\longrightarrow$	$\frac{SU(3,3)}{SU(3) \times SU(3) \times U(1)}$	$\longrightarrow$	$\frac{E_{6(+2)}}{SU(6) \times SU(2)}$
$\frac{SO(2,1)}{SO(2)}$	$\longrightarrow$	$\frac{SL(3, \mathbb{R})}{SO(3)}$	$\longrightarrow$	$\frac{Sp(6, \mathbb{R})}{U(3)}$	$\longrightarrow$	$\frac{F_{4(+4)}}{USp(6) \times USp(2)}$

Table: Scalar target spaces of magical supergravities in 6, 5, 4 and 3 dimensions.

6d origins of magical supergravities was first investigated by Romans and later by van Proeyen and others.



# Magical Supergravity Theories in 6 Dimensions coupled to Hypermultiplets and Their Gaugings

MG, Samtleben , Sezgin , 1012.1818

Chiral  $N = (1,0)$  supergravity coupled to  $n_T$  tensor multiplets,  $n_V$  vector multiplets and  $n_H$  hypermultiplets:

supergravity and tensor multiplets :  $\{e_\mu^m, \psi_\mu^i, B_{\mu\nu}^l, \chi^{ai}, L^l\}$ ,

vector multiplets :  $\{A_\mu^A, \lambda^{Ai}\}$ ,

hypermultiplets :  $\{\phi^X, \psi^r\}$ ,

$l = 0, 1, \dots, n_T$ ,  $a = 1, \dots, n_T$ ,  $A = 1, \dots, n_V$ ,  $X = 1, \dots, 4n_H$ ,  $r = 1, \dots, 2n_H$ . The gravitino, tensorino and gaugino are all doublets of the R-symmetry group  $USp(2)_R$  ( $i = 1, 2$ ). All fermions are symplectic Majorana-Weyl, where  $(\psi_\mu^i, \lambda^{Ai})$  have positive chirality and  $(\chi^{ai}, \psi^r)$  have negative chirality.  $L^l$  denotes a representative of the coset space  $SO(n_T, 1)/SO(n_T)$  parameterized by  $n_T$  real scalars of tensor multiplets.

- ▶  $n_T$  scalars of tensor multiplets parameterize the coset  $SO(n_T, 1)/SO(n_T)$ .
- ▶ Magical supergravities have  $n_T = 2, 3, 5, 9$  tensor multiplets with the vectors transforming in the spinor representation of  $SO(n_T, 1)$

$G_T$	$\mathcal{R}_v$	$A_\mu^A$	$\Gamma_{AB}^I$	$\mathcal{R}_{ten}$
$SO(9, 1)$	$16_c$	MW	$\Gamma_{AB}^I$	<b>10</b>
$SO(5, 1) \times USp(2)$	$(4_c, 2)$	SMW, $A = (\alpha r)$	$\Gamma_{\alpha r, \beta s}^I = \Gamma_{\alpha\beta}^I \epsilon_{rs}$	<b>(6, 1)</b>
$SO(3, 1) \times U(1)$	$(2, 1)_+ + (1, 2)_-$	W, $A = \{\alpha, \dot{\beta}\}$	$\begin{pmatrix} 0 & \Gamma_{\alpha\dot{\beta}}^I \\ \bar{\Gamma}_{\dot{\alpha}\beta}^I & 0 \end{pmatrix}$	<b>(2, 2)<sub>0</sub></b>
$SO(2, 1)$	<b>2</b>	M	$\Gamma_{AB}^I$	<b>3</b>

**Table:**  $G_T$  = global symmetry groups,  $\mathcal{R}_v$  = the representation of vector fields under  $G_T$ ;  $\mathcal{R}_{ten}$  = representation of tensors under  $G_T$ .

- ▶ Magical supergravities satisfy

$$\Gamma_{I(AB} \Gamma_{C)D}^I = 0$$

which are the Fierz identities of supersymmetric Yang-Mills theories in the critical dimensions 3, 4, 6, 10 and follow from the adjoint identities satisfied by their  $C_{IJK}$  tensors (Sierra 1987). Hence the Abelian gauge anomaly vanishes:

$$\mathcal{A} = c \epsilon^{\mu\nu\lambda\rho\sigma\epsilon} \Gamma_{IAB} \Gamma_{CD}^I \Lambda^A F_{\mu\nu}^B F_{\lambda\rho}^C F_{\sigma\epsilon}^D$$

## M/Superstring Theoretic Origins of Magical Sugras

Quaternionic magical 4d MESGT without any hypermultiplets can be obtained from IIB superstring by compactification on an orbifold of  $T^4 \times S^1 \times S^1$ . It has the same bosonic field content as the  $N = 6$  supergravity. Sen, Vafa (1995) & MG 2006

The 6d octonionic magical MESGT coupled to 28 hypermultiplets parametrizing the quaternionic symmetric space  $E_{8(-24)}/E_7 \times SU(2)$  is free of gravitational anomalies. ( $n_H - n_V + 29n_T = 273$ ). Its 4d scalar manifold contains the moduli space of FHSV model which is obtained from IIB superstring by compactification on a self-mirror Calabi-Yau manifold with Hodge numbers  $h^{(1,1)} = h^{(2,1)} = 11$  : MG (Paris 2006).

$$\mathcal{M}_V \times \mathcal{M}_H = \frac{E_{7(-25)}}{E_6 \times U(1)} \times \frac{E_{8(-24)}}{E_7 \times SU(2)} \supset \frac{SO(10,2) \times SU(1,1)}{SO(10) \times U(1) \times U(1)} \times \frac{SO(12,4)}{SO(12) \times SO(4)}$$

Bianchi and Ferrara (2008) reconsidered the string derivation of FSHV model over the Enriques Calabi-Yau and argued that the octonionic magical MESGT admits a string interpretation closely related to the Enriques model.

Type IIB superstring theory compactified over a CY threefold constructed by Todorov leads to the complex magical  $N = 2$  MESGT in  $d = 4$  with moduli space

$$\mathcal{M}_4 = \frac{SU(3,3)}{SU(3) \times SU(3) \times U(1)}$$

coupled to  $(h^{(1,1)} + 1) = 30$  hypermultiplets.

The hyper-free  $N = 2$  string models based on asymmetric orbifolds with  $N = (4, 1)$  worldsheet superconformal symmetry using 2D fermionic construction given by Dolivet, Julia and Kounnas include the complex and quaternionic magical supergravities in  $d = 4$ .

# U-duality Orbits of Extremal , Spherically Symmetric Stationary Black Hole Solutions of 5D Supergravity Theories with Symmetric Target Spaces: ( MG and Ferrara, 1997 )

- ▶ The black hole potential that determines the attractor flow takes on the following form for  $N = 2$  MESGTs: ( Ferrara, Gibbons, Kallosh, Strominger )

$$V(\phi, q) = q_I \overset{\circ}{a}{}^{IJ} q_J$$

where  $\overset{\circ}{a}{}_{IJ}$  is the "metric" of the kinetic energy term of the vector fields. The  $(n + 1)$  dimensional charge vector in an extremal BH background is given by

$$q_I = \int_{S^3} H_I = \int_{S^3} \overset{\circ}{a}{}_{IJ} *F^J \quad (I = 0, 1, \dots, n)$$

The entropy  $S$  of an extremal black hole solution of  $N = 2$  MESGT with charges  $q_I$  is determined by the value of the black hole potential  $V$  at the attractor points

$$S_{BPS} = (V_{critical})^{3/4} = (C^{IJK} q_I q_J q_K)^{3/4}$$

$J$	$\mathcal{O}_{BPS} = Str_0(J)/Aut(J)$
$J_3^{\mathbb{R}}$	$SL(3, \mathbb{R})/SO(3)$
$J_3^{\mathbb{C}}$	$SL(3, \mathbb{C})/SU(3)$
$J_3^{\mathbb{H}}$	$SU^*(6)/USp(6)$
$J_3^{\mathbb{O}}$	$E_{6(-26)}/F_4$
$\mathbb{R} \oplus \Gamma_{(1, n-1)}$	$SO(n-1, 1) \times SO(1, 1)/SO(n-1)$

**Table:** Orbits of spherically symmetric stationary BPS black hole solutions in 5D MESGTs defined by Euclidean Jordan algebras  $J$  of degree three. U-duality and stability groups are given by the Lorentz ( reduced structure ) and rotation ( automorphism) groups of  $J$ .

$J$	$\mathcal{O}_{non-BPS} = Str_0(J)/Aut(J_{(1,2)})$
$J_3^{\mathbb{R}}$	$SL(3, \mathbb{R})/SO(2, 1)$
$J_3^{\mathbb{C}}$	$SL(3, \mathbb{C})/SU(2, 1)$
$J_3^{\mathbb{H}}$	$SU^*(6)/USp(4, 2)$
$J_3^{\mathbb{O}}$	$E_{6(-26)}/F_{4(-20)}$
$\mathbb{R} \oplus \Gamma_{(1, n-1)}$	$SO(n-1, 1) \times SO(1, 1)/SO(n-2, 1)$

**Table:** Orbits of non-BPS extremal BHs with non-zero entropy.

# The orbits of BPS black hole solutions of $N = 8$ supergravity theory in five dimensions.

The  $1/8$  BPS black holes with non-vanishing entropy has the orbit

$$\mathcal{O}_{1/8-BPS} = \frac{E_{6(6)}}{F_{4(4)}}$$

Maximal supergravity theory admits  $1/4$  and  $1/2$  BPS black hole solutions with vanishing entropy. Their orbits under U-duality are

$$\mathcal{O}_{1/4-BPS} = \frac{E_{6(6)}}{O(5,4) \otimes T_{16}}$$

$$\mathcal{O}_{1/2-BPS} = \frac{E_{6(6)}}{O(5,5) \otimes T_{16}}$$

Vanishing entropy means vanishing cubic norm. Thus the black hole solutions corresponding to vanishing entropy has additional symmetries beyond the five dimensional U-duality group.

$\implies$  They are invariant under the generalized special conformal transformations of the underlying Jordan algebras.

- ▶ The proposal: the conformal groups  $Conf[J]$  of underlying Jordan algebras  $J$  of supergravity theories must act as spectrum generating symmetry groups.

$Conf[J]$  leaves invariant light-like separations with respect to a cubic distance function  $\mathcal{N}_3(J_1 - J_2)$  and admits a 3-grading with respect to their Lorentz subgroups

$$Conf[J] = K_J \oplus Lor(J) \times \mathcal{D} \oplus T_J$$

$Lor(J)$  is the 5D U-duality group that leaves the cubic norm invariant.

- ▶  $Conf[J] \Leftrightarrow$  U-duality group of corresponding 4D supergravity.  
 $\Rightarrow$  4D U-duality groups must act as spectrum generating symmetry groups of corresponding five dimensional supergravity theories.

- ▶ U-duality group  $G_4$  of a 4D Maxwell-Einstein supergravity defined by a Jordan algebra  $J$  of degree three  $\Leftrightarrow$   
 $G_4 \equiv \text{Aut}(\mathcal{F}(J)) \equiv \text{Conf}[J]$
- ▶ The entropy of an extremal dyonic black hole with charges  $(p^0, p^I, q_0, q_I)$  is given by the quartic invariant  $Q_4(q, p)$  of the Freudenthal triple system  $\mathcal{F}(J)$ .
- ▶ black hole attractor equations  $\Rightarrow$  criticality conditions for black hole scalar potential

$$V_{BH} \equiv |Z|^2 + G^{I\bar{J}}(D_I Z)(\bar{D}_{\bar{J}} \bar{Z})$$

$Z \equiv$  central charge function.



$$\partial_I V_{BH} = 0$$

implies

$$2\bar{Z}D_I Z + iC_{IJK}G^{J\bar{J}}G^{K\bar{K}}\bar{D}_{\bar{J}}\bar{Z}\bar{D}_{\bar{K}}\bar{Z} = 0$$



J	$\frac{1}{2}$ -BPS orbits $\mathcal{O}_{\frac{1}{2}-BPS}$	non-BPS, $Z \neq 0$ orbits $\mathcal{O}_{non-BPS, Z \neq 0}$	non-BPS, $Z = 0$ orbits $\mathcal{O}_{non-BPS, Z=0}$
–	$\frac{SU(1, n+1)}{SU(n+1)}$	–	$\frac{SU(1, n+1)}{SU(1, n)}$
$\mathbb{R} \oplus \Gamma_{(1, n-1)}$	$\frac{SU(1,1) \otimes SO(2, 2+n)}{SO(2) \otimes SO(2+n)}$	$\frac{SU(1,1) \otimes SO(2, 2+n)}{SO(1,1) \otimes SO(1, 1+n)}$	$\frac{SU(1,1) \otimes SO(2, 2+n)}{SO(2) \otimes SO(2, n)}$
$J_3^{\mathcal{O}}$	$\frac{E_7(-25)}{E_6}$	$\frac{E_7(-25)}{E_6(-26)}$	$\frac{E_7(-25)}{E_6(-14)}$
$J_3^{\mathbb{H}}$	$\frac{SO^*(12)}{SU(6)}$	$\frac{SO^*(12)}{SU^*(6)}$	$\frac{SO^*(12)}{SU(4, 2)}$
$J_3^{\mathbb{C}}$	$\frac{SU(3, 3)}{SU(3) \otimes SU(3)}$	$\frac{SU(3, 3)}{SL(3, \mathbb{C})}$	$\frac{SU(3, 3)}{SU(2, 1) \otimes SU(1, 2)}$
$J_3^{\mathbb{R}}$	$\frac{Sp(6, \mathbb{R})}{SU(3)}$	$\frac{Sp(6, \mathbb{R})}{SL(3, \mathbb{R})}$	$\frac{Sp(6, \mathbb{R})}{SU(2, 1)}$

**Table:** Non-degenerate orbits of  $N = 2$ ,  $D = 4$  MESGTs with symmetric scalar manifolds. Except for the first row all such theories originate from five dimensions and are defined by Jordan algebras that are indicated in the first column. (MG & Ferrara, 1997) ( Bellucci, Ferrara , MG, Marrani , 2006 )

- ▶ The orbits of BH solutions of 4D  $N = 8$  supergravity under  $E_{7(7)}$ :

$$I_4 > 0 : \mathcal{O}_{\frac{1}{8}\text{-BPS}} = \frac{E_{7(7)}}{E_{6(2)}} \iff \frac{1}{8}\text{-BPS};$$

$$I_4 < 0 : \mathcal{O}_{\text{non-BPS}} = \frac{E_{7(7)}}{E_{6(6)}} \iff \text{non-BPS}.$$

Generic light-like orbit with 1 vanishing eigenvalue:

$$\frac{E_{7(7)}}{F_{4(4)} \otimes T_{26}}$$

Critical light-like orbit with 2 vanishing eigenvalues:

$$\frac{E_{7(7)}}{O(6,5) \otimes (T_{32} \oplus T_1)}$$

Doubly critical light-like orbit with 3 vanishing eigenvalues :

$$\frac{E_{7(7)}}{E_{6(6)} \otimes T_{27}}$$

- ▶ Classification of the orbits of spherically symmetric stationary extremal black holes of  $4D$ ,  $N = 8$  sugra and  $4D$ ,  $N = 2$  MESGTs with symmetric target spaces. ( Ferrara and MG , 1997 ).
- ▶ Question: Can the  $3D$  U-duality groups act as spectrum generating conformal symmetries of corresponding  $4D$  supergravity theories ? ( MG, Koepsell, Nicolai 1997 )
- ▶ No conformal realization for any real forms of  $E_8, G_2$  and  $F_4$   
 $\Leftrightarrow$  No 3-grading with respect to a subgroup of maximal rank.
- ▶ However, all simple Lie algebras admit a 5-grading with respect to a subalgebra of maximal rank

$$\mathfrak{g} = \mathfrak{g}^{-2} \oplus \mathfrak{g}^{-1} \oplus \mathfrak{g}^0 \oplus \mathfrak{g}^{+1} \oplus \mathfrak{g}^{+2}$$

such that the grade  $\pm 2$  subspaces are one-dimensional.

► QUASICONFORMAL REALIZATION OF  $E_{8(8)}$  MG, Koepsell, Nicolai, 2000

$$\begin{aligned}
 E_{8(8)} &= \mathfrak{1}_{-2} \oplus 5\mathfrak{6}_{-1} \oplus E_{7(7)} + SO(1, 1) \oplus 5\mathfrak{6}_{+1} \oplus \mathfrak{1}_{+2} \\
 \mathfrak{g} &= \tilde{K} \oplus \tilde{U}_A \oplus [S_{(AB)} + \Delta] \oplus U_A \oplus K
 \end{aligned}$$

over a space  $\mathcal{T}$  coordinatized by the elements  $X$  of the exceptional FTS  $\mathcal{F}(J_3^{\text{OS}})$  plus an extra singlet variable  $x$ :  $5\mathfrak{6}_{+1} \oplus \mathfrak{1}_{+2} \Leftrightarrow (X, x) \in \mathcal{T}$ :

$$\begin{aligned}
 K(X) &= 0 & U_A(X) &= A & S_{AB}(X) &= (A, B, X) \\
 K(x) &= 2 & U_A(x) &= \langle A, X \rangle & S_{AB}(x) &= 2 \langle A, B \rangle x \\
 \tilde{U}_A(X) &= \frac{1}{2} (X, A, X) - Ax \\
 \tilde{U}_A(x) &= -\frac{1}{6} \langle (X, X, X), A \rangle + \langle X, A \rangle x \\
 \tilde{K}(X) &= -\frac{1}{6} (X, X, X) + Xx \\
 \tilde{K}(x) &= \frac{1}{6} \langle (X, X, X), X \rangle + 2x^2
 \end{aligned}$$

Freudenthal triple product  $\Leftrightarrow (X, Y, Z)$

Skew-symmetric invariant form  $\Leftrightarrow \langle X, Y \rangle = -\langle Y, X \rangle$

Quartic invariant of  $E_{7(7)}$   $\Leftrightarrow \langle (X, X, X), X \rangle$

$A, B, .. \in \mathcal{F}(J_3^{\text{OS}})$

- ▶ Geometric meaning of the quasiconformal action of the Lie algebra  $\mathfrak{g}$  on the space  $\mathcal{T}$ ?
- ▶ Define a quartic norm of  $\mathcal{X} = (X, x) \in \mathcal{T}$  as
 
$$\mathcal{N}_4(\mathcal{X}) := Q_4(X) - x^2$$
 $Q_4(X)$  is the quartic norm of the underlying Freudenthal system and  $X \in \mathcal{F}$ .
- ▶ Define a quartic “distance” function between any two points  $\mathcal{X} = (X, x)$  and  $\mathcal{Y} = (Y, y)$  in  $\mathcal{T}$  as

$$d(\mathcal{X}, \mathcal{Y}) := \mathcal{N}_4(\delta(\mathcal{X}, \mathcal{Y}))$$

$\delta(\mathcal{X}, \mathcal{Y})$  is the “symplectic” difference of  $\mathcal{X}$  and  $\mathcal{Y}$  :

$$\delta(\mathcal{X}, \mathcal{Y}) := (X - Y, x - y + \langle X, Y \rangle) = -\delta(\mathcal{Y}, \mathcal{X})$$

- ▶ Light-like separations  $d(\mathcal{X}, \mathcal{Y}) = 0$  are left invariant under quasiconformal group action.
  - Quasiconformal groups are the invariance groups of “light-cones” defined by a quartic distance function.
- ▶  $E_{8(8)}$  is the invariance group of a quartic light-cone in 57 dimensions!

- ▶ Scalar manifolds of  $N = 2$  MESGTs defined by Jordan algebras of degree three ( 8 real supersymmetries)

$$\mathcal{M}_5 = \frac{Lor(J)}{Rot(J)}$$

$$\mathcal{M}_4 = \frac{Conf(J)}{\widetilde{Lor(J)} \times U(1)}$$

$$\mathcal{M}_3 = \frac{QConf(J)}{\widetilde{Conf(J)} \times SU(2)}$$

- ▶ Quasiconformal extensions of  $4D$  U-duality groups  $\equiv 3D$  U-duality groups.
- ▶ Extremal black hole solution of  $4D$  ,  $N = 8$  sugra with 56 charges  $q_I, p^I$  and entropy  $s$ : light-like vectors in 57 dimensional charge-entropy space

$$\mathcal{N}_4(q_I, p^I, s) = 0 \rightarrow Q_4 = s^2$$

- ▶ Proposal:  $3D$  U-duality groups must act as spectrum generating symmetry groups of the extremal black hole solutions of  $4D$  supergravity theories.

$J$	$\mathcal{M}_5 =$ $\text{Lor}(J) / \text{Rot}(J)$	$\mathcal{M}_4 =$ $\text{Conf}(J) / \widetilde{\text{Lor}}(J) \times U(1)$	$\mathcal{M}_3 =$ $\text{QConf}(\mathcal{F}(J)) / \widetilde{\text{Conf}}(J) \times \text{SU}(2)$
$J_3^{\mathbb{R}}$	$\text{SL}(3, \mathbb{R}) / \text{SO}(3)$	$\text{Sp}(6, \mathbb{R}) / \text{U}(3)$	$\text{F}_{4(4)} / \text{USp}(6) \times \text{SU}(2)$
$J_3^{\mathbb{C}}$	$\text{SL}(3, \mathbb{C}) / \text{SU}(3)$	$\text{SU}(3, 3) / \text{S}(\text{U}(3) \times \text{U}(3))$	$\text{E}_{6(2)} / \text{SU}(6) \times \text{SU}(2)$
$J_3^{\mathbb{H}}$	$\text{SU}^*(6) / \text{USp}(6)$	$\text{SO}^*(12) / \text{U}(6)$	$\text{E}_{7(-5)} / \text{SO}(12) \times \text{SU}(2)$
$J_3^{\mathbb{O}}$	$\text{E}_{6(-26)} / \text{F}_4$	$\text{E}_{7(-25)} / \text{E}_6 \times \text{U}(1)$	$\text{E}_{8(-24)} / \text{E}_7 \times \text{SU}(2)$
$\mathbb{R} \oplus \Gamma_{(1, n-1)}$	$\frac{\text{SO}(n-1, 1) \times \text{SO}(1, 1)}{\text{SO}(n-1)}$	$\frac{\text{SO}(n, 2) \times \text{SU}(1, 1)}{\text{SO}(n) \times \text{SO}(2) \times \text{U}(1)}$	$\frac{\text{SO}(n+2, 4)}{\text{SO}(n+2) \times \text{SO}(4)}$

Table: Scalar manifolds  $\mathcal{M}_d$  of  $N = 2$  MESGT's defined by Jordan algebras  $J$  of degree 3 in  $d = 3, 4, 5$  dimensions.  $\widetilde{\text{Lor}}(J)$  and  $\widetilde{\text{Conf}}(J)$  denote the compact real forms of the Lorentz group  $\text{Lor}(J)$  and conformal group  $\text{Conf}(J)$  of a Jordan algebra  $J$ .  $\text{QConf}(\mathcal{F}(J))$  denotes the quasiconformal group associated with  $J$ .

- ▶ A concrete implementation of the proposal that 3D U-duality groups must act as spectrum generating quasisymmetric groups of spherically symmetric stationary BPS black holes of 4D supergravity theories:

MG, Neitzke, Pavlyk, Pioline, Waldrom 2005,2006

- ▶ Equations of motion for a spherically symmetric stationary black hole of four dimensional supergravity theories are equivalent to equations for geodesic motion of a fiducial particle on the moduli space  $\mathcal{M}_3^*$  of 3D supergravity obtained by reduction on a time-like circle.

Breitenlohner, Gibbons, Maison 1987

- ▶ 3D scalar manifold from compactification on a space-like (time-like) circle

$$\mathcal{M}_3 = \frac{G_3}{K_3} \quad \left( \mathcal{M}_3^* = \frac{G_3}{H_3} \right)$$

where  $K_3$  is the maximal compact subgroup of  $G_3$  and  $H_3$  is a noncompact real form of  $K_3$ . For  $N = 2$  MESGTs defined by Jordan algebras  $\mathcal{M}_3^*$  is a para-quaternionic symmetric space.



$n_Q$	$n_V$	$M_4$	$\mathcal{M}_3^*$	$J$
8	1	$\emptyset$	$\frac{U(2,1)}{U(1,1) \times U(1)}$	$\mathbb{R}$
8	2	$\frac{SL(2, \mathbb{R})}{U(1)}$	$\frac{G_{2,2}}{SO(2,2)}$	$\mathbb{R}$
8	7	$\frac{Sp(6, \mathbb{R})}{SU(3) \times U(1)}$	$\frac{F_{4(4)}}{Sp(6, \mathbb{R}) \times SL(2, \mathbb{R})}$	$J_3^{\mathbb{R}}$
8	10	$\frac{SU(3,3)}{SU(3) \times SU(3) \times U(1)}$	$\frac{E_{6(2)}}{SU(3,3) \times SL(2, \mathbb{R})}$	$J_3^{\mathbb{C}}$
8	16	$\frac{SO^*(12)}{SU(6) \times U(1)}$	$\frac{E_{7(-5)}}{SO^*(12) \times SL(2, \mathbb{R})}$	$J_3^{\mathbb{H}}$
8	28	$\frac{E_{7(-25)}}{E_6 \times U(1)}$	$\frac{E_{8(-24)}}{E_{7(-25)} \times SL(2, \mathbb{R})}$	$J_3^{\mathbb{O}}$
8	$n+2$	$\frac{SL(2, \mathbb{R})}{U(1)} \times \frac{SO(n,2)}{SO(n) \times SO(2)}$	$\frac{SO(n+2,4)}{SO(n,2) \times SO(2,2)}$	$\mathbb{R} \oplus \Gamma(1, n-1)$
16	$n+2$	$\frac{SL(2, \mathbb{R})}{U(1)} \times \frac{SO(n-4,6)}{SO(n-4) \times SO(6)}$	$\frac{SO(n-2,8)}{SO(n-4,2) \times SO(2,6)}$	$\mathbb{R} \oplus \Gamma(5, n-5)$
24	16	$\frac{SO^*(12)}{SU(6) \times U(1)}$	$\frac{E_{7(-5)}}{SO^*(12) \times SL(2, \mathbb{R})}$	$J_3^{\mathbb{H}}$
32	28	$\frac{E_{7(7)}}{SU(8)}$	$\frac{E_{8(8)}}{SO^*(16)}$	$J_3^{\mathbb{O}_s}$

**Table:** Above we give the number of supercharges  $n_Q$ , 4D vector fields  $n_V$ , scalar manifolds of supergravity theories before and after reduction along a timelike Killing vector from  $D = 4$  to  $D = 3$ , and associated Jordan algebras  $J$ . Isometry groups of 4D and 3D supergravity theories are given by  $Conf(J)$  and  $QConf(J)$ , of  $J$ , respectively.

- ▶ The quantization of the motion of fiducial particle on  $\mathcal{M}_3^*$  leads to quantum mechanical wave functions that provide the basis of a unitary representation of the isometry group  $G_3$  of  $\mathcal{M}_3^*$ .
- ▶ BPS black holes correspond to a special class of geodesics which lift holomorphically to the twistor space  $\mathcal{Z}_3$  of  $\mathcal{M}_3^*$ . Spherically symmetric stationary BPS black holes of  $N = 2$  MESGT's are described by holomorphic curves in  $\mathcal{Z}_3$
- ▶ The relevant unitary representations of the  $3D$  isometry groups  $QConf(J)$  for BPS black holes are those induced by their holomorphic actions on the corresponding twistor spaces  $\mathcal{Z}_3$ , which belong to quaternionic discrete series representations.
- ▶ For rank two quaternionic groups  $SU(2,1)$  and  $G_{2(2)}$  unitary representations induced by the geometric quasiconformal actions and their spherical vectors were studied in ( MG, Neitzke, Pavlyk , Pioline , 2007).

- ▶ Construction of the quaternionic discrete series representations of  $SU(2,1)$  and  $G_{2(2)}$  using the quasiconformal realizations twisted with a unitary character  $\nu$  : GNPP (2007)  
Starting point is the determination of the spherical vector  $|\mathcal{O}\rangle$  of the QCG group  $G$  with the maximal compact subgroup  $K$ :

$$K|\mathcal{O}\rangle = |\mathcal{O}\rangle$$

Consider the Verma module generated by the action of the noncompact generators  $P_a$  of  $G$ :

$$|\mathcal{O}\rangle \oplus P_a|\mathcal{O}\rangle \oplus P_a P_b|\mathcal{O}\rangle \oplus \dots$$

The quaternionic discrete series representations are obtained as submodules of the above Verma module for certain discrete values of  $\nu$ .

U-duality groups of magical sugras in  $d=3$  are the quaternionic real forms of exceptional groups  $F_4, E_6, E_7$  and  $E_8$ .

- ▶ Unified realization of 3D U-duality groups of all  $N = 2$  MESGTs defined by Jordan algebras as spectrum generating quasiconformal groups covariant with respect to their 5D U-duality groups. MG, Pavlyk (2009)
- ▶ Unified construction of the spherical vectors of quasiconformal realizations of  $F_{4(4)}$ ,  $E_{6(2)}$ ,  $E_{7(-5)}$ ,  $E_{8(-24)}$  and  $SO(d+2, 4)$  twisted by a unitary character  $\nu$  with respect to their maximal compact subgroups and determination of their quadratic Casimir operators
- ▶ For  $\nu = -(n_V + 2) + i\rho$  the quasiconformal action induces unitary representations that belong to the principle series. For special discrete values of  $\nu$  the quasiconformal action leads to unitary representations belonging to the quaternionic discrete series and their analytic continuations. ( $n_V =$  number of vector fields of the  $5d$ ,  $N = 2$  MESGT.)
- ▶ Extension of the above results to  $F_{4(4)}$ ,  $E_{6(6)}$ ,  $E_{7(7)}$ ,  $E_{8(8)}$  and of  $SO(n+3, m+3)$ . as QCGs covariant with respect to  $SL(3, R)$ ,  $SL(3, R) \times SL(3, R)$ ,  $SL(6, R)$ ,  $E_{6(6)}$  and  $SO(n, m) \times SO(1, 1)$ , respectively.  
 Quaternionic discrete series of  $E_{8(-24)} \Rightarrow$  "Octonionic discrete series" of  $E_{8(8)}$  !

## Minimal Unitary Representations and Quasiconformal Groups:

- ▶ Quantization of the quasiconformal realization of a non-compact Lie group leads directly to its minimal unitary representation  $\Rightarrow$  **Unitary representation over an Hilbert space of square integrable functions of smallest number of variables possible.**
- ▶ **Minimal unitary representation of  $E_{8(8)}$  over  $L^2(\mathbb{R}^{29})$  from its geometric realization as a quasiconformal group** MG, Koepsell & Nicolai 2000

$$E_{8(8)} = 1_{-2} \oplus 56_{-1} \oplus E_{7(7)} + SO(1, 1) \oplus 56_{+1} \oplus 1_{+2}$$

- ▶ **Minimal unitary representation of  $E_{8(-24)}$  over  $L^2(\mathbb{R}^{29})$  from its geometric realization as a quasiconformal group** MG, Pavlyk, 2004

$$E_{8(-24)} = 1_{-2} \oplus 56_{-1} \oplus E_{7(-25)} + SO(1, 1) \oplus 56_{+1} \oplus 1_{+2}$$

**56 of  $E_7 \Rightarrow 28$  coordinates and 28 momenta. These 28 coordinates plus the singlet coordinate yield the minimal number ( 29) of variables for  $E_8$ .**

# QCG Approach to Minimal Unitary Representations

GKN & MG, Pavlyk

- ▶ Lie algebra  $\mathfrak{g}$  of a quasiconformal realization of a group  $G$  can be decomposed as :

$$\mathfrak{g} = \mathfrak{g}^{-2} \oplus \mathfrak{g}^{-1} \oplus (\mathfrak{h} \oplus \Delta) \oplus \mathfrak{g}^{+1} \oplus \mathfrak{g}^{+2}$$

$$\mathfrak{g} = E \oplus E^\alpha \oplus (J^a + \Delta) \oplus F^\alpha \oplus F$$

$\Delta = -\frac{i}{2}(yp + py)$  ( $[y, p] = i$ ) determines the 5-grading and  $\Omega^{\alpha\beta}$  is the symplectic invariant tensor of  $\mathfrak{h}$  generated by  $J^a$  ( $\alpha, \beta, .. = 1, 2, \dots, 2n$ ) and  $[\xi^\alpha, \xi^\beta] = \Omega^{\alpha\beta}$

$$E = \frac{1}{2}y^2 \quad E^\alpha = y\xi^\alpha, \quad J^a = -\frac{1}{2}\lambda^a_{\alpha\beta}\xi^\alpha\xi^\beta$$

$$F = \frac{1}{2}p^2 + \frac{\kappa I_4(\xi^\alpha)}{y^2}, \quad F^\alpha = [E^\alpha, F]$$

$I_4(\xi^\alpha) = S_{\alpha\beta\gamma\delta}\xi^\alpha\xi^\beta\xi^\gamma\xi^\delta \Leftrightarrow$  *quartic invariant of  $\mathfrak{h}$*

Choosing a polarization  $\xi^\alpha = (x^i, p_j)$  one has  $[x^i, p_j] = i\delta_j^i$  ( $i, j = 1, 2, \dots, n$ )

Minimal Gelfand-Kirillov dimension for the minimal unitary representation is  $n + 1 \Leftrightarrow (x^i, y)$ .

- ▶  $(E, F, \Delta) \implies SL(2, \mathbb{R})$  of conformal quantum mechanics with the quartic invariant  $I_4$  playing the role of coupling constant.

# 4D, $N = 2$ $\sigma$ -models coupled to Supergravity in Harmonic Superspace and Minimal Unitary Representations

MG 2007

- In HSS the metric on a quaternionic target space is given by a quartic potential  $\mathcal{L}^{(+4)}$ . The action is Galperin, Ogievetsky 1993

$$S = \int d\zeta^{(-4)} du \{ Q_\alpha^+ D^{++} Q^{+\alpha} - q_i^+ D^{++} q^{+i} + \mathcal{L}^{(+4)}(Q^+, q^+, u^-) \}$$

$\zeta, u_i^\pm$  are analytic superspace coordinates

$Q_\alpha^+(\zeta, u)$ ,  $\alpha = 1, \dots, 2n$  are hypermultiplets and  $q_i^+(\zeta, u)$ , ( $i = 1, 2$ ) are supergravity compensators.  $Q_\alpha^+$  and  $q_i^+$  are analytic  $N = 2$  superfields.

"Hamiltonian mechanics" with  $D^{++}$  playing the role of time derivative and  $Q^+$  and  $q^+$  corresponding to phase space coordinates under Poisson brackets

$$\{f, g\} = \frac{1}{2} \Omega^{\alpha\beta} \frac{\partial f}{\partial Q^{+\alpha}} \frac{\partial g}{\partial Q^{+\beta}} - \frac{1}{2} \epsilon^{ij} \frac{\partial f}{\partial q^{+i}} \frac{\partial g}{\partial q^{+j}},$$

Isometries are generated by Killing potentials  $K_A(Q^+, q^+, u^-)$  that obey the "conservation law"  $\partial^{++} K_A + \{K_A, \mathcal{L}^{(+4)}\} = 0$

$$\mathcal{L}^{(+4)} = \frac{P^{(+4)}(Q^+)}{(q^+ u^-)^2}$$

$$P^{(+4)}(Q^+) = \frac{1}{12} S_{\alpha\beta\gamma\delta} Q^{+\alpha} Q^{+\beta} Q^{+\gamma} Q^{+\delta}$$

$S_{\alpha\beta\gamma\delta}$  is a completely symmetric invariant tensor of  $H$ .

The Killing potentials that generate the isometry group  $G$  are given by

$$\mathbf{Sp}(2): \quad K_{ij}^{++} = 2(q_i^+ q_j^+ - u_i^- u_j^- \mathcal{L}^{(+4)}),$$

$$\mathbf{H}: \quad K_a^{++} = t_{a\alpha\beta} Q^{+\alpha} Q^{+\beta},$$

$$\mathbf{G/H} \times \mathbf{Sp}(2): \quad K_{i\alpha}^{++} = 2q_i^+ Q_\alpha^+ - u_j^- (q^+ u^-) \delta_\alpha^- \mathcal{L}^{(+4)}$$

$N = 2$ $\sigma$ -model in HSS	Minrep of Isometry Grp $G$
compensator $w$	coordinate $y$
conjugate $p^{++}$	momentum $p$
$\{, \}$	$i[, ]$
harmonic superfields $Q^{+\alpha}$	symplectic coords $\xi^\alpha$
$P^{(+4)}(Q^+)$	$I_4(\xi)$
$K^{a++} = t_{\alpha\beta}^a Q^{+\alpha} Q^{+\beta}$	$J^a = \lambda_{\alpha\beta}^a \xi^\alpha \xi^\beta$
$T_\alpha^{++++} = \{T_\alpha^+, M^{++++}\}$	$F_\alpha = [E_\alpha, F]$
$T_\alpha^+ = -\sqrt{2}wQ_\alpha^+$	$E_\alpha = y \xi_\alpha$
$M^{++++} = \frac{1}{2}(p^{++})^2 - \frac{P^{(+4)}(Q^+)}{w^2}$	$F = \frac{1}{2}p^2 + \frac{\kappa(I_4(\xi^\alpha))}{y^2}$
$M^0 = \frac{1}{2}w^2$	$E = \frac{1}{2}y^2$
$M^{++} = \frac{1}{2}(wp^{++} + p^{++}w)$	$\Delta = -\frac{i}{2}(yp + py)$

**Table:** The correspondence between the harmonic superspace formulation of  $N = 2$  sigma models coupled to supergravity and the minimal unitary realizations of their isometry groups  $G$  obtained by quantization of quasiconformal action of  $G$ .



- ▶ The Poisson brackets (PB)  $\{, \}$  in HSS formulation go over to  $i$  times the commutator  $[, ]$  in the minimal unitary realization and the harmonic superfields  $w, p^{++}$  corresponding to supergravity hypermultiplet compensators, that are canonically conjugate under PB map to the canonically conjugate coordinate and momentum operators  $y, p$ . Similarly, the harmonic superfields  $Q^{+\alpha}$  that form  $N/2$  conjugate pairs under Poisson brackets map into the symplectic bosonic oscillators  $\xi^\alpha$  on the MINREP side. One finds a normal ordering ambiguity in the quantum versions of the quartic invariants. The classical expression relating the quartic invariant polynomial  $P^{(+4)}$  to the quadratic Casimir function in HSS formulation differs from the expression relating the quartic quantum invariant  $I_4$  to the quadratic Casimir of  $H$  by an additive c-number depending on the ordering chosen.
- ▶ On the MINREP side we are working with a realization in terms of quantum mechanical coordinates and momenta, while in HSS side the corresponding quantities are classical harmonic analytic superfields. The above correspondence can be extended to the full quantum correspondence on both sides by reducing the  $4D N = 2 \sigma$  model to one dimension and quantizing it to get a supersymmetric quantum mechanics (with 8 supercharges). The bosonic spectrum of the corresponding quantum mechanics must furnish a minimal unitary representation of the isometry group, which extends to a fully supersymmetric spectrum.

- ▶ The above mapping implies that the *fundamental spectra* of the "quantum"  $N = 2$ , quaternionic Kähler  $\sigma$  models coupled to sugra in  $d = 4$  must fit into the minimal unitary representations of their isometry groups.
- ▶ The  $N = 2$ ,  $d = 4$  MESGT's, under dimensional reduction, lead to  $d = 3$  supersymmetric  $\sigma$  models with quaternionic Kähler manifolds  $\mathcal{M}_3$  (C-map). After T-dualizing the three dimensional theory one can lift it back to four dimensions, thereby obtaining an  $N = 2$  sigma model coupled to supergravity that is in the mirror image of the original  $N = 2$  MESGT.
- ▶ The above results suggest that there must be a correspondence between the fundamental and full quantum spectrum of a  $4D$ ,  $N = 2$  MESGT and the minimal unitary representation of its three dimensional U-duality group and those representations obtained by tensoring of its minrep.
- ▶ Important problem: decomposition of tensor products of minreps into irreducible representations of the spectrum generating symmetry groups. In particular, how to obtain quaternionic series representations by tensoring minimal unitary representations.

# AdS/CFT: Aspen Summer 1984

- ▶ The Kaluza-Klein spectrum of IIB supergravity on  $AdS_5 \times S^5$  was first obtained via the oscillator method by simple tensoring of the CPT self-conjugate doubleton supermultiplet of  $N = 8$   $AdS_5$  superalgebra  $PSU(2, 2 | 4)$ .
- ▶ The CPT self-conjugate doubleton supermultiplet of  $PSU(2, 2 | 4)$  of  $AdS_5 \times S^5$  solution of IIB supergravity does not have a Poincaré limit in five dimensions and decouples from the Kaluza-Klein spectrum as gauge modes and the field theory of CPT self-conjugate doubleton supermultiplet of  $PSU(2, 2 | 4)$  lives on the boundary of  $AdS_5$ , which can be identified with 4D Minkowski space on which  $SO(4, 2)$  acts as a conformal group, and the unique candidate for this theory is the four dimensional  $N = 4$  super Yang-Mills theory that was known to be conformally invariant. MG, Marcus (1984)
- ▶ The spectra of 11D supergravity over  $AdS_4 \times S^7$  and  $AdS_7 \times S^4$  were fitted into supermultiplets of the symmetry superalgebras  $OSp(8 | 4, \mathbb{R})$  and  $OSp(8^* | 4)$  constructed by oscillator methods. The entire Kaluza-Klein spectra over these two spaces were obtained by tensoring the singleton and doubleton supermultiplets of  $OSp(8 | 4, \mathbb{R})$  and  $OSp(8^* | 4)$ , respectively.
- ▶ The relevant singleton supermultiplet of  $OSp(8 | 4, \mathbb{R})$  and doubleton supermultiplet of  $OSp(8^* | 4)$  do not have a Poincaré limit in four and seven dimensions, respectively, and decouple from the respective spectra as gauge modes. Again it was proposed that field theories of the singleton and scalar doubleton supermultiplets live on the boundaries of  $AdS_4$  and  $AdS_7$  as superconformally invariant theories. MG, Warner (1984), MG, Pvn, Warner (1984).
- ▶ Singletons of  $Sp(4, \mathbb{R})$  are the remarkable representations of Dirac (1963). Subsequent important work of Fronsdal and collaborators.

# Minimal Unitary Representations versus Singletons & Doubletons and Supersymmetry

MG, Pavlyk (2006), MG, Fernando (2009/10)

- ▶ The minimal unitary representations of symplectic groups  $Sp(2n, \mathbb{R})$  are the singletons and their generators can be written as bilinears of bosonic oscillators since their quartic invariants vanish. Tensoring procedure becomes simple for the symplectic groups. Minreps of  $OSp(2n|2m, \mathbb{R})$  are supersingletons
- ▶ Minimal unitary representation of  $SU(2, 2) = SO(4, 2)$  over  $L^2$  functions in 3 variables  $\Leftrightarrow$  conformal scalar = scalar doubleton
- ▶ Minrep of  $SU(2, 2)$  admits a one-parameter,  $\zeta$ , family of deformations corresponding to massless conformal fields in  $d = 4$  with helicity  $\zeta/2$ .
- ▶ Minrep of  $PSU(2, 2|4)$  is the 4D Yang-Mills supermultiplet (CPT-self-conjugate doubleton)
- ▶ Minrep of  $SU(2, 2|N)$  admits a one-parameter family of deformations  $\Leftrightarrow$  Higher spin doubleton supermultiplets studied in MG, Minic, Zagermann (1998).
- ▶ Minrep of  $SO^*(8) \simeq SO(6, 2)$  realized over the Hilbert space of functions of five variables and its deformations labeled by the spin  $t$  of an  $SU(2)$  subgroup correspond to massless conformal fields in six dimensions. Minimal unitary supermultiplet of  $OSp(8^*|2N)$  admits deformations labeled by the spin  $t$  of an  $SU(2)$  subgroup of the little group  $SO(4)$  of lightlike vectors in  $6D$ .
- ▶ Minrep of  $OSp(8^*|4)$  is the massless supermultiplet of  $(2, 0)$  conformal field theory that is dual to M-theory on  $AdS_7 \times S^4$ .

- ▶ To summarize: minreps and their deformations describe the *CFT* side of *AdS/CFT* dualities. For *AdS* supergroups the minreps act like "quarks" of positive energy unitary representations  $\rightarrow$  *AdS/CFT*
- ▶ Furthermore the above results imply that the minrep of any noncompact group  $G$  must admit deformations if it occurs as a factor in the even subgroup  $G \times K$  of a supergroup. ( For  $K$  compact such supergroups admit unitary representations in general.) Groups  $F_4, E_6, E_7$  and  $E_8$  do not occur as subgroups of any simple Lie supergroups !
- ▶ The mapping between the HSS formulation of  $N = 2$  sigma models coupled to sugra in  $d = 4$  and the minimal unitary realizations of their isometry group  $\implies$  suggest that the minimal unitary representations of  $3d$  U-duality must act as the "quarks" of the spectrum of these theories whose quantum completion may require extension to M/superstring theory.  
The corresponding sigma models reduced to one dimension give supersymmetric sigma models. Indeed the quantization of certain one-dimensional sigma models with 8 super symmetries yields the minimal unitary representation of  $D(2, 1; \alpha) \supset SU(1, 1) \times SU(2) \times SU(2)$  and its deformations.

Govil & MG (2012)

# OPEN PROBLEMS

- ▶ Decomposition of tensor products of minreps of U-duality groups into irreps. Of particular interest are  $E_{8(8)}$  and  $E_{8(-24)}$ .
- ▶ Physical meaning of the states of minrep of U-duality groups. For  $AdS$  supergroups the minreps act like "quarks" of positive energy unitary representations  $\rightarrow AdS/CFT$
- ▶ For 3D U-duality groups of MESGTs with 8 real supersymmetries the fundamental spectrum corresponds to "super BPS" states, i.e they preserve full susy!
- ▶ Explicit construction of the quaternionic discrete series for  $F_{4(4)}$ ,  $E_{6(2)}$ ,  $E_{7(-5)}$  and  $E_{8(-24)}$  and the corresponding discrete series for  $E_{8(8)}$
- ▶ Construction of the full spectrum by tensoring of minreps and embedding into their quantum completion ( M/Superstring theory).
- ▶ Supersymmetry of the black hole spectra & relation to the work of Townsend et.al. on the black holes and conformal quantum mechanics and Calogero models.
- ▶ At the non-perturbative level one expects only the discrete subgroups of U-duality groups to be symmetries of M/superstring theory compactified to various dimensions on tori or their orbifolds ( Hull and Townsend) . How to extend the above results to the discrete subgroups? Automorphic representations ?

HAPPY BIRTHDAY PAUL !