

# Evolution and magnetic topology of the M 1.0 flare of October 22, 2002

A. Berlicki<sup>1,2</sup>, B. Schmieder<sup>1,3</sup>, N. Vilmer<sup>1</sup>, G. Aulanier<sup>1</sup>, and G. Del Zanna<sup>4</sup>

<sup>1</sup> Observatoire de Paris, Section de Meudon, LESIA, F-92195 Meudon Principal Cedex, France

<sup>2</sup> Astronomical Institute of the Wrocław University, ul. Kopernika 11, 51-622 Wrocław, Poland

<sup>3</sup> ITA, Oslo, Norway

<sup>4</sup> DAMTP, University of Cambridge, Cambridge, UK

Received date / Accepted date

**Abstract.** In this paper we analyse an M1.0 confined flare observed mainly during its gradual phase. We used the data taken during a coordinated observational campaign between ground based instruments (THEMIS and VTT) and space observatories (SoHO/CDS and MDI, TRACE and RHESSI). We use these multi-wavelength observations to study the morphology and evolution of the flare, to analyse its gradual phase and to understand the role of various heating mechanisms. During the flare, RHESSI observed emission only within 3 – 25 keV spectral range. The RHESSI spectra indicate that the emission of the flare was mainly of thermal origin with a small non-thermal component observed between 10 and 20 keV. The temperature of plasma obtained from the fitting of the RHESSI X-ray spectra was between 8.5 and 14 MK. The lower temperature limit is typical for a plasma contained in post flare loops observed in X-rays. Higher temperatures were observed during a secondary peak of emission corresponding to a small impulsive event. The SoHO/CDS observations performed in EUV Fe XIX line also confirm the presence of a hot plasma at temperatures similar to those obtained from RHESSI spectra. Besides, the EUV structures were located at the same place as RHESSI X-ray emission. The magnetic topology analysis of the AR coming from a linear force-free field extrapolation explains the observed features of the gradual phase of the flare i.e. the asymmetry of the ribbons and their fast propagation. The combination of the multi-wavelength observations with the magnetic model further suggests that the onset of the flare would be due to the reconnection of an emerging flux in a sheared magnetic configuration.

**Key words.** Sun: flares – Sun: X-rays – Sun: magnetic fields

## 1. Introduction


The gradual phase of flares can last half an hour to few hours. Detailed observations of flares during their gradual phase have been recorded by Solar Maximum Mission (SMM) and later by Yohkoh (Švestka et al. 1982, Schmieder et al. 1996 and 1997, van Driel et al. 1997). Soft X-ray instruments in these missions observed the formation of hot loops (10–20 MK). These loops subsequently cool and become visible as post-flare loops in lower temperatures (EUV and H $\alpha$  lines). The hot loops are formed at higher and higher altitudes as the time is increasing. Such events were interpreted with models based on the Kopp and Pneuman (1976) concept. These models exist also in two dimensions (Forbes and Malherbe 1986, Forbes and Acton 1996). During the gradual phase of flares, magnetic field lines reconnect at an X point or neutral sheet in the corona. Particles are accelerated as

a consequence of the reconnection. Plasma in the loop is heated and "gentle" chromospheric evaporation can occur at the footpoints of the loops in high density plasma region. These models are well adapted for the interpretation of flare events in general, not only for eruptive two ribbon flares, but also for confined flares (Švestka 1986) that are due to magnetic reconnection at (quasi-) separatrix layers (Démoulin et al. 1997) and which do not result in an eruption.

During the reconnection process and the gradual phase of flares, SMM with its HXIS telescope could image the plasma heated by high energy particle in four energy bands of spectra with a low spatial resolution. The Hard X-ray telescope on-board Yohkoh had a better spatial resolution but no low energy band. A long debate about the long duration emission of particles started in order to understand if the particles accelerated during the impulsive phase were trapped at the reconnection point or if they were continuously accelerated.

---


Send offprint requests to: A. Berlicki,  
e-mail: [arkadiusz.berlicki@obspm.fr](mailto:arkadiusz.berlicki@obspm.fr)



**Fig. 1.** Image of AR 0162 obtained from THEMIS/MSDP observations taken between 15:41 and 16:29 UT in the Na I D<sub>1</sub> spectral line wing at 200 mÅ from the line centre (left panel). In the right panel we present the line-of-sight (LOS) magnetic field deduced from *I* and *V* Stokes profiles at 100 mÅ from the Na I D<sub>1</sub> line centre overlaid with the contours of the H $\alpha$  flare observed by VTT/MSDP at 15:30:01 UT. The field-of-view (FOV) is 400''  $\times$  250''. The cross denotes the place separating two main ribbons R1 and R2.

To distinguish between both processes and to further investigation this question we use the X-ray observations obtained with high spectral resolution in the 3–20 keV range by the Reuven Ramaty High Energy Solar Spectroscopic Imager (RHESSI: Lin et al. 2002). RHESSI was launched in 2002 and continuing to operate at the present time, are observing the solar spectrum in the energy range 3 keV – 17 MeV.

We had the opportunity to perform multi-wavelength observations a long duration M1.0 flare during a coordinated observational campaign between ground based instruments (THEMIS and VTT) and space observatories (SoHO/CDS and MDI, TRACE and RHESSI). The flare was followed by several local increases of X-ray emission in its decay phase. The flare occurs in a mainly bipolar active region (as observed by THEMIS). Looking at SoHO/EIT and LASCO data, we noticed that neither coronal mass ejection nor any related coronal signature of eruptive events (such as transient coronal dimming and fast propagating wave-like disturbance) were associated to the flare. Therefore, this event was certainly a confined flare. We observe mainly one long bright ribbon in H $\alpha$  and with the Transition Region and Corona Explorer (TRACE) at 195 Å. To understand the magnetic configuration of the flare we have performed magnetic extrapolations using the *lfff* (linear force-free field) code developed by Démoulin et al. (1997). The three-dimensional analysis of the extrapolation allows us to have a good idea of the topology of the flare. Long low sheared horizontal field lines were drawn, they could correspond to the TRACE loops. RHESSI and CDS Fe XIX images show elongated emissions over these field lines. This agrees with the models of gradual phase already explained above. The analysis of the RHESSI spectra allowed us to find if non-thermal particles can still exist during the gradual phase of the flares.




**Fig. 2.** Time evolution of GOES X-ray flux observed during the flare of October 22, 2002. The timings of the RHESSI observations (three orbits) are indicated. THEMIS observed continuously between 15:41–17:35 UT. TRACE and VTT/MSDP observations were performed during whole flare.

The paper is organized as follows. In Sect. 2 we describe the observations and their coalignment. In the third section we discuss on the topology of the flare using *lfff* extrapolation. Section 4 provides description of the RHESSI X-ray spectral observations of the flare. In this section we also discussed the nature of X-ray emission. Section 5 summarizes our analysis and provides a description of our results.

## 2. Observations

The M1.0 flare was observed in NOAA 0162 active region (AR) located about N26 E21 on October 22, 2002. This active region was a target during the coordinated SoHO and ground-based observational campaign in October 2002. The active region campaign included EUV spectral observations with the SoHO/CDS Normal Incidence Spectrometer (NIS) (Harrison et al. 1995). During its passage through solar disk (17–31 October, 2002) the active region consisted of a big leading spot located in a positive



**Fig. 3.** Approximate orientation of FOVs of different observations used in the analysis of the flare of October 22, 2002. The north is up. For RHESSI we marked the part of FOV which was used in our analysis.

magnetic field polarity and a cluster of following spots of negative magnetic field polarity (Fig. 1).

At X-ray range, this flare was observed by GOES 9 and in its decay phase by one of the RHESSI orbits (see Fig. 2). According to the GOES 9 flux curve the flare onset was at 15:29 UT and the  $1 - 8 \text{ \AA}$  X-ray flux reached its maximum at 15:36 UT. During the decay phase, several local increases of the X-ray flux were observed, the most notable one with a peak at 16:28 UT.


### 2.1. Summary of the data

Within the X-ray spectral range, the most interesting observations are those obtained by RHESSI orbit between 16:03 – 16:28 UT. During this time the X-ray emission was detected from 3 keV to about 25 keV.

The spectro-polarimetric observations of the AR 0162 were performed with the THEMIS telescope, operating in the MSDP (Multichannel Subtractive Double Pass spectrograph) observing mode (Mein 2002).

In our analysis we also used the VTT/MSDP observations in  $H\alpha$  line (Mein 1991). During the flare between 15:30 and 16:17 UT we completed 9 scans of the MSDP entrance window covering the whole active region with a field-of-view (FOV) of about  $380'' \times 170''$ .

TRACE (Handy et al. 1999) obtained high cadence (40 s) filtergrams in the  $195 \text{ \AA}$  band between 15:29 and 15:41 UT (with a FOV  $384'' \times 384''$  and a  $1''$  resolution). Before and after this time the TRACE observations were obtained with slightly lower time resolution (50 s) and larger FOV ( $512'' \times 512''$ ). The solar coordinates of the TRACE images were corrected by using the SoHO/EIT and MDI maps. In the EUV spectral range this flare was also observed by the SoHO/CDS instrument. In Fig. 3 we



**Fig. 4.** Time evolution of RHESSI X-ray counts rates observed during the flare of October 22, 2002 in 3–6, 6–12 and 12–25 keV energy bands. The counts rates observed in 12–25 keV energy band were multiplied by a factor of 10 to make the plots clear. Letters A, B and C denote the phases in the X-ray flux evolution (see Sect. 2.2).

present the relative pointings and approximate fields-of-view of the various instruments used in our analysis.

### 2.2. Magnetic field observation

On October 22, 2002 the spectro-polarimetric observations of the AR 0162 were performed with the THEMIS telescope, operating in the MSDP spectrograph observing mode. THEMIS is a polarization free telescope, designed especially for polarimetric observations. The polarization analyser provides successively the Stokes parameters  $I \pm S$  ( $S = V, Q$  or  $U$ ) without differential blurring due to seeing effects. Detailed descriptions of the specific instrumental set up of THEMIS in the MSDP observing mode can be found in Mein (2002).

During the flare the observations in Na I  $D_1$  sodium line were performed only for  $S = V$  and we obtained the Stokes  $I$  and  $V$  profiles simultaneously in the 16 MSDP channels. The line-of-sight (LOS) magnetic field can be estimated from the observed Stokes profiles. The center-of-gravity method (Semel 1967) was used to determine the magnetic field. In this method the longitudinal (or LOS) component  $B_l$  of magnetic field is proportional to the relative shift between the centres of  $I + V$  and  $I - V$  profiles (for details see the paper Eibe et al., 2002).

On October 22, 2002 we have obtained five scans of the AR 0162 during the flare. Each scan lasted about 25 minutes. From these scans five sets of data were reconstructed. Each set of data consists of the images ( $600'' \times 130''$ ) in the Na  $D_1$  sodium line (intensity, velocity and LOS magnetic field maps at different wavelengths). The FOV of the THEMIS scans was too small to cover the whole active re-

gion and to make an image of the whole AR thus we need to make two scans with two different FOV.

On October 22, 2002 THEMIS observations of the AR 0162 were performed from about 10:18 until 18:00 UT. Analysis of all images showed that there were no significant changes of LOS magnetic field configuration during all day. Finally, during the flare from five scans we obtained two full images ( $400'' \times 250''$ ) of the AR at 15:41–16:29 and 16:47–17:35 UT. The pixel size of these images is  $0.25''$  but the spatial resolution of the images was limited by seeing (about  $1''$ ).

Fig. 1 presents THEMIS observations of AR 0162 obtained in the Na I D<sub>1</sub> spectral line wing at  $200 \text{ m}\text{\AA}$  from the line centre (left panel). In the right panel we present the LOS magnetic field deduced from *I* and *V* Stokes profiles of the Na I D<sub>1</sub> line overlaid with the isophotes of the H $\alpha$  flare observed by VTT/MSDP. The area of high H $\alpha$  emission (more concentrated isophotes) is located close to the emerging bipole – a two polarity emerging flux of high gradient magnetic field (see Sect. 3).

### 2.3. X-ray observations

The decay phase of this flare was observed by RHESSI during the orbit between 16:03 – 16:28 UT (Fig. 4). From RHESSI data, it is possible to construct images in some chosen energy bands, and to build low-resolution spectra. During the decay phase the count rates detected from 3 keV to 25 keV were sufficiently high to make images with spatial resolution of  $4''$  but it was necessary to accumulate count rates over few minutes periods to reconstruct the images with valuable quality. We obtained the images in three energy bands 3–6 keV, 6–12 keV and 12–25 keV using the CLEAN algorithm and front collimators 3F–9F. During this flare there was no attenuator in front of the grids. We concentrated the analysis of the RHESSI data on three periods of time (ranges A,B,C in Fig. 4). During the period A (16:03–16:13 UT), the X-ray count rates exhibited a continuous decrease. At 16:16 UT a weak increase of the count rates in 6–12 and 12–25 keV bands, lasting about 2 minutes, was observed (range B). This small event was not visible on the RHESSI flux curves for higher energies. After it, from about 16:18 UT we can see a continuous increase of the X-ray flux (range C). It reached its maximum at about 16:26 UT, just before the passage of RHESSI through the radiative belts.

Between 16:03 and 16:09 UT during the decaying X-ray flux (range A) we could reconstruct the images only in the 3–6 and 6–12 keV energy bands (Fig. 5 – left panel). Images shows an elongated loop-like structures with the same shape both in 3–6 keV as well as in 6–12 keV range. We have also examined the small event (period B), by reconstructing RHESSI images during its peak (16:15–16:19 UT) in the 3–6 and 6–12 keV energy bands (Fig. 5 – middle panel). The X-ray emission was concentrated in two compact sources located slightly to the south of the emission observed at 16:03 UT (Fig. 6 – left panel). In the

12–25 keV energy band the count rates were too low to make images but analysis of the spectra can be done in this range.


The following increase (period C) was well visible in the 3–6, 6–12 and 12–25 keV energy bands, and peaked at about 16:26 UT. We have reconstructed images around this peak (16:25–16:28 UT), shown in Fig. 5 – right panel and Fig. 6. Also in this case elongated loop-like structures are seen, that, however, are now located to the south of the structures previously observed at 16:15–16:19 UT. There were no significant shifts between the positions of these X-ray structures as observed at different energies (Fig. 6 – right panel).

### 2.4. Coronal Diagnostic Spectrometer (CDS)


By rastering the CDS slit (from west to east), monochromatic images and velocity maps of the FOV ( $4' \times 4'$ ) can be constructed. The NIS observes the 305–635 Å spectral range with two gratings, and a spatial resolution of approximately  $5''$ . The NIS spectral range includes a large number of lines, but due to telemetry limitations, only a limited number of NIS lines are normally extracted and telemetered to the ground. For more details of the CDS see e.g. Harrison et al. (1995) and for data reduction Del Zanna et al. (2001).

Two types of NIS ‘rasters’ were used for these observations: 1) a slow (1hour 20 mins), diagnostic raster that extracts many diagnostic lines, 2) a fast (20 mins) raster, that only extracts a few lines, selected to cover a wide range of temperatures (He I 522.2 Å,  $\log T = 4.5$ ; O IV 554.5 Å,  $\log T = 5.3$ ; O V 630 Å,  $\log T = 5.4$ ; Mg X 625 Å,  $\log T = 6.0$ ; Si XII 520.8 Å,  $\log T = 6.3$ ; Fe XIX 592.2 Å,  $\log T = 6.9$ ). The two rasters used the  $4'' \times 240''$  slit and were repeated a few times over the active region during the day. In this paper, only some results concerning these two CDS rasters are presented. A more detailed discussion of the CDS observations is deferred to a future paper.

NOAA 0162 was observed by CDS during the periods 08:02–13:07 UT and 15:18–18:33 UT with the two types of rasters. A relatively strong flare was observed during the fast raster taken during 12:43–13:07 UT, approximately at the same location where the M1.0 flare took place. The M1.0 flare that is the subject of this paper was recorded by CDS with a fast raster which scanned the active region between 15:18 and 15:43 UT. CDS also observed the same region with a diagnostic raster (15:43–17:07 UT). In our analysis we used the CDS images obtained in Fe XIX line at 592.2 Å. The CDS slit was located in the flare area during the peak X-ray emission. It should be noted that spatially-resolved EUV spectroscopic observations of flares are quite rare, although CDS has already observed a number of them, as described in Del Zanna et al. (2002a) and Del Zanna et al. (2002b).



**Fig. 5.** Sequence of RHESSI images ( $224'' \times 224''$ ) reconstructed in 6–12 keV energy band during the gradual phase of the flare on October 22, 2002.



**Fig. 6.** Left panel: The time evolution of RHESSI X-ray structures observed in 6–12 keV energy band. RHESSI image ( $224'' \times 224''$ ) taken at 16:03–16:09 UT is overlaid with the X-ray contours at 16:15–16:19 and 16:25–16:28 UT. Right panel: RHESSI images between 16:25 and 16:28 UT observed in three different (3–6, 6–12 and 12–25 keV) energy band. The image shows the X-ray structures observed in 3–6 keV energy bands overlaid with the contours of the structures observed at this time in 6–12 and 12–25 keV energy bands.

### 2.5. Multi-wavelength observations of the flare structure


On the  $H\alpha$  VTT/MSDP images well developed flare ribbons were observed from 15:30 UT. The orientation of the two main ribbons R1 and R2 indicates the presence of a large shear in this AR (Fig. 7). The brightest parts of  $H\alpha$  flare ribbons, well visible at 15:30 UT, were much weaker after 7 minutes and almost disappeared at about 15:59 UT but the other parts of the flare ribbons were still well visible. The two brightest parts of  $H\alpha$  ribbons were located in the following spots group, within emerging bipole, each of them in an area of opposite polarity magnetic field (Fig. 1 – right panel).

We point out that the two-ribbon shape of this flare is not contradictory with its confined nature, since flare models (Démoulin et al. 1997) have shown that quasi-bipolar photospheric flux distributions can be associated


with quasi-separatrix layers whose intersection with the photosphere can form only two (and not four) thin ribbons.

The first increase of emission in the 195 Å band was observed by TRACE on the image taken at 15:32:18 UT while the first increase of GOES X-ray flux of M1.0 flare was observed at 15:29 UT. Initially, the TRACE emission was concentrated in the compact source well visible at 15:34:56 and 15:40:39 UT (Fig. 8). Later on, from about 15:58:55 UT the TRACE UV emission was concentrated in small, rather low, thin loops spreaded along the east-west axis of the AR and in few brightenings located close to the  $H\alpha$  flare emission (Fig. 9 – lower panels).

In the part of the AR where the magnetic field gradient was very high the initial energy release probably took place. This idea may be supported by the TRACE 195 Å observations. In the image taken at 15:34:56 UT the



**Fig. 7.** The flare of October 22, 2002 observed in  $H\alpha$  line by MSDP/VTT spectrograph. The FOV is  $340'' \times 200''$ . R1 and R2 denote the two main ribbons of the flare. LS marks the leading spot of the AR 0162.




**Fig. 8.** TRACE 195 Å image of the AR 0162 at 15:34:56 UT. R1 and R2 denote the two main ribbons of the flare and LS marks the leading spot of the AR 0162 (left panel). In the right panel we present the contours of longitudinal magnetic field of the AR 0162 observed with THEMIS in the Na I  $D_1$  spectral line at 15:41–16:29 UT. Thick and thin contours denote respectively the positive and negative polarity magnetic field.

brightest UV emission was concentrated just above the emerging bipole (Fig. 8 – lower right panel) and above two bright  $H\alpha$  flare kernels observed at 15:30:01 UT.

The coalignment of RHESSI images with the VTT/MSDP filtergrams shows that X-ray emission observed between 16:03 and 16:09 UT is located not on the  $H\alpha$  flare ribbons but, in projection, it is shifted above (Fig. 9 – upper left panel). There is no shift between the emission observed in 3–6 and 6–12 keV energy bands. Later, the X-ray structures observed by RHESSI at 16:25 – 16:28 UT were located (in projection) in the same place as  $H\alpha$  flare emission observed half an hour before. In the upper right panel of Fig. 9 we present also the contours of RHESSI X-ray structures observed in three time intervals between 16:03 and 16:25 UT overlaid on THEMIS/MSDP image of LOS magnetic field. In the lower left panel of

Fig. 9, the contours of RHESSI image at 16:03–16:09 UT are overlaid on the TRACE image taken at 16:04:38 UT. Note that X-ray emission of the flare is located just above the bright thin loop-like structure seen in UV 195 Å. Later, the X-ray emission observed by RHESSI at 16:25–16:28 UT was located (in projection) in the same place as TRACE emission observed at 16:27:04 UT (Fig. 9 - lower right panel).

During the flare, the Fe XIX line observed by CDS was very bright (reaching 3000 photon-events in the NIS detector), and clearly showed where strong heating (to at least temperatures of 8 MK) takes place. The peak emission, recorded between 15:35 and 15:43 UT, occurred just above the region of strongly mixed polarity and where the  $H\alpha$  flare ribbons observed by VTT/MSDP at 15:30:01 UT were located (Fig. 10 – lower left panel). The Fe XIX emis-



**Fig. 9.** VTT/MSDP image of the H $\alpha$  flare in the AR 0162 at 15:59:52 UT (left upper panel) and the longitudinal magnetic field deduced from THEMIS/MSDP observations (15:41–16:29 UT) in the Na I D<sub>1</sub> spectral line at 100 mÅ from the line centre (right upper panel) overlaid with the contours of the X-ray structures observed by RHESSI at 16:03–16:09, 16:15–16:19 and 16:25–16:28 UT (right upper panel). In the lower panels we present TRACE 195 Å images of the AR 0162 (16:04:38 and 16:27:04 UT) overlaid with the contours of the X-ray structures observed by RHESSI at 16:03–16:09 and 16:25–16:28 UT time intervals.

sion was located also just above the brightest TRACE 195 Å emission (Fig. 10 – upper left panel). Note that the TRACE 195 Å band is dominated at the flare site by Fe XXIV emission at 192.028 Å, while in general is multi-thermal, with contributions from plasma at temperatures as low as 700 000 K (see Del Zanna & Mason, 2003).

During the decay phase, the Fe XIX emission becomes more diffuse, a common feature already noted in Del Zanna et al. (2002a, 2002b) and shown in Fig. 10 (right panels). The Fe XIX emission was recorded while the slit was covering the eastern half of the FOV, between 16:20 and 16:51 UT, i.e. the RHESSI 16:25–16:28 UT image and the CDS scan are almost simultaneous. Note the good agreement between positions of the structures observed by CDS and RHESSI.


### 3. Interpretation of the flare ribbons in a simple topology

In this section we provide an interpretation for the triggering and the development of the confined flare by the means of a topological analysis of the coronal magnetic field above NOAA 0162 calculated from a linear force-free field model.

#### 3.1. Linear force-free field extrapolation

For the extrapolation the LOS magnetic fields was measured with the THEMIS/MSDP in the Na I D<sub>1</sub> line at 12:32–13:20 UT. We used the image obtained before the flare because the FOV was extended towards east and additional magnetic structures could be taken into account. The LOS magnetic fields were assumed to be equal to the vertical fields at the altitude  $z = 0$  (i.e.  $\mathbf{B}(x, y, z = 0)$ ) divided by the projection angle, as often done in extrap-






**Fig. 10.** TRACE 195 Å images of the AR 0162 (15:34:56 and 16:14:45 UT) overlaid with the contours of the UV structures observed by CDS between 15:18:16–15:43:16 and 15:43:30–17:07:33 UT in the Fe XIX line (592.2 Å) (upper panels) and VTT/MSDP image of the H $\alpha$  flare in the AR 0162 at 15:30:01 UT overlaid with the contours of the UV structures observed by CDS (Fe XIX line, 592.2 Å) between 15:18:16–15:43:16 UT (left lower panel) and RHESSI image (245''  $\times$  245'') reconstructed for the time 16:25–16:28 UT overlaid with CDS Fe XIX structures observed between 15:43:30–17:07:33 UT (right lower panel).

ulations using longitudinal magnetograms only (see e.g. Démoulin et al. 1996, Eibe et al. 2002). The extrapolation was performed in the linear force-free field approximation ( $\nabla \times \mathbf{B} = \alpha \mathbf{B}$  with  $\alpha = \text{cst}$ ) with the FFT method, so with periodic boundary conditions in  $(x, y)$  (Alissandrakis 1981). In order to minimize the boundary effects, a large computational box was chosen, with  $\Delta x = \Delta y = 10^3$  Mm, and the magnetic field was computed up to  $z = 150$  Mm above the photosphere. The calculation was done with  $nx = ny = 1024$  mesh points so as to sufficiently resolve the emerging bipole in the trailing part of the active region. The force-free parameter  $\alpha$  was iteratively tuned so as to obtain a general fit between the orientations of the flare loops observed by TRACE and of the calculated field

lines rooted in the vicinity of these loops. This constraint led to fix  $\alpha = -5 \times 10^{-3} \text{ Mm}^{-1}$ , whose sign is compatible with the hemispheric helicity rules found by Pevtsov et al. (1995).

Even though the results of this calculation were used to study the triggering and the development of the flare, we must point out that they are uncertain for two reasons. Firstly, the selected  $\alpha$  is equal to 80% of the resonant value well known for extrapolations performed with a FFT in a large computational box. This leads to a very probable artificial over-shearing of the large scale loops. Secondly, it was impossible to find a single value of  $\alpha$  with, even in a 20% interval, which allowed to fit all the observed TRACE loops with calculated magnetic field lines. Therefore, it ap-





**Fig. 11.** (*top left:*) superposition of the THEMIS/MSDP Na D1 magnetogram of 12:32:02 UT with a sample of magnetic field lines resulting from the *lfff* extrapolation. (*bottom right:*) partial view of the flare as observed in  $H\alpha$  with the VTT/MSDP at 15:59:52 UT. The field lines are the ones which are rooted in the flare ribbons R1 and R2 as well as in the emerging bipole. (*bottom left:*) same as *bottom right*, except that the underlying image is the THEMIS magnetogram. (*top right:* projection view of the field lines shown in the *bottom* panels. Continuous (resp. dashed) contours stand for positive (resp. negative) vertical fields. North is toward the left.


pears that the  $\alpha = \text{cst}$  hypothesis of the *lfff* extrapolation may be strongly disputed in this region. Nevertheless, as the global magnetic topology is a well preserved property of magnetic fields since it is an integral quantity, we still used the *lfff* model to pursue our analysis, although it is clear that it is disputable on small scales (where electric currents are concentrated) and on large scales (where the field is too much sheared).

### 3.2. Origins of the asymmetrical ribbons

Fig. 11 shows the results of the *lfff* extrapolation. The large scale shear is evident on the scale of the full active region, as seen on the *top left* panel. The other panels show the

connectivities of the magnetic field lines in the vicinity of the emerging bipole and around the flare ribbons as observed in  $H\alpha$ .

In this region, even though a few nulls points are present around the negative moving magnetic features on the edges of the leading spot, the global topology is *not* complex: no thin quasi-separatrix layers were found to be associated with the flare loops, contrary to many previous studies of confined flares (see e.g. Mandrini et al. 1996, Schmieder et al. 1997, Aulanier et al. 2000). Also, the connectivities look very continuous around the region of the emerging bipole. So at first sight it appears that the flare studied in this paper cannot be interpreted with the standard reconnection model. However emerging bipoles are




**Fig. 12.** RHESSI spectra reconstructed during the gradual phase of the solar flare in four intervals: 16:03–16:04, 16:13–16:14, 16:16–16:17 and 16:25–16:26 UT. Dotted lines denote the calculated thermal component, dashed lines represent non-thermal component, thin continuous lines are the Fe-line feature ( $\sim 6.7$  keV) calculated in our model. In our modelling of the spectrum at 16:25–16:26 UT we also used Fe/Ni-line feature ( $\sim 7.9$  keV). Thick continuous lines are the sum of all components – predicted photon spectrum. With crosses we marked observed flare spectrum. The temperature of plasma  $T$ , emission measure  $EM$ , power-law index of non-thermal component  $\gamma$ , and normalization of broken power-law  $A$  at 10 keV for these spectra are: 16:03–16:04 UT:  $T_1 = 9.8$  MK,  $EM_1 = 2.3 \times 10^{48} \text{ cm}^{-3}$ ,  $\gamma_1 = -7.5$ ,  $A_1 = 3.3 \text{ photons s}^{-1} \text{ cm}^{-2} \text{ keV}^{-1}$ ; 16:13–16:14 UT:  $T_2 = 9.1$  MK,  $EM_2 = 1.5 \times 10^{48} \text{ cm}^{-3}$ ,  $\gamma_2 = -6.9$ ,  $A_2 = 1.1 \text{ photons s}^{-1} \text{ cm}^{-2} \text{ keV}^{-1}$ ; 16:16–16:17 UT:  $T_3 = 9.4$  MK,  $EM_3 = 9.4 \times 10^{47} \text{ cm}^{-3}$ ,  $\gamma_3 = -6.7$ ,  $A_3 = 2.1 \text{ photons s}^{-1} \text{ cm}^{-2} \text{ keV}^{-1}$ ; 16:25–16:26 UT:  $T_4 = 13.7$  MK,  $EM_4 = 2.4 \times 10^{47} \text{ cm}^{-3}$ ,  $\gamma_4 = -7.0$ ,  $A_4 = 4.1 \text{ photons s}^{-1} \text{ cm}^{-2} \text{ keV}^{-1}$ .

known to often emerge twisted (Leka et al. 1996, Lopez-Fuentes et al. 2000), which is supported by the Huairou vector magnetograms for this region (Y.H. Yan, private communication). Therefore we conjecture that the flare was still probably due to magnetic reconnection, but here between small-scale twisted emerging fields (not calculated in the extrapolation) and the large-scale sheared magnetic fields of the global bipole of NOAA 0162. Since the flare was confined, it may then be interpreted by the global disruption model calculated by Amari et al. (1999) in a very symmetrical configuration, where a lo-

cally twisted flux tube undergoes an ideal kink instability and reconnects with overlaying potential fields.

If we assume the global disruption scenario, the *lfff* model naturally provides an explanation for the appearance of the compact ribbon R1 and its elongated counterpart R2, as well as its fast propagation away from the emerging bipole and toward the leading sunspot. Figure 11 indeed shows that the field lines located higher and higher above the emerging bipole are rooted in R1 and R2, and have footpoint separations that are small on the R1 side and quite large on the R2 side. The continuous reconnect-



**Fig. 13.** The time evolution of the temperature  $T$  and emission measure  $EM$  of the flare on October 22, 2002. These parameters were obtained from fitting the observed RHESSI spectra with the model.

tion of the growing emerging fields with higher and higher overlaying fields provides a natural explanation for the sequential brightening of their footpoints along the flare ribbons, included along the very elongated R2 ribbon. In this frame, the fast propagation of the brightenings as observed by TRACE are not due to MHD waves, but are rather due to the electron beams accelerated from the reconnection regions above the emerging bipole in the long overlaying loops of NOAA 0162, which progressively reach larger and larger chromospheric areas toward the West.

It is also noteworthy that the shape of the field lines anchored in the  $H\alpha$  ribbons follows the elongated shape of the RHESSI images recorded in the 6–12 keV range of energy between 16:03–16:09 UT. Thus, the extrapolation supports the idea that RHESSI observes thermal emission of coronal loops heated by the confined flare. It is, however, not contradictory that the RHESSI also detects a non-thermal component since it is known that reconnection can accelerate particles along reconnected field lines. The RHESSI emission at 16:25–16:28 UT being similar in shape to the one recorded at 16:03–16:09 UT, though slightly shifted southward, suggests that this secondary event may be associated to the resistive relaxation of the system which may have been forced too rapidly by the flare of 15:59 UT, as in the reconnection calculations of Karpen et al. (1998). Unfortunately, the present data and model cannot provide more clues to this issue.

#### 4. Thermal and non-thermal component in the gradual phase

Using RHESSI data and the SPEX package we performed spectral analysis of X-ray emission from 16:03 to 16:27 UT. The background emission was evaluated using data taken before the flare during the previous RHESSI orbit between 15:12 and 15:24 UT. Spectral accumulation

was done for all 1 minute time bins between 16:03 and 16:27 UT and was performed using 0.25 keV energy bins in the 3–10 keV energy range and 1 keV energy bins above 10 keV. To obtain the spectra we used front segments of detectors 1F–9F (except 2F and 7F.) We omitted the data from these detectors because they have a poorer energy resolution than intended (Smith et al. 2002). We performed the full calculation of all diagonal and off-diagonal matrix elements.

All the spectra accumulated between 16:03 and 16:27 UT were analysed with SPEX. To reproduce the observed count spectra, we used a model consisting of a thermal component (characterized by an emission measure and a temperature) and a power-law non-thermal component. In addition to these continuum components we considered two lines that Phillips (2004) predicted to be significant in X-ray spectra observed by RHESSI for hot ( $T_e > 10$  MK) plasma. The first line is a Fe-line feature at  $\sim 6.7$  keV and the second line is a Fe/Ni complex line feature at  $\sim 8$  keV emitted from even hotter plasma. The 6.7 keV feature corresponds mainly to the Fe XXV lines and associated dielectronic satellites, centred on 6.6 keV and to Fe XXVI lines and associated satellites, centred on 6.9 keV. The 8 keV line feature is due to Fe XXV and Fe XXVI lines as well as Ni XXVII and Ni XXVIII lines. The lines are excited at electron  $T_e > 10$  MK. The contributions of individual lines to the total emission within both line features is temperature dependent (Phillips 2004). The Fe/Ni-line feature starts to be visible with RHESSI at higher temperatures than Fe-line feature, and for the  $T_e < 20$  MK the Fe/Ni-line feature is very weak.

Fig. 12 shows the photon spectrum resulting from the fitting of the model to the observed count spectrum within the 3–20 keV energy range. For all spectra a non-thermal continuum is necessary to reproduce the observation above 10 keV. Furthermore, for all spectra the Fe-line feature at

$\sim 6.7$  keV is seen in the count and photon spectra. For the spectra after 16:25 UT, the complex Fe/Ni-line feature at  $\sim 8$  keV gives a marginally significant contribution to the observed spectrum in good consistency with the higher temperatures deduced from the thermal component. The main difference in the spectra obtained between 16:03 and 16:27 UT is the increase of the temperature of the thermal component at 16:26 UT as well as the contribution of the non-thermal component to the spectra above 10 keV. The contribution of the non-thermal component (see Fig. 12) is larger in the 16:16 – 16:17 UT and 16:25 – 16:26 UT, during time intervals corresponding to some additional increases of the X-ray flux in 12 – 25 keV energy range. In the figures the fitted spectra are shown assuming no low energy cut-off for the non-thermal part of the photon spectrum. The effect of a low energy cut-off around 10 keV for the non-thermal photon spectrum were analysed. Below the cut-off we assumed a photon spectrum with an index of  $-1.5$ . We found that below 10 keV the fit is not strongly dependent on the existence and the value of this cut-off which means that the thermal emission is dominant below 10 keV. For all spectra and independent on the cut-off value the contribution of a non-thermal component is necessary to explain the photon spectrum between 10 and 20 keV.

Fig. 13 shows the time evolution of the temperature  $T$  and emission measure  $EM$  obtained by the fitting of the observed count spectra with the model. First, the deduced temperature decreases from about 9.7 MK to 8.8 MK and after, at 16:10 UT starts to increase to the value of 13.6 MK. It means that from about 16:10 UT some additional heating processes took place. The maximum value of the temperature at 16:25 UT is consistent with the marginal appearance of the Fe/Ni-line feature at the same time. Between 16:03 and 16:09 UT the emission measure exhibited some increase. At about 16:09 UT  $EM$  started to decrease from  $2.53 \times 10^{48} \text{ cm}^{-3}$  to  $2.35 \times 10^{47} \text{ cm}^{-3}$  at 16:27 UT what indicate for continuous decrease of amount of emitting plasma.

## 5. Conclusions

In this paper we performed an analysis of the morphology and evolution of an M1.0 flare on October 22, 2002 observed mainly during its gradual phase. We used multi-wavelength observations obtained in optical range as well as in X-ray and EUV. We also analysed line-of-sight magnetic field of the AR 0162 which allowed us to understand the mechanisms which are responsible for the observed evolution of this event. The magnetic field (photospheric and extrapolated) and the aligned  $H\alpha$  ribbons shows that this active region was highly sheared, thus was storing energy. The triggering of  $H\alpha$  flare occurs in a region of high gradient of  $B$  which could correspond to the emergence of a locally very twisted magnetic flux. The *loff* extrapolation of the magnetic field provides an explanation for the appearance two  $H\alpha$  flare ribbons: ribbon R1 was compact and its position was almost constant during the

flare; ribbon R2 was elongated and exhibited fast propagation along it, from the emerging bipole towards the leading sunspot. This propagation can be explained by the reconnection of the growing emerging field lines with higher and higher overlaying field. The shift of RHESSI X-ray and CDS/EUV emission observed with time above the AR also confirms the idea of subsequential reconnection of magnetic field lines. The elongated shape of X-ray and EUV structures observed during the gradual phase follow the shape of the magnetic field lines anchored in the  $H\alpha$  ribbons. Thus, we can say that RHESSI and CDS permit to see thermal emission of successive coronal loops heated probably by non-thermal electrons, accelerated by reconnection processes.

The presence of non-thermal component during the gradual phase of flares is still not clear. It is well known that non-thermal particles occur during the impulsive phase of flares and they can still exist during late phase of an eruptive flares. Analysis of magnetic field configuration showed that the flare of October 22, 2002 was of confined type. During this event we did not observe neither CME nor any other signatures of an eruption. Nevertheless, the presence of non-thermal particles can be deduced from the X-ray spectra reconstructed during the gradual phase. RHESSI spectra show that the non-thermal component is necessary to explain the spectra between the energy 10 and 20 keV but the emission below 10 keV observed during gradual phase was mainly of thermal origin.

From the RHESSI spectra we found a temperature of the flare plasma to be around 8.5–14 MK. That is typical for the structures observed during the gradual phase of solar flares. The Fe-line feature ( $\sim 6.7$  keV) also indicates the presence of high temperature plasma (about 9–10 MK) (Phillips 2004). At 16:26 UT, during the increase of X-ray flux, we also observed Fe/Ni-line feature ( $\sim 7.9$  keV). The presence of this line emission is due to temperature increase to about 14 MK. The CDS observations in Fe XIX flare line (592.2 Å) confirm the existence of a hot plasma of that temperature. The Fe XIX emission arises at the temperature around 8–9 MK which is in agreement with the temperatures obtained from the RHESSI spectra. Besides, the Fe XIX emission was nearly co-spatial with the RHESSI X-ray emission.

The high plasma temperatures observed during the gradual phase of this flare are in agreement with the commonly-accepted model for gradual phase of flares (Forbes and Malherbe, 1986, Forbes and Acton, 1996). During the decay phase, X-ray emission was observed at the reconnection site of the magnetic field lines. The X-ray and EUV observations suggest that this emission originated rather from continuous reconnection processes than from trapped particles. There was no X-ray emission at the footpoints of the loops which indicate that during the gradual phase non-thermal heating of  $H\alpha$  flare ribbons was not present. The thermal conduction heating can be responsible for the flare ribbons observed by TRACE but is not sufficient to explain the  $H\alpha$  flare emission (Berlicki

and Heinzel 2004). We need to investigate other processes (for instance the effect of high coronal pressure).

*Acknowledgements.* The magnetic field extrapolations used in this paper have been obtained from the code base FRENCH Online MAGnetic Extrapolations (FROMAGE). FROMAGE is a joint project between LESIA (Observatoire de Paris), CPhT (Ecole Polytechnique) and the Centre National d'Etudes Spatiales (CNES). The work of A.B. was supported by the European Commission through the RTN programme (European Solar Magnetism Network, contract HPRN-CT-2002-00313). VTT/MSDP observations have been provided and reduced by P. N. Mein and J. Staiger. These observations were performed during a MEDOC campaign (JOP 157). GDZ acknowledges support from PPARC (UK). The TRACE analysis at SAO is supported by a contract from Lockheed Martin. The authors are grateful to the RHESSI Team (PI: R. P. Lin) for the free access to the RHESSI data and the development of the software. N.V. would like to acknowledge many useful discussion on RHESSI imaging techniques and spectroscopy with R. P. Lin, G. Hurford, S. Krucker and R. S. Schwartz. The support of CNES to the RHESSI project is also acknowledged.

## References

- Alissandrakis, C. E. 1981 *A&A*, 100, 197
- Amari, T., & Luciani, J. F. 1999, *ApJ*, 515, 81L
- Aulanier, G., DeLuca, E. E., Antiochos, S. K., McMullen, R. A., & Golub, L. 2000, *ApJ*, 540, 1126
- Berlicki, A., & Heinzel, P. 2004, *A&A*, in press
- Del Zanna, G., Bromage B. J. I., E. Landi, E., & Landini, M. 2001, *A&A*, 379, 708
- Del Zanna, G., Mason, H. E., Gibson, S. E., Pike, C. D., & Mandrini, C. H. 2002a, *Adv. Space Res.*, Vol. 30, No. 3, 551
- Del Zanna, G., Mason, H. E., & Foley, C. 2002b, *Proc. 10th European Sol. Phys. Meeting*, ESA-SP 506, 585
- Del Zanna, G., & Mason, H. E. 2003, *A&A*, 406, 1089
- Démoulin, P., Bagala, L. G., Mandrini, C. H., Henoux, J. C., & Rovira, M. G. 1997, *A&A*, 325, 305
- Eibe, M. T., Aulanier, G., Faurobert, M., Mein, P., & Malherbe, J. M. 2002, *A&A*, 381, 290
- Forbes, T. G., & Acton, L. W. 1996, *ApJ*, 459, 330
- Forbes, T. G., & Malherbe, J. M. 1986, *ApJ*, 302, L67
- Handy, B. N., et al. 1999, *Sol. Phys.*, 187, 229
- Harrison, R. A., et al. 1995, *Sol. Phys.*, 162, 233.
- Karpen, J. T., Antiochos, S. K., Devore, C. R., & Golub, L. 1998, *ApJ*, 495, 491
- Kopp, R. A., & Pneuman, G. W. 1976, *Sol. Phys.*, 93, 351
- Leka, K. D., Canfield, R. C., McClymont, A. N., & van Driel-Gesztelyi, L. 1996, *ApJ*, 462, 547
- Lin, R. P. 2002, *Sol. Phys.*, 210, 3
- Lopez Fuentes, M. C., Démoulin, P., Mandrini, C. H., & van Driel-Gesztelyi, L. 2000, *ApJ*, 544, 540
- Mandrini, C. H., Démoulin, P., van Driel-Gesztelyi, L., Schmieder, B., Cauzzi, G., & Hofmann, A. 1996, *Sol. Phys.*, 168, 115
- Mein, P. 1991, *A&A*, 248, 669
- Mein, P. 2002, *A&A*, 381, 271
- Pevtsov, A. A., Canfield, R. C., & Metcalf, T. R. 1995, *ApJ*, 440, L109
- Phillips, K. J. H. 2004, *ApJ*(submitted)
- Schmieder, B., Heinzel, P., Van Driel-Gesztelyi, L., & Lemen, J. R. 1996, *Sol. Phys.*, 165, 303
- Schmieder, B., Aulanier, G., Démoulin, P., van Driel-Gesztelyi, L., Roudier, T., Nitta, N., & Cauzzi, G. 1997, *A&A*, 325, 1213
- Semel, M. D. 1967, *Ann. Astroph.*, 381, 271
- Smith, D. M., Lin, R. P., Turin, P., et al. 2002, *Sol. Phys.*, 210, 33
- Švestka, Z., Hoyng, P., van Tend, W., et al. 1982, *Sol. Phys.*, 75, 305
- Švestka, Z. 1986, in *The Lower Atmosphere in Solar Flares*, NSO, 332
- van Driel-Gesztelyi, L., Wiik, J. E., Schmieder, B., Tarbell, T., Kitai, R., Funakoshi, Y., & Anwar, B. 1997, *Sol. Phys.*, 174, 151