

Mathematical Tripos Part II: Michaelmas Term 2015

Numerical Analysis – Lecture 2

Problem 1.8 Finite-difference discretization of $\nabla^2 u = f$ replaces the PDE by a large system of linear equations. In the sequel we pay special attention to the *five-point formula*, which results in the approximation

$$h^2 \nabla^2 u(x, y) \approx u(x - h, y) + u(x + h, y) + u(x, y - h) + u(x, y + h) - 4u(x, y). \quad (1.5)$$

For the sake of simplicity, we restrict our attention to the important case of Ω being a *unit square*, where $h = \frac{1}{m+1}$ for some positive integer m . Thus, we estimate the m^2 unknown function values $u(ih, jh)_{i,j=1}^m$ (where $(ih, jh) \in \Omega$) by letting the right-hand side of (1.5) equal $h^2 f(ih, jh)$ at each value of i and j . This yields an $n \times n$ system of linear equations with $n = m^2$ unknowns $u_{i,j}$:

$$u_{i-1,j} + u_{i+1,j} + u_{i,j-1} + u_{i,j+1} - 4u_{i,j} = h^2 f(ih, jh) \quad (1.6)$$

(Note that when i or j is equal to 1 or m , then the values $u_{0,j}, u_{i,0}$ or $u_{i,m+1}, u_{m+1,j}$ are known boundary values and they should be moved to the right-hand side, thus leaving fewer unknowns on the left.) Having ordered grid points, we can write (1.6) as a linear system, say

$$A\mathbf{u} = \mathbf{b}.$$

Our present concern is to prove that, as $h \rightarrow 0$, the numerical solution (1.6) tends to the exact solution of the Poisson equation $\nabla^2 u = f$ (with appropriate Dirichlet boundary conditions).

Example 1.9 (Natural ordering) *The way the matrix A of this system looks depends of course on the way how the grid points (ih, jh) are being assembled in the one-dimensional array. In the natural ordering, when the grid points are arranged by columns, A is the following block tridiagonal matrix:*

$$A = \begin{bmatrix} S & I & & & \\ I & S & I & & \\ & & \ddots & \ddots & \ddots \\ & & & I & S & I \\ & & & & I & S \end{bmatrix}, \quad S = \begin{bmatrix} -4 & 1 & & & \\ 1 & -4 & 1 & & \\ & & \ddots & \ddots & \ddots \\ & & & 1 & -4 & 1 \\ & & & & 1 & -4 \end{bmatrix}.$$

Matlab demo: Check out the online animation for natural ordering at <http://www.maths.cam.ac.uk/undergrad/course/na/ii/partii.php>.

Before heading on let us prove the following simple but useful theorem whose importance will become apparent in the course of the lecture.

Theorem 1.10 (Gershgorin theorem) *All eigenvalues of an $n \times n$ matrix A are contained in the union of the Gershgorin discs in the complex plane:*

$$\sigma(A) \subset \cup_{i=1}^n \Gamma_i, \quad \Gamma_i := \{z \in \mathbb{C} : |z - a_{ii}| \leq r_i\}, \quad r_i := \sum_{j \neq i} |a_{ij}|.$$

Proof. For any matrix A , if $A\mathbf{x} = \lambda\mathbf{x}$ and $|x_i| = \max |x_j|$, then the i th equation of the relation $A\mathbf{x} = \lambda\mathbf{x}$ gives

$$|\lambda - a_{ii}| \cdot |x_i| = \left| \sum_{j \neq i} a_{ij} x_j \right| \leq \sum_{j \neq i} |a_{ij}| |x_j| \leq |x_i| \sum_{j \neq i} |a_{ij}| =: |x_i| r_i,$$

and after dividing by $|x_i|$ we obtain $|\lambda - a_{ii}| \leq r_i$. So, for any eigenvalue λ of A , the inequality $|\lambda - a_{ii}| \leq r_i$ is valid for at least one value of i , hence the theorem. \square

Lemma 1.11 For any ordering of the grid points, the matrix A of the system (1.6) is symmetric and negative definite.

Proof. Equation (1.6) implies that if $a_{ij} \neq 0$ for $i \neq j$, then the i -th and j -th points of the grid (ph, qh) , are nearest neighbours. Hence $a_{ij} \neq 0$ implies $a_{ji} = a_{ji} = 1$, which proves the symmetry of A . Therefore A has real eigenvalues and eigenvectors.

It remains to prove that all the eigenvalues are negative. The arguments are parallel to the proof of Gershgorin theorem. Let $Ax = \lambda x$, and let i be an integer such that $|x_i| = \max |x_j|$. With such an i we address the following identity (which is a reordering of the equation $(Ax)_i = \lambda x_i$):

$$\underbrace{|(\lambda - a_{ii}) x_i|}_{|\lambda+4||x_i|} = \underbrace{|\sum_{j \neq i}^n a_{ij} x_j|}_{\leq 4|x_i|} \quad (1.7)$$

Here $a_{ii} = -4$ and $a_{ij} \in \{0, 1\}$ for $j \neq i$, with at most four nonzero elements on the right-hand side. It is seen that the case $\lambda > 0$ is impossible. Assuming $\lambda = 0$, we obtain $|x_j| = |x_i|$ whenever $a_{ij} = 1$, so we can alter the value of i in (1.7) to any of such j and repeat the same arguments. Thus, the modulus of every component of x would be $|x_i|$, but then the equations (1.7) that occur at the boundary of the grid and have fewer than four off-diagonal terms (see (1.6)) could not be true. Hence, $\lambda = 0$ is impossible too, hence $\lambda < 0$ which proves that A is negative definite. \square

Proposition 1.12 The eigenvalues of the matrix A are

$$\lambda_{k,\ell} = -4 \left(\sin^2 \frac{k\pi h}{2} + \sin^2 \frac{\ell\pi h}{2} \right), \quad h = \frac{1}{m+1}, \quad k, \ell = 1 \dots m,$$

Proof. Let us show that, for every pair (k, ℓ) , the vectors

$$v = (v_{i,j}), \quad v_{i,j} = \sin ix \sin jy, \quad \text{where } x = k\pi h, \quad y = \ell\pi h,$$

are the eigenvectors of A . Indeed, for $i, j = 1 \dots m$, we have

$$\begin{aligned} (Av)_{i,j} &= \sin(jy) [\sin(ix - x) - 2\sin(ix) + \sin(ix + x)] \\ &\quad + \sin(ix) [\sin(jy - y) - 2\sin(jy) + \sin(jy + y)] \\ &= \sin(jy) \sin(ix) [2\cos x - 2] + \sin(ix) \sin(jy) [2\cos y - 2] = \lambda v_{i,j}. \end{aligned}$$

Note that the terms $u_{i \pm 1, j}$, $u_{i, j \pm 1}$ do not appear in (1.6) for $i, j = 1$ or $i, j = m$, respectively, therefore (for such i, j) we should have dropped the corresponding components from above equation, but they are equal to zero because $\sin(i-1)x = 0$ for $i = 1$, while $\sin(i+1)x = 0$ for $i = m$, since $x = \frac{k\pi}{m+1}$. Thus, the eigenvalues are

$$\lambda_{k,\ell} = [2\cos x - 2] + [2\cos y - 2] = -4 \left(\sin^2 \frac{x}{2} + \sin^2 \frac{y}{2} \right) = -4 \left(\sin^2 \frac{k\pi h}{2} + \sin^2 \frac{\ell\pi h}{2} \right). \quad \square$$

Remark 1.13 As a matter of independent mathematical interest, note that for $1 \leq k, \ell \ll m$ we have $\sin x \approx x$, hence the eigenvalues for the discretized Laplacian ∇_h^2 are

$$\frac{\lambda_{k,\ell}}{h^2} \approx -\frac{4}{h^2} \left[\frac{k^2 \pi^2 h^2}{4} + \frac{\ell^2 \pi^2 h^2}{4} \right] = -(k^2 + \ell^2) \pi^2.$$

Now, recall (e.g. from the solution of the Poisson equation in a square by separation of variables in Maths Methods) that the *exact* eigenvalues of ∇^2 (in the unit square) are $-(k^2 + \ell^2) \pi^2$, $k, \ell \in \mathbb{N}$, with the corresponding eigenfunctions $V_{k,\ell}(x, y) = \sin k\pi x \sin \ell\pi y$. So, the eigenvectors of the discretized ∇_h^2 are the values of $V_{k,\ell}(x, y)$ on the grid-points, and the eigenvalues of ∇_h^2 approximate those for continuous case.