

3P1c **Quantum Field Theory: Example Sheet 3** Michaelmas 2018

Corrections and suggestions should be emailed to [B.C.Allanach@damtp.cam.ac.uk](mailto:B.C.Allanach@damtp.cam.ac.uk). Starred questions may if you wish be handed in to your supervisor for feedback prior to the class.

1. The chiral representation of the Clifford algebra is

$$\gamma^0 = \begin{pmatrix} 0 & 1_2 \\ 1_2 & 0 \end{pmatrix}, \quad \gamma^i = \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix}.$$

Show that these indeed satisfy  $\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu}\mathbf{1}$ . Find a unitary matrix  $U$  such that  $(\gamma')^\mu = U\gamma^\mu U^\dagger$ , where  $(\gamma')^\mu$  form the Dirac representation of the Clifford algebra

$$(\gamma')^0 = \begin{pmatrix} 1_2 & 0 \\ 0 & -1_2 \end{pmatrix}, \quad (\gamma')^i = \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix}.$$

2. Show that if  $\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu}$ , then

$$[\gamma^\mu \gamma^\nu, \gamma^\rho \gamma^\sigma] = 2\eta^{\nu\rho} \gamma^\mu \gamma^\sigma - 2\eta^{\mu\rho} \gamma^\nu \gamma^\sigma + 2\eta^{\nu\sigma} \gamma^\rho \gamma^\mu - 2\eta^{\mu\sigma} \gamma^\rho \gamma^\nu.$$

Show further that  $S^{\mu\nu} \equiv \frac{1}{4}[\gamma^\mu, \gamma^\nu] = \frac{1}{2}(\gamma^\mu \gamma^\nu - \eta^{\mu\nu})$ . Use this to confirm that the matrices  $S^{\mu\nu}$  form a representation of the Lie algebra of the Lorentz group.

3. Using just the algebra  $\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu}$  (that is to say without resorting to any particular representation of the gamma matrices), and defining  $\gamma^5 = i\gamma^0\gamma^1\gamma^2\gamma^3$ ,  $\not{p} = p_\mu \gamma^\mu$  and  $S^{\mu\nu} \equiv \frac{1}{4}[\gamma^\mu, \gamma^\nu]$ , prove the following results (these are useful when calculating cross-sections or decay widths involving spinor fields):

- $\text{Tr}\gamma^\mu = 0$
- $\text{Tr}(\gamma^\mu \gamma^\nu) = 4\eta^{\mu\nu}$
- $\text{Tr}(\gamma^\mu \gamma^\nu \gamma^\rho) = 0$
- $(\gamma^5)^2 = 1$
- $\text{Tr}\gamma^5 = 0$
- $\not{p}\not{q} = 2p \cdot q - \not{q}\not{p} = p \cdot q + 2S^{\mu\nu} p_\mu q_\nu$
- $\text{Tr}(\not{p}\not{q}) = 4p \cdot q$
- $\text{Tr}(\not{p}_1 \dots \not{p}_n) = 0$  if  $n$  is odd
- $\text{Tr}(\not{p}_1 \not{p}_2 \not{p}_3 \not{p}_4) = 4[(p_1 \cdot p_2)(p_3 \cdot p_4) + (p_1 \cdot p_4)(p_2 \cdot p_3) - (p_1 \cdot p_3)(p_2 \cdot p_4)]$
- $\text{Tr}(\gamma^5 \not{p}_1 \not{p}_2) = 0$
- $\gamma_\mu \not{p} \gamma^\mu = -2\not{p}$
- $\gamma_a \not{p}_1 \not{p}_2 \gamma^a = 4p_1 \cdot p_2$
- $\gamma_\mu \not{p}_1 \not{p}_2 \not{p}_3 \gamma^\mu = -2\not{p}_3 \not{p}_2 \not{p}_1$
- $\text{Tr}(\gamma^5 \not{p}_1 \not{p}_2 \not{p}_3 \not{p}_4) = 4i \epsilon_{\mu\nu\rho\sigma} p_1^\mu p_2^\nu p_3^\rho p_4^\sigma$

4\* The plane-wave solutions to the Dirac equation are

$$u^s(\vec{p}) = \begin{pmatrix} \sqrt{p \cdot \vec{\sigma}} \xi^s \\ \sqrt{p \cdot \vec{\sigma}} \xi^s \end{pmatrix} \text{ and } v^s(\vec{p}) = \begin{pmatrix} \sqrt{p \cdot \vec{\sigma}} \xi^s \\ -\sqrt{p \cdot \vec{\sigma}} \xi^s \end{pmatrix},$$

where  $\sigma^\mu = (1, \vec{\sigma})$  and  $\bar{\sigma}^\mu = (1, -\vec{\sigma})$  and  $\xi^s$ , with  $s \in \{1, 2\}$ , is a basis of orthonormal two-component spinors, satisfying  $(\xi^r)^\dagger \cdot \xi^s = \delta^{rs}$ . Show that

$$\begin{aligned} u^r(\vec{p})^\dagger \cdot u^s(\vec{p}) &= 2p_0 \delta^{rs} \\ \bar{u}^r(\vec{p}) \cdot u^s(\vec{p}) &= 2m \delta^{rs} \end{aligned} \tag{1}$$

and similarly,

$$\begin{aligned} v^r(\vec{p})^\dagger \cdot v^s(\vec{p}) &= 2p_0 \delta^{rs} \\ \bar{v}^r(\vec{p}) \cdot v^s(\vec{p}) &= -2m \delta^{rs}. \end{aligned} \tag{2}$$

Show also that the orthogonality condition between  $u$  and  $v$  is

$$\bar{u}^s(\vec{p}) \cdot v^r(\vec{p}) = 0,$$

while taking the inner product using  $\dagger$  requires an extra minus sign

$$u^s(\vec{p})^\dagger \cdot v^r(-\vec{p}) = 0. \tag{3}$$

5. Using the same notation as Question 4, show that

$$\sum_{s=1}^2 u^s(\vec{p}) \bar{u}^s(\vec{p}) = \not{p} + m, \tag{4}$$

$$\sum_{s=1}^2 v^s(\vec{p}) \bar{v}^s(\vec{p}) = \not{p} - m, \tag{5}$$

where, rather than being contracted, the two spinors on the left-hand side are placed back to back to form a  $4 \times 4$  matrix.

6. The Fourier decomposition of the Dirac field operator  $\psi(x)$  and the hermitian conjugate field  $\psi^\dagger(\vec{x})$  is given by

$$\begin{aligned} \psi(\vec{x}) &= \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_p}} \sum_{s=1}^2 [b_p^s u^s(\vec{p}) e^{i\vec{p} \cdot \vec{x}} + c_p^{s\dagger} v^s(\vec{p}) e^{-i\vec{p} \cdot \vec{x}}], \\ \psi^\dagger(\vec{x}) &= \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_p}} \sum_{s=1}^2 [b_p^{s\dagger} u^s(\vec{p})^\dagger e^{-i\vec{p} \cdot \vec{x}} + c_p^s v^s(\vec{p})^\dagger e^{i\vec{p} \cdot \vec{x}}]. \end{aligned} \tag{6}$$

The creation and annihilation operators are taken to satisfy

$$\begin{aligned} \{b_p^r, b_q^{s\dagger}\} &= (2\pi)^3 \delta^{rs} \delta^{(3)}(\vec{p} - \vec{q}), \\ \{c_p^r, c_q^{s\dagger}\} &= (2\pi)^3 \delta^{rs} \delta^{(3)}(\vec{p} - \vec{q}), \end{aligned}$$

with all other anticommutators vanishing. Show that these imply that the field and its conjugate field satisfy the anti-commutation relations

$$\begin{aligned} \{\psi_\alpha(\vec{x}), \psi_\beta(\vec{y})\} &= \{\psi_\alpha^\dagger(\vec{x}), \psi_\beta^\dagger(\vec{y})\} = 0, \\ \{\psi_\alpha(\vec{x}), \psi_\beta^\dagger(\vec{y})\} &= \delta_{\alpha\beta} \delta^{(3)}(\vec{x} - \vec{y}). \end{aligned}$$

*Note: the calculation is very similar to that for the bosonic field, but at some point you will need to make use of the identities Eqs. (4), (5).*

7\* Using the results of Question 6, show that the quantum Hamiltonian

$$H = \int d^3x \bar{\psi}(-i\gamma^i \partial_i + m)\psi$$

can be written, after normal ordering, as

$$H = \int \frac{d^3p}{(2\pi)^3} E_{\vec{p}} \sum_{r=1}^2 [b_{\vec{p}}^{r\dagger} b_{\vec{p}}^r + c_{\vec{p}}^{r\dagger} c_{\vec{p}}^r].$$

*Note: again, the calculation is very similar to that of the bosonic field. This time you will need to make use of the identities in Eqs. (1), (2) and (3).*

8. Standard fermion Yukawa theory has the Lagrangian density

$$\mathcal{L} = \bar{\psi}(i \not{\partial} - m)\psi + \frac{1}{2} \partial^\mu \phi \partial_\mu \phi - \frac{1}{2} \mu^2 \phi^2 - \lambda \bar{\psi} \psi \phi.$$

Show that the differential cross-section in the centre of mass frame for nucleon-nucleon scattering ( $\psi\psi \rightarrow \psi\psi$ ) including the masses  $m$  and  $\mu$  is

$$\frac{d\sigma}{dt} = \frac{\lambda^4}{16\pi s(s-4m^2)} \left[ \frac{(u-4m^2)^2}{(u-\mu^2)^2} + \frac{(t-4m^2)^2}{(t-\mu^2)^2} + \frac{1}{2} \frac{s^2 - (u-4m^2)^2 - (t-4m^2)^2}{(u-\mu^2)(t-\mu^2)} \right].$$

What values of  $t$  should this be integrated between to obtain the total cross-section  $\sigma$ ?