

# String Theory: Example Sheet 2

Malcolm Perry, February 2011

1. Construct the open string states at level 2 in the lightcone formalism and determine their representation under  $SO(D-1)$ . Construct the states at level 3. Show that they fit into a traceless symmetric-3-tensor and an anti-symmetric-2-tensor representation of  $SO(D-1)$ .

2. Write down the mode expansion describing an open string stretched between two parallel  $Dp$ -branes. Interpret the result.

3. Verify that

$$\partial\bar{\partial} \ln |z|^2 = 2\pi\delta(z, \bar{z}) ,$$

firstly by using the divergence theorem, and secondly by regulating the singularity at  $z = 0$ .

4. Show that  $: e^{ikX} :$  is a primary operator for the theory of a free scalar field. Compute the weight.

5. A theory of a free scalar field  $X$  has OPE

$$\partial X(z) \partial X(w) = -\frac{\alpha'}{2} \frac{1}{(z-w)^2} + \dots$$

Consider a putative candidate for the stress-energy tensor

$$T(z) = -\frac{1}{\alpha'} : \partial X(z) \partial X(z) : -Q \partial^2 X(z)$$

a. Use the  $TX$  OPE to determine the transformation of  $X$  under conformal transformations  $\delta z = \epsilon(z)$ .

b. Show that  $\partial X$  is not primary unless  $Q = 0$ , but is quasi-primary with weight  $h = 1$ . Show that  $: e^{ikX} :$  is primary and compute its weight.

c. Determine the  $TT$  OPE and show that it does have the structure expected of a stress-energy tensor. What is the central charge of the theory?

**An Aside:** What's going on here? How can the same free scalar field have different stress-energy tensors? The point is that the stress energy tensor tells us how the system couples to a curved metric. Several such couplings could all give the same physics when restricted to a flat background. The theory considered here is called the *linear dilaton* theory. We will see where it comes from later in the course.

6. A theory of several free, non-interacting scalars  $X^\mu$ ,  $\mu = 1, \dots, D$ , has the operators

$$\zeta_\mu : \partial X^\mu e^{ik \cdot X} : \quad \text{and} \quad \zeta_{\mu\nu} : \partial X^\mu \bar{\partial} X^\nu e^{ik \cdot X} :$$

where  $\zeta_\mu$ ,  $k_\mu$  are constant vectors and  $\zeta_{\mu\nu}$  is a constant tensor. If the stress-energy tensor is given by  $T = (-1/\alpha') : \partial X^\mu \partial X_\mu :$  determine the conditions for these operators to be primary. What are their weights?

7. A free fermion Majorana fermion in two dimensions has action,

$$S = \frac{1}{2\pi} \int d^2z \psi \bar{\partial} \psi + \bar{\psi} \partial \psi$$

The propagator is given by the OPE

$$\psi(z)\psi(w) = -\psi(w)\psi(z) = \frac{1}{z-w}$$

and similar for  $\bar{\psi}$ . (Remember,  $\psi$  and  $\bar{\psi}$  are Grassmann-valued fields, a fact which is reflected in the OPE). The energy momentum tensor is

$$T_{zz} = -\frac{1}{2} : \psi \partial \psi :$$

Show that  $\psi$  is a primary operator of weight  $1/2$ . Determine the central charge of this theory.

8. The bc ghost system consist of two free Grassmann fields  $b$  and  $c$ . (Note: do not confuse the field  $c$  with the central charge  $c$ . They are not the same thing!) The OPE is given by

$$b(z)c(w) = -c(w)b(z) = \frac{1}{z-w}$$

Consider the stress-energy tensor

$$T = : (\partial b)c : - \lambda \partial : bc :$$

Show that  $b$  is primary with weight  $h = \lambda$  and  $c$  is primary with weight  $h = 1 - \lambda$ . Show that the central charge of this system is equal to

$$c = -12\lambda^2 + 12\lambda - 2$$

**An Aside:** This peculiar looking theory is *extremely* important. We will come across it later in the course when we discuss the path integral approach to string theory.

**9.** Show that the Schwarzian transformation of the stress tensor reproduces the correct infinitesimal transformation. Show, moreover, that it has the correct property under successive conformal transformations.

**8.** Use the OPE for a free scalar field to determine the commutation relations of the Fourier modes  $\alpha_m$ .