

## Part III Applications of Differential Geometry to Physics, Sheet Two

Maciej Dunajski, Lent Term 2016

1. Let  $\phi : \mathbb{R}^{2,1} \rightarrow S^2$ . Set

$$\phi^1 + i\phi^2 = \frac{2u}{1 + |u|^2}, \quad \phi^3 = \frac{1 - |u|^2}{1 + |u|^2},$$

and deduce that the Bogomolny equations

$$\partial_i \phi^a = \pm \varepsilon_{ij} \varepsilon^{abc} \phi^b \partial_j \phi^c, \quad \phi_t = 0$$

imply that  $u$  is holomorphic or antiholomorphic in  $z = x_1 + ix_2$ .

Find the expression for the total energy

$$E[\phi] = \frac{1}{2} \int \partial_j \phi^a \partial_j \phi^a d^2x$$

in terms of  $u$ .

By counting the pre-images or otherwise find the topological degree of  $\phi$  corresponding to  $u(z) = u_0 + u_1z + \dots + u_kz^k$ , where  $u_0, \dots, u_k$  are constants with  $u_k \neq 0$ .

2. Derive the  $SU(2)$  Yang–Mills theory on  $\mathbb{R}^4$  from the action. Let  $A_a(x)$  be a solution to these equations. Show that, for any nonzero constant  $c$ , the potential  $\tilde{A}_a(x) = cA_a(cx)$  is also a solution and that it has the same action.
3. Consider the map  $g : S^3 \rightarrow SU(2)$  defined by

$$g(x_1, x_2, x_3, x_4) = x_4 + i(x_1\sigma_1 + x_2\sigma_2 + x_3\sigma_3),$$

where  $\sigma_i$  are Pauli matrices and  $x_1^2 + x_2^2 + x_3^2 + x_4^2 = 1$  and find its degree. By calculating  $\text{Tr}((dg g^{-1})^3)$  at the point on  $S^3$  where  $x_4 = 1$ , or otherwise deduce that the formula

$$\text{deg}(g) = \frac{1}{24\pi^2} \int_{S^3} \text{Tr}((dg g^{-1})^3)$$

is correctly normalised.

4. Let  $T_1, T_2, T_3$  form a basis of  $\mathfrak{su}(2)$  such that

$$[T_\alpha, T_\beta] = -\varepsilon_{\alpha\beta\gamma} T_\gamma, \quad \alpha, \beta, \gamma = 1, 2, 3,$$

and let the symbols  $\sigma_{ab} = -\sigma_{ba}$  where  $a, b = 1, \dots, 4$  be defined by

$$\sigma_{\alpha\beta} = \varepsilon_{\alpha\beta\gamma} T_\gamma, \quad \sigma_{\alpha 4} = T_\alpha.$$

Show that

$$\sigma_{ab} = \frac{1}{2} \varepsilon_{ab}{}^{cd} \sigma_{cd}, \quad \text{and} \quad \sigma_{ab} \sigma_{ac} = -\frac{3}{4} \mathbf{1} \delta_{bc} - \sigma_{bc}.$$

Identify  $\Lambda^2(\mathbb{R}^4)$  with the Lie algebra  $\mathfrak{so}(4)$  and deduce that  $\mathfrak{so}(4) = \mathfrak{so}(3) \oplus \mathfrak{so}(3)$ .

5. Let  $V = 1 + r^{-2}$ , where  $r^2 := \delta_{ab} x^a x^b$ . Show that the one-form

$$A = \sigma_{ab} \frac{1}{V} \frac{\partial V}{\partial x^b} dx^a \tag{1}$$

is a solution of the anti-self-dual Yang–Mills equations on  $\mathbb{R}^4$ .

The one-form  $A$  is singular at  $r = 0$ . What can you say about the behaviour of the field strength  $F$  at  $r = 0$ ?

6. Find, by explicit integration, the Chern number of the solution (1).
7. Let  $F$  be a two-form on  $\mathbb{R}^4$ . Show, from the definition of the Hodge operator, that
- (a)  $**F = \pm F$  depending on the signature.
  - (b)  $*F \wedge *F = F \wedge F$ .

Show that in the  $U(1)$  theory  $F \rightarrow *F$  interchanges the electric and magnetic fields with factors of  $\pm 1$  or  $\pm i$  and determine the different cases in the corresponding signatures.

Let  $F$  be a non-zero real self-dual two-form on  $\mathbb{R}^4$  such that  $F \wedge F = 0$ . What is the signature of the underlying metric?

8. Let  $A$  be a 1-form gauge potential on  $\mathbb{R}^n$  with values in  $\mathfrak{su}(2)$ , and let  $F$  be its curvature. Verify that  $Tr(A), Tr(A \wedge A), Tr(A \wedge A \wedge A \wedge A)$  and  $Tr(F)$  all vanish.

Verify that  $C_2 = dY_3$ , where  $C_2$  and  $Y_3$  are the second Chern form, and the Chern–Simons three-form respectively.

9. Let  $A = A_i dx^i$ ,  $i = 1, 2, 3$  be a gauge potential on  $\mathbb{R}^3$  with values in the Lie algebra  $\mathfrak{g}$ . Find the Euler–Lagrange equations arising from varying the Chern–Simons functional

$$W[A] = \int_{\mathbb{R}^3} \text{Tr}(A \wedge dA + \frac{2}{3} A \wedge A \wedge A)$$

with respect to  $A$ .

Now consider a one parameter family of  $\mathfrak{g}$ -valued one-forms  $A = A(t)$  on  $\mathbb{R}^3$ , and define a one-form on  $\mathbb{R}^4$  by  $\mathcal{A} = A + \phi dt$ , where the function  $\phi = \phi(x^i, t)$  takes its values in  $\mathfrak{su}(2)$ . Show that, in a gauge where  $\phi = 0$ , the anti-self-dual Yang–Mills equations on  $\mathcal{A}$  take the gradient flow form

$$\frac{dA_i(t)}{dt} = \frac{\delta W[A]}{\delta A_i}.$$

10. Consider a connection  $\omega = \gamma^{-1}A\gamma + \gamma^{-1}d\gamma$  on a principal  $G$ -bundle  $P \rightarrow B$ , where  $A$  is a one-form on  $B$  and  $\gamma^{-1}d\gamma$  is the Maurer–Cartan form on  $G$ .
- Show that the transformation of the fibres  $\gamma' = g\gamma$ , where  $g \in G$  depends on the coordinates on  $B$ , does not change  $\omega$  if  $A$  transforms like a gauge potential.
  - Let  $\Omega = d\omega + \omega \wedge \omega$ . Show that  $\Omega = \gamma^{-1}F\gamma$  for some  $F$  which should be found.
  - Let  $D_a$ ,  $a = 1, \dots, \dim(B)$  be linearly independent vector fields on  $P$  such that

$$D_a \lrcorner \omega = 0.$$

Show that  $D_a = \partial_a - A_a^\alpha R_\alpha$ , where  $\partial_a = \partial/\partial x^a$  are vector fields on  $B$  and  $R_\alpha$  are right-invariant vector fields on  $G$ . Demonstrate that

$$[D_a, D_b] = -F_{ab}^\alpha R_\alpha.$$