

Example Sheet 1

1. A steady two-dimensional flow (pure straining) is given by $u = \alpha x$, $v = -\alpha y$ with α constant.
 - (i) Find the equation for a general streamline of the flow, and sketch some of them.
 - (ii) At $t = 0$ the fluid on the curve $x^2 + y^2 = a^2$ is marked (by an electro-chemical technique). Find the equation for this material fluid curve for $t > 0$.
 - (iii) Does the area within the curve change in time, and why?

2. Repeat question 1 for the two-dimensional flow (simple shear) given by $u = \gamma y$, $v = 0$ with γ constant. Which of the two flows stretches the curve faster at long times?

3. A two-dimensional flow is represented by a streamfunction $\psi(x, y)$ with $u = \partial\psi/\partial y$ and $v = -\partial\psi/\partial x$. Show that
 - (i) the streamlines are given by $\psi = \text{const}$,
 - (ii) $|\mathbf{u}| = |\nabla\psi|$, so that the flow is faster where the streamlines are closer,
 - (iii) the volume flux crossing any curve from \mathbf{x}_0 to \mathbf{x}_1 is given by $\psi(\mathbf{x}_1) - \psi(\mathbf{x}_0)$,
 - (iv) $\psi = \text{const}$ on any *fixed* (i.e. stationary) boundary.

[Hint for (iii): $\mathbf{n} ds = (dy, -dx)$.]

4. Verify that the two-dimensional flow given in Cartesian coordinates by

$$u = \frac{y - b}{(x - a)^2 + (y - b)^2}, \quad v = \frac{a - x}{(x - a)^2 + (y - b)^2}$$

satisfies $\nabla \cdot \mathbf{u} = 0$, and then find the streamfunction $\psi(x, y)$ such that $u = \partial\psi/\partial y$ and $v = -\partial\psi/\partial x$. Sketch the streamlines.

5. Verify that the two-dimensional flow given in polar coordinates by

$$u_r = U \left(1 - \frac{a^2}{r^2} \right) \cos \theta, \quad u_\theta = -U \left(1 + \frac{a^2}{r^2} \right) \sin \theta$$

satisfies $\nabla \cdot \mathbf{u} = 0$, and find the streamfunction $\psi(r, \theta)$. Sketch the streamlines.

$$\left[\text{Take: } \nabla \cdot \mathbf{u} = \frac{1}{r} \frac{\partial}{\partial r} (ru_r) + \frac{1}{r} \frac{\partial}{\partial \theta} (u_\theta) \quad \text{and} \quad u_r = \frac{1}{r} \frac{\partial \psi}{\partial \theta}, \quad u_\theta = -\frac{\partial \psi}{\partial r} \right]$$

6. Verify that the axisymmetric flow (uniaxial straining) given in cylindrical polar coordinates by $u_r = -\frac{1}{2}\alpha r$, $u_z = \alpha z$ satisfies $\nabla \cdot \mathbf{u} = 0$, and find the Stokes streamfunction $\Psi(r, z)$. Sketch the streamlines.

$$\left[\text{Take: } \nabla \cdot \mathbf{u} = \frac{1}{r} \frac{\partial}{\partial r} (ru_r) + \frac{\partial u_z}{\partial z} \quad \text{and} \quad u_r = -\frac{1}{r} \frac{\partial \Psi}{\partial z}, \quad u_z = \frac{1}{r} \frac{\partial \Psi}{\partial r} \right]$$

7. Consider the two-dimensional flow $u = 1/(1+t)$, $v = 1$ in $t > -1$. Find and sketch
- the streamline at $t = 0$ which passes through the point $(1, 1)$,
 - the path of a fluid particle which is released from $(1, 1)$ at $t = 0$,
 - the position at $t = 0$ of a streak of dye released from $(1, 1)$ during the time interval $-1 < t \leq 0$.

8. An axisymmetric jet of water of speed 1 m s^{-1} and cross-section $6 \times 10^{-4} \text{ m}^2$ strikes a wall at right angles and spreads out over it. By using the momentum integral equation, calculate the force on the wall due to the jet. [Neglect gravity.]

9. Starting from the Euler momentum equation for a fluid of constant density with a potential force $-\nabla\chi$, show that for a fixed volume V enclosed by surface A

$$\frac{d}{dt} \int_V \frac{1}{2} \rho u^2 dV + \int_A H \mathbf{u} \cdot \mathbf{n} dA = 0$$

where $H = \frac{1}{2} \rho u^2 + p + \chi$ is the Bernoulli quantity, so concluding that H is the transportable energy.

10. How high can water rise up one's arm hanging in the river from a lazy (1 m s^{-1}) punt? [Use Bernoulli on surface streamline, where $p = 1$ atmosphere.]

11. A rotating circular tank of radius a is filled with a volume V of fluid of density ρ . The tank is allowed to rotate for a long time, until the flow is a two-dimensional rigid-body motion with constant angular velocity ω around the axis (so that $u = -\omega y$, $v = \omega x$, $w = 0$, $\rho = \text{const}$). Derive the equation for the pressure, p , at any point in the rotating fluid. What is the equation for the height $h(r)$ of the free surface? (Hint: Integrate the Euler equation to find the pressure and determine the constant of integration from the volume conservation).

12. Waste water flows into a tank at $10^{-4} \text{ m}^3 \text{ s}^{-1}$ and out of a short exit pipe of cross-section $4 \times 10^{-5} \text{ m}^2$ into the air. In steady state, estimate how high above the pipe is the water in the tank?

13. A water clock is an axisymmetric vessel with a small exit pipe in the bottom. Find the shape for which the water level falls equal heights in equal intervals of time.