

Example Sheet 2

1. Show that $\mathbf{u}(\mathbf{x} + \delta\mathbf{x}) - \mathbf{u}(\mathbf{x})$ can be written as $(e_{ij} + \frac{1}{2}\varepsilon_{jik}\omega_k)\delta x_j + O(|\delta\mathbf{x}|^2)$, where $e_{ij} = e_{ji}$ and $\omega_k = (\nabla \times \mathbf{u})_k$. Find e_{ij} and ω_k for the case of linear shear flow $\mathbf{u} = (y, 0, 0)$, and for this case sketch the streamlines of the two flows whose velocity fields are $e_{ij}x_j$ and $\frac{1}{2}\varepsilon_{jik}\omega_k x_j$.

2. Calculate the vorticity $\boldsymbol{\omega}$ of the velocity field

$$u = -\alpha x - yrf(t), \quad v = -\alpha y + xrf(t), \quad w = 2\alpha z$$

where $r^2 = x^2 + y^2$. Show that $\nabla \cdot \mathbf{u} = 0$ for any function $f(t)$, and that the vorticity equation $\partial\boldsymbol{\omega}/\partial t + (\mathbf{u} \cdot \nabla)\boldsymbol{\omega} = (\boldsymbol{\omega} \cdot \nabla)\mathbf{u}$ is satisfied if and only if $f(t) \propto e^{3\alpha t}$.

Calculate the velocity components u_r and u_θ in cylindrical polar coordinates. Consider a material curve that starts at $t = 0$ as $z = 0, r = a_0$. Show that this becomes $z = 0, r = a_0 e^{-\alpha t}$. Verify the circulation theorem for this material curve.

3. If $\mathbf{u} = \boldsymbol{\Omega} \times \mathbf{x}$ (uniform rotation with angular velocity $\boldsymbol{\Omega}$) show that $\boldsymbol{\omega} = 2\boldsymbol{\Omega}$, where $\boldsymbol{\omega}$ is the vorticity. For a two-dimensional flow $(u(x, y), v(x, y), 0)$, show that $\boldsymbol{\omega} = (0, 0, -\nabla^2\psi)$, where ψ is the streamfunction.

A long cylinder filled with water has elliptical cross-section with major and minor semi-axes a and b . While $t < 0$, both the cylinder and the water within it rotate about the axis of the cylinder with uniform angular velocity $(0, 0, \Omega)$. Sketch the streamlines, noting that they intersect the elliptical boundary of the cylinder. (Why?)

At $t = 0$, the cylinder is suddenly brought to rest. Assuming that the flow remains two-dimensional, what does the vorticity equation say about $\nabla^2\psi$ for $t > 0$? Verify that the flow for $t > 0$ can be described by

$$\psi = \frac{a^2 b^2 \Omega}{a^2 + b^2} \left(1 - \frac{x^2}{a^2} - \frac{y^2}{b^2} \right)$$

in suitable coordinates, and sketch the streamlines.

4. Show that velocity potential ϕ cannot have an extreme value at an interior point of the fluid. Deduce that the speed cannot have a maximum value at an interior point. (Hint: ϕ and its partial derivatives satisfy Laplace's equation.)

5. (1) Is the motion incompressible for the flows given by the following velocity potentials: (a) $\phi = C(x^2 + y^2)$ (b) $\phi = C(x^2 - y^2)$? If so, determine the corresponding stream functions. (2) Is the motion irrotational for the flows given by the following stream functions: (a) $\psi = C(x^2 + y^2)$ (b) $\psi = C(x^2 - y^2)$? If so, determine the corresponding velocity potentials. Sketch the streamlines for all cases ((1) and (2)) and the lines of constant ϕ where possible.

6. A sphere of radius a moves with constant velocity U in a fluid otherwise at rest. How far ahead of the sphere is there a disturbance of magnitude $\frac{1}{20}U$ relative to the far-field? Show that the acceleration of a fluid particle at distance x ahead of the centre of the sphere is

$$3U^2 \left(\frac{a^3}{x^4} - \frac{a^6}{x^7} \right).$$

7. In spherical polar coordinates (r, θ, ξ) the velocity potential

$$\phi = -kr^2 P_2(\cos \theta)$$

represents the axisymmetric flow produced by the confluence of two equal and opposite streams. Sketch the stream line pattern in any plane through the axis of symmetry. If now a solid sphere with surface $r = a$ is placed in this flow, how is the velocity potential modified? Hence, determine the distribution of velocity and pressure over the surface of the sphere.

8. Write down the velocity potential $\phi(r, \theta, z)$ for the axisymmetric flow produced by a point source of strength m located at the origin, in the presence of a uniform stream $(0, 0, U)$. Show that there is a stagnation point at $(0, 0, -\frac{1}{2}a)$, where $a = (m/\pi U)^{1/2}$. Sketch the streamlines. Show that the Stokes streamfunction is given by $\Psi = \frac{1}{2}Ur^2 - mz/4\pi(z^2 + r^2)^{1/2}$. From the sketch and the streamfunction show that ϕ represents the flow around a semi-infinite body whose radius tends to a far downstream.

Sketch the flow if the source is re-located at $(0, 0, -l)$ and a sink of equal strength introduced at $(0, 0, l)$.

9. An orifice in the side of a large open vessel full of water leads smoothly into a horizontal tube of uniform cross-section and length L . The diameter of the tube is small in comparison with L and with the size of the vessel and the depth h of the orifice below the free surface. A plug at the end of the tube is suddenly removed and the water begins to flow. Neglecting small changes in h , show from the relevant form of Bernoulli's theorem, or otherwise, that the outflow velocity at subsequent times t is approximately

$$\sqrt{2gh} \tanh \left(\frac{t\sqrt{2gh}}{2L} \right) .$$

Deduce that the time it takes for the flow to accelerate to a fraction $(e^2 - 1)/(e^2 + 1) = 0.7616$ of its limiting value is $2L/\sqrt{(2gh)}$, and verify that this time is about 3 seconds for a garden hose $L = 9$ m long supplied by a rainwater tank with $h = 1.8$ m.

10. A U-tube consists of two long uniform vertical tubes of different cross-sectional areas A_1, A_2 connected at the base by a short tube of large cross-section, and contains an inviscid, incompressible fluid whose surface, in equilibrium, is at height h above the base. Derive the equation governing the nonlinear oscillations of the displacement ζ of the surface in the tube of cross-section A_2

$$(h + r\zeta) \frac{d^2\zeta}{dt^2} + \frac{r}{2} \left(\frac{d\zeta}{dt} \right)^2 + g\zeta = 0 \quad \text{where} \quad r = 1 - A_2/A_1 .$$

[Hint: Take $\phi = 0$ at the bottom of the U-tube, and remember that the irrotational form of Bernoulli's theorem involves $\partial\phi/\partial t$, implying time differentiation at a fixed location.]

11. A sphere is immersed in an infinite calm ocean of incompressible fluid of density ρ . Its radius is given by $R(t) = a + b \sin nt$, where a, b and n are constants with $b < a$. The fluid moves radially under no external forces and the constant pressure at infinity is P . If $a \geq 5b$, show that the maximum pressure attained on the surface of the sphere is $P + \rho n^2 b(a - b)$. What is the corresponding formula if $b < a \leq 5b$?