

Mathematical Tripos Part II
STATISTICAL PHYSICS Examples 3

H.S. Reall
 Lent 2018

1. A *Wigner crystal* is a triangular lattice of electrons in a two dimensional plane. The longitudinal vibration modes of this crystal are bosons with dispersion relation $\omega = \alpha\sqrt{|\mathbf{k}|}$. Show that, at low temperatures, these modes provide a contribution to the heat capacity that scales as $C \sim T^4$.
2. Use the fact that the density of states is constant in $d = 2$ dimensions to show that Bose-Einstein condensation does not occur no matter how low the temperature.
3. Consider N non-interacting, non-relativistic bosons, each of mass m , in a cubic box of side L . Show that the transition temperature scales as $T_c \sim N^{2/3}/mL^2$ and the 1-particle energy levels scale as $E_n \propto 1/mL^2$. Show that when $T < T_c$, the mean occupancy of the first few excited 1-particle states is large, but not as large as $\mathcal{O}(N)$.
4. Consider an ideal gas of bosons whose density of states is given by $g(E) = CE^{\alpha-1}$ for some constants C and $\alpha > 1$. Derive an expression for the critical temperature T_c , below which the gas experiences Bose-Einstein condensation.
5. In experiments on Bose-Einstein condensates, atoms are confined in magnetic traps which can be modelled by a quadratic potential of the type discussed in Question 9 of Example Sheet 2. Determine T_c for bosons in a three dimensional trap. Show that bosons in a two dimensional trap will condense at suitably low temperatures. In each case, calculate the number of particles in the condensate as a function of $T < T_c$.
6. A system has two energy levels with energies 0 and ϵ . These can be occupied by (spinless) fermions from a particle and heat bath with temperature T and chemical potential μ . The fermions are non-interacting. Show that there are four possible microstates, and show that the grand partition function is

$$\mathcal{Z}(T, V, \mu) = 1 + z + ze^{-\beta\epsilon} + z^2e^{-\beta\epsilon}$$

where $z = e^{\beta\mu}$. Evaluate the average occupation number of the state of energy ϵ , and show that this is compatible with the result of the calculation of the average energy of the system using the Fermi-Dirac distribution. How could you take account of fermion interactions?

7. In an ideal Fermi gas the average occupation numbers of the single particle state $|r\rangle$ is n_r . Show that the entropy

$$S = \frac{\partial}{\partial T}(kT \log \mathcal{Z})_{V, \mu}$$

can be written as

$$S = -k \sum_r [(1 - n_r) \log(1 - n_r) + n_r \log n_r]$$

Find the corresponding expression for an ideal Bose gas.

8. Show that $(\Delta n_r)^2 = n_r(1 - n_r)$ for the ideal Fermi gas. Comment on this result, especially for very low T . What is the corresponding result for an ideal Bose gas? How does $\Delta n_0/n_0$ behave at low T for the Bose gas?
9. As a simple model of a semiconductor, suppose that there are N bound electron states, each having energy $-\Delta < 0$, which are filled at zero temperature. At non-zero temperature some electrons are excited into the conduction band, which is a continuum of positive energy states. The density of these states is given by $g(E)dE = A\sqrt{E}dE$ where A is a constant. Show that at temperature T the mean number \bar{n}_c of excited electrons is determined by the pair of equations

$$n_c = \frac{N}{e^{(\mu+\Delta)/kT} + 1} = \int_0^\infty \frac{g(E) dE}{e^{(E-\mu)/kT} + 1}.$$

Show also that, if $n_c \ll N$ and $kT \ll \Delta$ and $e^{\mu/kT} \ll 1$, then

$$2\mu \approx -\Delta + kT \log \left[\frac{2N}{A\sqrt{\pi}(kT)^3} \right].$$

10. Consider an almost degenerate Fermi gas of electrons with spin degeneracy $g_s = 2$. At high temperatures, show that the equation of state is given by

$$pV = NkT \left(1 + \frac{\lambda^3 N}{4\sqrt{2}g_s V} + \dots \right)$$

At low temperatures, show that the chemical potential is

$$\mu = E_F \left(1 - \frac{\pi^2}{12} \left(\frac{kT}{E_F} \right)^2 + \dots \right)$$

and the average energy is

$$E = \frac{3NE_F}{5} \left(1 + \frac{5\pi^2}{12} \left(\frac{kT}{E_F} \right)^2 + \dots \right)$$

11. Consider a gas of non-interacting ultra-relativistic electrons, whose mass may be neglected. Find an integral for the grand potential Φ . Show that $3pV = E$. Show that at zero temperature $pV^{4/3} = \text{const}$. Show that at high temperatures $E = 3NkT$ and the equation of state coincides with that of a classical ultra-relativistic gas.

12. A crude non-relativistic model of a white dwarf star consists of a sphere of radius R of free electrons at zero temperature together with a sufficient number of protons to make the star electrically neutral. Determine the energy E_{el} of all the electrons. Assuming the gravitational energy of the star is given by $E_{\text{grav}} = -\gamma M^2/R$, where M is the total mass of the star, show that if the state of equilibrium of the star is given by minimising the total energy ($E_{\text{grav}} + E_{\text{el}}$) then R is proportional to $M^{-\frac{1}{3}}$. What justification can be given for neglecting the proton zero-point energy?