

## Examples Sheet #3

**1. Nonlinear Diffusion.** For the nonlinear diffusion equation  $C_t = D(C^p C_x)_x$ , where  $p$  and  $D$  are strictly positive constants, show that the self similar solution defined in  $x \geq 0$  of the form

$$C(x, t) = \frac{M^{2/(2+p)}}{(Dt)^{1/(2+p)}} F(\xi), \quad \xi = \frac{x}{(M^p Dt)^{1/(2+p)}}$$

which satisfies  $\int_0^\infty C(x, t) dx = M$ ,  $C_x(0, t) = 0$  and  $C(x \rightarrow \infty, t) \rightarrow 0$  for some constant  $M > 0$ , is given by

$$F(\xi) = \left[ A - \frac{p\xi^2}{2(2+p)} \right]^{1/p}, \quad 0 < \xi < \left[ \frac{2(2+p)A}{p} \right]^{1/2}$$

and  $F = 0$  otherwise. For the case  $p = 1$ , prove that  $A = (3/8)^{1/3}$ .

**2. The Fitzhugh-Nagumo model.** The Fitzhugh-Nagumo model for nerve impulse propagation is given by

$$\begin{aligned} u_t &= u_{xx} + u(1-u)(u-a) + v \\ v_t &= bu - cv, \end{aligned}$$

with  $b, c > 0$  and  $0 < a < 1$ . (i) Show that the homogeneous system has a stable fixed point for which neither  $u$  nor  $v$  is zero. (ii) Seek a travelling wave solution of the equations of the form

$$u = \phi(\xi), v = \psi(\xi), \xi = x - \gamma t.$$

Deduce the ODEs to be satisfied by  $\phi, \psi$ . (iii) Verify that if  $b = c = \psi(0) = 0$  then there is a solution of the form

$$\phi = \frac{1}{1 + e^{-\alpha\xi}},$$

for two possible values of  $\alpha$  which should be found. What is the wave speed  $\gamma$  in each case? In what direction does the wave propagate?

**3. Chemotaxis.** Consider the chemostatic system

$$n_t = Dn_{xx} + bn \left( 1 - \frac{n}{n_0} \right) - (\chi(a)na_x)_x$$

$$a_t = D_A a_{xx} + hn - ka,$$

where  $\chi(a) = \chi_0 K / (K + a)^2$ . Find a scaling such that this reduces to

$$\dot{u} = u'' + u(1-u) - \beta \left[ \frac{uv'}{(\alpha + v)^2} \right]'$$

$$\dot{v} = \delta v'' + \gamma(u - v),$$

where  $\cdot$  and  $\prime$  refer to differentiation with respect to the scaled  $t$  and  $x$  respectively, and  $\alpha, \beta, \gamma, \delta$  are positive constants. Show that the uniform, steady solution  $u = v = 1$  is unstable if

$$\frac{\beta\gamma}{(1+\alpha)^2} > (\sqrt{\gamma} + \sqrt{\delta})^2,$$

and find the wavenumber at which the system first becomes unstable as  $\chi_0$  is increased from zero, in the case  $\alpha = \gamma = \delta = 1$ .

**4. Turing instability.** Investigate the possibility of Turing instability for the reaction-diffusion system.

$$\begin{aligned}\frac{\partial u}{\partial t} &= \nabla^2 u + \frac{u^2}{v} - bu, \\ \frac{\partial v}{\partial t} &= d\nabla^2 v + u^2 - v.\end{aligned}$$

In particular, find the region of the parameter space  $(b, d)$  in which Turing instability can occur, and give a value for the critical wavenumber at the onset of instability.

**5. Phytoplankton-zooplankton.** A space-dependent phytoplankton - zooplankton model can be reduced to the following equations

$$\begin{aligned}u_t &= \nabla^2 u + u + u^2 - \gamma uv \\ v_t &= d\nabla^2 v + \beta uv - v^2.\end{aligned}$$

Find the regions in the  $\beta - \gamma$  plane (a) in which there is a stable, homogeneous state  $(u_0, v_0)$  in which neither  $u_0$  nor  $v_0$  is zero and (b) in which that state may be unstable to a Turing instability. In case (b), for what values of  $d$  will the instability occur, and what is the critical wavenumber for the onset of the instability?