

Mathematical Tripos Part II: Michaelmas Term 2022

Numerical Analysis – Lecture 9

Problem 2.25 (The diffusion equation in two space dimensions) We are solving

$$\frac{\partial u}{\partial t} = \nabla^2 u, \quad 0 \leq x, y \leq 1, \quad t \geq 0, \quad (2.11)$$

where $u = u(x, y, t)$, together with initial conditions at $t = 0$ and Dirichlet boundary conditions at $\partial\Omega$, where $\Omega = [0, 1]^2 \times [0, \infty)$. It is straightforward to generalize our derivation of numerical algorithms, e.g. by semi-discretization (also known as the method of lines). Thus, let $u_{\ell,m}(t) \approx u(\ell h, mh, t)$, where $h = \Delta x = \Delta y$, and let $u_{\ell,m}^n \approx u_{\ell,m}(nk)$ where $k = \Delta t$. The five-point formula results in

$$u'_{\ell,m} = \frac{1}{h^2} (u_{\ell-1,m} + u_{\ell+1,m} + u_{\ell,m-1} + u_{\ell,m+1} - 4u_{\ell,m}),$$

or in the matrix form

$$\mathbf{u}' = \frac{1}{h^2} A_* \mathbf{u}, \quad \mathbf{u} = (u_{\ell,m}) \in \mathbb{R}^N, \quad (2.12)$$

where A_* is the block TST (Toeplitz Symmetric Tridiagonal) matrix of the five-point scheme:

$$A_* = \begin{bmatrix} H & I & & & \\ & I & \ddots & & \\ & & \ddots & \ddots & \\ & & & \ddots & I \\ & & & & I & H \end{bmatrix}, \quad H = \begin{bmatrix} -4 & 1 & & & \\ & 1 & \ddots & & \\ & & \ddots & \ddots & \\ & & & \ddots & 1 \\ & & & & 1 & -4 \end{bmatrix}.$$

Thus, the Euler method yields

$$u_{\ell,m}^{n+1} = u_{\ell,m}^n + \mu (u_{\ell-1,m}^n + u_{\ell+1,m}^n + u_{\ell,m-1}^n + u_{\ell,m+1}^n - 4u_{\ell,m}^n), \quad (2.13)$$

or in the matrix form

$$\mathbf{u}^{n+1} = A \mathbf{u}^n, \quad A = I + \mu A_*$$

where, as before, $\mu = \frac{k}{h^2} = \frac{\Delta t}{(\Delta x)^2}$. The local error is $\eta = \mathcal{O}(k^2 + kh^2) = \mathcal{O}(h^4)$. To analyse stability, we notice that A is symmetric, hence normal, and its eigenvalues are related to those of A_* by the rule

$$\lambda_{k,\ell}(A) = 1 + \mu \lambda_{k,\ell}(A_*) \stackrel{\text{Prop. 1.12}}{=} 1 - 4\mu \left(\sin^2 \frac{\pi k h}{2} + \sin^2 \frac{\pi \ell h}{2} \right).$$

Consequently,

$$\sup_{h>0} \rho(A) = \max\{1, |1 - 8\mu|\}, \quad \text{hence} \quad \mu \leq \frac{1}{4} \Leftrightarrow \text{stability}.$$

Method 2.26 (Fourier analysis) Fourier analysis generalizes to two dimensions: of course, we now need to extend the range of (x, y) in (2.11) from $0 \leq x, y \leq 1$ to $x, y \in \mathbb{R}$. A 2D Fourier transform reads

$$\hat{u}(\theta, \psi) = \sum_{\ell, m \in \mathbb{Z}} u_{\ell, m} e^{-i(\ell\theta + m\psi)}$$

and all our results readily generalize. In particular, the Fourier transform is an isometry from $\ell_2[\mathbb{Z}^2]$ to $L_2([-\pi, \pi]^2)$, i.e.

$$\left(\sum_{\ell, m \in \mathbb{Z}} |u_{\ell, m}|^2 \right)^{1/2} =: \|\mathbf{u}\| = \|\hat{u}\|_* := \left(\frac{1}{4\pi^2} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} |\hat{u}(\theta, \psi)|^2 d\theta d\psi \right)^{1/2},$$

and the method is stable iff $|H(\theta, \psi)| \leq 1$ for all $\theta, \psi \in [-\pi, \pi]$. The proofs are an easy elaboration on the one-dimensional theory. Insofar as the Euler method (2.13) is concerned,

$$H(\theta, \psi) = 1 + \mu (e^{-i\theta} + e^{i\theta} + e^{-i\psi} + e^{i\psi} - 4) = 1 - 4\mu \left(\sin^2 \frac{\theta}{2} + \sin^2 \frac{\psi}{2} \right),$$

and we again deduce stability if and only if $\mu \leq \frac{1}{4}$.

Method 2.27 (Crank-Nicolson for 2D) Applying the trapezoidal rule to our semi-discretization (2.12) we obtain the two-dimensional Crank-Nicolson method:

$$(I - \frac{1}{2}\mu A_*) \mathbf{u}^{n+1} = (I + \frac{1}{2}\mu A_*) \mathbf{u}^n, \quad (2.14)$$

in which we move from the n -th to the $(n+1)$ -st level by solving the system of linear equations $B\mathbf{u}^{n+1} = C\mathbf{u}^n$, or $\mathbf{u}^{n+1} = B^{-1}C\mathbf{u}^n$. For stability, similarly to the one-dimensional case, the eigenvalue analysis implies that $A = B^{-1}C$ is normal and shares the same eigenvectors with B and C , hence

$$\lambda(A) = \frac{\lambda(C)}{\lambda(B)} = \frac{1 + \frac{1}{2}\mu\lambda(A_*)}{1 - \frac{1}{2}\mu\lambda(A_*)} \Rightarrow |\lambda(A)| < 1 \text{ as } \lambda(A_*) < 0$$

and the method is stable for all μ . The same result can be obtained through the Fourier analysis.

Implementing the Crank-Nicolson method requires solving the linear system $B\mathbf{u}^{n+1} = C\mathbf{u}^n$ at each step. The matrix $B = I - \frac{1}{2}\mu A_*$ has a structure similar to that of A_* , so we may apply the fast Poisson solver seen in Lectures 3 and 4. The total computational cost per iteration is $\mathcal{O}(M^2 \log M)$ for a $M \times M$ discretization grid.

Matlab demo: Download the Matlab GUI for *Solving the Wave and Diffusion Equations in 2D* from http://www.damtp.cam.ac.uk/user/hf323/M21-II-NA/demos/pdes_2d/pdes_2d.html and solve the diffusion equation (2.11) for different initial conditions. For the numerical solution of the equation you can choose from the Euler method and the Crank-Nicolson scheme. The GUI allows you to solve the wave equation as well. Compare the behaviour of solutions!