

Mathematical Tripos Part II: Michaelmas Term 2022

Numerical Analysis – Lecture 16

Definition 4.10 (Strictly diagonally dominant matrices) A matrix A is called strictly diagonally dominant by rows (resp. by columns) if

$$|a_{ii}| > \sum_{j \neq i} |a_{ij}|, \quad i = 1..n \quad (\text{resp. } |a_{jj}| > \sum_{i \neq j} |a_{ij}|, \quad j = 1..n).$$

From Gershgorin theorem, it follows that strictly diagonally dominant matrices are nonsingular.

Theorem 4.11 *If A is strictly diagonally dominant (either by rows or columns), then both the Jacobi and the Gauss-Seidel methods converge.*

Proof. Jacobi method: We have $H_J = -D^{-1}(A - D) = I - D^{-1}A$. The diagonal elements of H_J are all zero, and the sum of the off-diagonal entries on the i 'th row is $\sum_{j \neq i} |(H_J)_{ij}| = \sum_{j \neq i} |A_{ij}|/|A_{ii}| < 1$ if A is strictly diagonally dominant by rows. Applying Gershgorin's theorem to H_J , we get that all the eigenvalues of H_J have modulus < 1 , which is what we wanted. If A is strictly diagonally dominant by columns (instead of by rows), we get that $\rho(I - AD^{-1}) < 1$ using the same argument, and use the fact that $I - D^{-1}A$ and $I - AD^{-1}$ have the same eigenvalues (since $I - D^{-1}A = D^{-1}(I - AD^{-1})D$).

Gauss-Seidel: If λ is an eigenvalue of $H_{GS} = -(L_0 + D)^{-1}U_0$, then this means that $H_{GS} - \lambda I = -(L_0 + D)^{-1}U_0 - \lambda I$ is singular, which in turn implies that $U_0 + \lambda(L_0 + D)$ is singular. It is easy to see that if $A = L_0 + D + U_0$ is strictly diagonally dominant, then the same is true for $A_\lambda = U_0 + \lambda(L_0 + D)$ for all $|\lambda| \geq 1$, and in particular A_λ is nonsingular in this case. This implies that any eigenvalue λ of $-(L_0 + D)^{-1}U_0$ must satisfy $|\lambda| < 1$. This shows convergence of the Gauss-Seidel method. (Note: a similar argument can also be used for Jacobi.) \square

Theorem 4.12 (The Householder–John theorem) *If A and B are real matrices such that both A and $A - B - B^T$ are symmetric positive definite, then the spectral radius of $H = -(A - B)^{-1}B$ is strictly less than one.*

Proof. Let λ be an eigenvalue of H , so $Hw = \lambda w$ holds, where $w \neq 0$ is an eigenvector. (Note that both λ and w may have nonzero imaginary parts when H is not symmetric, e.g. in the Gauss–Seidel method.) By definition of H we have $-Bw = \lambda(A - B)w$, and we note that $\lambda \neq 1$ since otherwise A would be singular (which it is not). Thus, we deduce

$$\bar{w}^T B w = \frac{\lambda}{\lambda - 1} \bar{w}^T A w, \quad (4.3)$$

where the bar means complex conjugation. Moreover, writing $w = u + iv$, where u and v are real, we find (for $C = C^T$) the identity $\bar{w}^T C w = u^T C u + v^T C v$, so symmetric positive definiteness in the assumption implies $\bar{w}^T A w > 0$ and $\bar{w}^T (A - B - B^T) w > 0$. In the latter inequality, we use relation (4.3) and its conjugate transpose to obtain

$$0 < \bar{w}^T A w - \bar{w}^T B w - \bar{w}^T B^T w = \left(1 - \frac{\lambda}{\lambda - 1} - \frac{\bar{\lambda}}{\bar{\lambda} - 1}\right) \bar{w}^T A w = \frac{1 - |\lambda|^2}{|\lambda - 1|^2} \bar{w}^T A w.$$

Now $\lambda \neq 1$ implies $|\lambda - 1|^2 > 0$. Hence, recalling that $\bar{w}^T A w > 0$, we see that $1 - |\lambda|^2$ is positive. Therefore every eigenvalue of H satisfies $|\lambda| < 1$ as required. \square

Corollary 4.13 1) *If A is symmetric positive definite, then the Gauss-Seidel method converges.*
2) *If both A and $2D - A$ are symmetric positive definite, then the Jacobi method converges.*

Proof. 1) For the Gauss-Seidel method, B is the superdiagonal part of symmetric A , hence $A - B - B^T$ is equal to D , the diagonal part of A , and if A is positive definite, then D is positive definite too (this is the first part of the Exercise 23 from Example Sheets).

2) For the Jacobi method, we have $B = A - D$, and if A is symmetric, then $A - B - B^T = 2D - A$. (The latter matrix is the same as A except that the signs of the off-diagonal elements are reversed.) \square

Example 4.14 (Poisson's equation on a square) As we have seen in the previous sections linear systems $A\mathbf{x} = \mathbf{b}$, where A is a real symmetric positive (negative) definite matrix, frequently occur in numerical methods for solving elliptic partial differential equations. A typical example we already encountered is Poisson's equation on a square where the *five-point formula* approximation yields an $n \times n$ system of linear equations with $n = m^2$ unknowns $u_{p,q}$:

$$u_{p-1,q} + u_{p+1,q} + u_{p,q-1} + u_{p,q+1} - 4u_{p,q} = h^2 f(ph, qh) \quad (4.4)$$

(Note that when p or q is equal to 1 or m , then the values $u_{0,q}$, $u_{p,0}$ or $u_{p,m+1}$, $u_{m+1,q}$ are known boundary values and they should be moved to the right-hand side, thus leaving fewer unknowns on the left.)

For any ordering of the grid points (ph, qh) we have shown in Lemma 1.11 that the matrix A of this linear system is symmetric and negative definite.

Corollary 4.15 For linear system (4.4), for any ordering of the grid, both Jacobi and Gauss-Seidel methods converge.

Proof. By Lemma 1.11, A is symmetric and negative definite, hence convergence of Gauss-Seidel. To prove convergence of the Jacobi method, we need negative definiteness of the matrix $2D - A$, and that follows by the same arguments as in Lemma 1.11: recall that the proof operates with the modulus of the off-diagonal elements and does not depend on their sign. \square

Method 4.16 (Relaxation) It is often possible to improve the efficiency of the recursive schemes above by *relaxation*. Specifically, instead of letting $\mathbf{x}^{(k+1)} = H\mathbf{x}^{(k)} + \mathbf{v}$, we let

$$\begin{aligned} \hat{\mathbf{x}}^{(k+1)} &= H\mathbf{x}^{(k)} + \mathbf{v}, \quad \text{and then} \quad \mathbf{x}^{(k+1)} = \omega\hat{\mathbf{x}}^{(k+1)} + (1-\omega)\mathbf{x}^{(k)} \\ &= H_\omega\mathbf{x}^{(k)} + \omega\mathbf{v} \end{aligned}$$

with

$$H_\omega = \omega H + (1-\omega)I,$$

where ω is a real constant called the *relaxation parameter*. (Note that $\omega = 1$ corresponds to the standard "unrelaxed" iteration.) Good choice of ω leads to a smaller spectral radius of the iteration matrix (compared with the "unrelaxed" method), and the smaller the spectral radius, the faster the iteration converges.

The eigenvalues of H_ω and H are related by the rule $\lambda_\omega = \omega\lambda + (1-\omega)$, therefore one may try to choose $\omega \in \mathbb{R}$ to minimize

$$\rho(H_\omega) = \max \{ |\omega\lambda + (1-\omega)| : \lambda \in \sigma(H) \}.$$

In general, $\sigma(H)$ is unknown, but often we have some information about it which can be utilized to find a "good" (rather than "best") value of ω . For example, suppose that it is known that $\sigma(H)$ is real and resides in the interval $[\alpha, \beta]$ where $-1 < \alpha < \beta < 1$. In that case we seek ω to minimize

$$\max \{ |\omega\lambda + (1-\omega)| : \lambda \in [\alpha, \beta] \}.$$

It is readily seen that, for a fixed $\lambda < 1$, the function $f(\omega) = \omega\lambda + (1-\omega)$ is decreasing, therefore, as ω increases (decreases) from 1 the spectrum of H_ω moves to the left (to the right) of the spectrum of H . It is clear that the optimal location of the spectrum $\sigma(H_\omega)$ (or of the interval $[\alpha_\omega, \beta_\omega]$ that contains $\sigma(H_\omega)$) is the one which is centralized around the origin:

$$-[\omega\alpha + (1-\omega)] = \omega\beta + (1-\omega) \Rightarrow \omega_{\text{opt}} = \frac{2}{2-(\alpha+\beta)}, \quad -\alpha_{\omega_{\text{opt}}} = \beta_{\omega_{\text{opt}}} = \frac{\beta-\alpha}{2-(\alpha+\beta)}.$$