

## 15 Sums of squares on the hypercube

In this lecture we look at polynomial optimisation on the hypercube  $S = \{-1, 1\}^n$ . One way to certify that a polynomial  $f$  is nonnegative on  $\{-1, 1\}^n$  is to try to express it in the following way:

$$f(x) = \sum_{i=1}^l q_i(x)^2 + \sum_{i=1}^n (x_i^2 - 1)h_i(x) \quad (1)$$

where  $q_i$  and  $h_i$  are arbitrary polynomials. It is clear that any  $f$  of the form (1) is nonnegative on  $\{-1, 1\}^n$ . For example consider the function  $f(x) = 1 + x_1$ . Clearly  $f$  is nonnegative on  $\{-1, 1\}^n$  and one can verify that we have the following certificate of nonnegativity  $1 + x_1 = \frac{1}{2}(1 + x_1)^2 + (x_1^2 - 1) \cdot (-1/2)$ .

Functions on the hypercube can be expressed in a specific basis, called the basis of *square-free monomials* (also known as *multilinear monomials*). A square-free monomial is a monomial of the form  $x^S := \prod_{i \in S} x_i$  where  $S \subseteq [n]$  (we use the notation  $[n] := \{1, \dots, n\}$ ).

**Proposition 15.1.** *Any function  $f : \{-1, 1\}^n \rightarrow \mathbb{R}$  can be expressed as*

$$f(x) = \sum_{S \subseteq [n]} f_S x^S \quad \forall x \in \{-1, 1\}^n \quad (2)$$

for some coefficients  $(f_S)_{S \subseteq [n]}$ .

*Proof.* Let  $a \in \{-1, 1\}^n$  and let  $\delta_a(x)$  be the function that takes value 1 at  $a$  and 0 elsewhere. Note that  $\delta_a$  can be expressed as:

$$\frac{1}{2^n} \prod_{i=1}^n (1 + a_i x_i). \quad (3)$$

Expanding the product we see that  $\delta_a$  is a linear combination of the square-free monomials. Finally since each function is a linear combination of the  $\delta_a$ s we get the desired result.  $\square$

**Definition 15.1.** We say that a function  $f : \{-1, 1\}^n \rightarrow \mathbb{R}$  is *k-sos on  $\{-1, 1\}^n$*  if it is a sum-of-squares of polynomials of degree at most  $k$  on  $\{-1, 1\}^n$ , i.e., if there exists polynomials  $q_1, \dots, q_l$  of degree at most  $k$  such that  $f(x) = \sum_{i=1}^l q_i(x)^2$  for all  $x \in \{-1, 1\}^n$ .

**Remark 1.** One can show (using e.g., the division algorithm for polynomials in more than one variable) that  $f$  is *k-sos on  $\{-1, 1\}^n$*  if and only if it can be expressed as (1) where  $\deg q_i \leq k$  for all  $i = 1, \dots, l$  and  $\deg h_i \leq 2k - 2$  for all  $i = 1, \dots, n$  (assuming  $\deg f \leq 2k$ ).

**Example 15.1.** • The function  $f(x) = 1 + x_1$  is 1-sos on  $\{-1, 1\}^n$  because  $1 + x_1 = \frac{1}{2}(1 + x_1)^2$  on  $\{-1, 1\}^n$ .

- Any nonnegative function  $f$  on  $\{-1, 1\}^n$  is *n-sos*. Indeed we have  $f = g^2$  on  $\{-1, 1\}^n$  where  $g : \{-1, 1\}^n \rightarrow \mathbb{R}$  is defined by  $g(x) = \sqrt{f(x)}$ . By Proposition 15.1 we know that  $g$  is a polynomial of degree at most  $n$ . Another way of seeing this same fact is to observe that the delta function  $\delta_a$  defined in (3) satisfies  $\delta_a = \delta_a^2$  and so we have:

$$f = \sum_{a \in \{-1, 1\}^n} f(a) \delta_a^2 \quad (4)$$

Since  $f(a) \geq 0$  and each  $\delta_a$  is a polynomial of degree at most  $n$  (cf. (3)), (4) shows that  $f$  is  $n$ -sos.

*Degree cancellations:* There is an important difference that one must keep in mind between (i) sum-of-squares certificates on the hypercube, and (ii) global sum-of-squares certificates. When writing a global sum of squares certificate for a polynomial  $f$  on  $\mathbb{R}^n$ , i.e.,  $f(x) = \sum_{i=1}^l q_i(x)^2$  for all  $x \in \mathbb{R}^n$  then necessarily  $\deg q_i \leq (\deg f)/2$ . When working on  $\{-1, 1\}^n$  however, such degree bounds on the  $q_i$ 's do not hold anymore as there can be *degree cancellations*. This is already evident in the two examples above (Example 15.1).

The next theorem shows that deciding whether a function  $f : \{-1, 1\}^n \rightarrow \mathbb{R}$  is  $k$ -sos is a semidefinite feasibility problem.

**Theorem 15.1.** *A function  $f : \{-1, 1\}^n \rightarrow \mathbb{R}$  is  $k$ -sos on  $\{-1, 1\}^n$  if and only if there exists a positive semidefinite matrix  $Q$  of size  $\binom{n}{0} + \binom{n}{1} + \dots + \binom{n}{k}$  such that*

$$f_S = \sum_{\substack{U, V \subseteq [n] \\ |U|, |V| \leq k \\ U \Delta V = S}} Q_{U, V} \quad \forall S \subseteq [n], |S| \leq 2k$$

where  $f_S$  is the coefficient of  $f$  in the expansion (2), and  $U \Delta V$  is the symmetric difference of  $U$  and  $V$ , i.e.,  $U \Delta V = (U \setminus V) \cup (V \setminus U)$ .

*Proof.* The proof is very similar to Theorem 13.2. Simply use the fact that  $x^U x^V = x^{U \Delta V}$  on  $\{-1, 1\}^n$ .  $\square$