

Project 2 - Stability of stratified shear flow

I. BACKGROUND

Although most natural flows are time-varying, and have complicated spatial structure, important insights can be gained by examining the stability of simple flows. For example, many identifiable features in the atmosphere and ocean, such as eddies and billow clouds (shown), are generated and derive their properties from fluid instabilities.



FIG. 1: Kelvin-Helmholtz billows developing in a cloud layer over Mount Shasta, California. Photo ©1999 Beverly Shannon.

II. INTRODUCTION

Here, we will examine the basic stability properties of a stratified shear flow, and will then use Diabolo to examine the nonlinear evolution of the unstable state.

Start by considering a stratified shear flow of the form:

$$\mathbf{U} = S_0 h \tanh\left(\frac{y - LY/2}{h}\right) \hat{i}, \quad (1)$$

$$B = N_0^2 h \tanh\left(\frac{y - LY/2}{h}\right), \quad (2)$$

where $B = -g\rho/\rho_0$ is the buoyancy, h is the height of the shear layer, and S_0 and N_0 are the shear and buoyancy frequency at the center of the shear layer. Pick some parameters that permit shear instability by the Miles-Howard theorem (with $Ri = N^2/S^2 < 1/4$ somewhere in the flow). For example, $LY = 1$, $h = 1/10$, $S_0 = 10$, and $N_0^2 = 10$ seem to work well.

III. LINEAR STABILITY ANALYSIS

Consider the stability of small perturbations the base state defined above: $\mathbf{u} = \mathbf{U} + \epsilon \mathbf{u}'$, $b = B + \epsilon b'$. We can then look for normal mode solutions to the linearised equations of the form

$$v' = \text{Re} [\hat{v}(y) \exp(\sigma t + i(kx + lz))] . \quad (3)$$

The MATLAB script, *linstab.m*, which you can find inside the folder *project2*, solves the viscous linear stability problem for stratified shear flow, returning the vertical velocity and buoyancy eigenfunctions, $\hat{v}(y)$ and $\hat{b}(y)$, and the corresponding growth rates, σ . Specifically, the code solves the following equations for 2D perturbations ($l = 0$):

$$\sigma(\hat{v}_{yy} - k^2 \hat{v}) = -ikU(y)\hat{v}_{yy} + ikU_{yy}\hat{v} + \nu(d_y^2 - k^2)^2 \hat{v} - k^2 \hat{b}, \quad (4)$$

$$\sigma \hat{b} = -B_y \hat{v} - ikU(y)\hat{b} + \kappa(d_y^2 - k^2)\hat{b}. \quad (5)$$

At the start of *linstab.m*, you can specify several parameters including the vertical domain size and number of gridpoints, h , and ν and κ . Optionally, edit these parameters to match your choice of parameters (the suggested ones listed above are in place now.)

Run the script *linstab.m* from within MATLAB, which will then plot the growth rate of the most unstable mode as a function of the horizontal wavenumber. How does the most unstable mode vary with the shear layer width, and the viscosity?

IV. NONLINEAR SIMULATIONS

Once you are happy with a choice of parameters, try to recreate the instability using Diablo. We will use periodic boundary conditions at both ends of the x -domain for all variables, and place free-slip, no normal flow walls at the top and bottom in the y -direction with gravity pointing down (in the negative y -direction). If we want to capture the most unstable mode, we need it to fit within our domain. Based on the results of your stability calculation from *linstab.m*, set the size of the domain in the x -direction to be a multiple (say 3) of the wavelength of the most unstable mode. Using the size of the most unstable mode to set the size of the domain ensures that we can capture that mode inside our domain with periodic boundary conditions.

Edit the following Diablo setup scripts:

set_params.m - A Reynolds number of 500 works reasonably well here. Match LY , the domain size in the y -direction with what you used in *linstab.m*. Start with $NY = 100$, and set NX so that the grid is close to isotropic $\Delta x \simeq \Delta y$ (within a factor of two should be fine).

set_bcs.m - Set up periodic boundary conditions at both ends of the x - *domain*. In the y -direction, let $U_2=0$ at the edges (no normal flow), and let $dU_1/dy=0$, $dP/dy=0$, and $dTH/dy=0$.

create_grid.m - Since most of the action will happen near the center of the domain, it is useful to cluster the grid in the y -direction to put more points there. Change `GRID_TYPE_Y=2` to do this, and set the stretching factor to 1.0

(higher numbers will lead to more stretching.).

create_flow.m - Here, in the User input area, create an initial flow with a horizontal velocity (U1) and buoyancy (TH(:,1)) matching the profiles that you used in *linstab.m*, and uniform in the x -direction. Keep the random perturbation added to the velocity field (you can play around with this later).

Now, run Diablo, and see what happens. Do Kelvin-Helmholtz billows develop with a wavelength close to what you predicted? If not, what might be happening? (Hint: you might need to change the random perturbation added to the velocity field in *create_flow.m* to make sure that you are seeding all horizontal wavenumbers with the same amount of energy.)

Plot the *rms* and mean velocity and buoyancy as a function of y and t . Can you identify an exponential growth phase at the beginning of the simulation? Calculate and plot the gradient Richardson number: $Ri_g = \frac{\langle \partial b / \partial y \rangle}{(\langle \partial u / \partial y \rangle)^2}$, where angle brackets represent an average over x . How does the minimum value of Ri_g evolve as a function of time? What does this say about the stability of the shear layer after the development of K-H billows and the resulting mixing?

V. REFERENCES

Several papers that might be useful for this project: Peltier and Caulfield¹ and Smyth and Moum² contain nice discussions of mixing in stratified shear flows. The appendix in Smyth et al.³ describes the setup of the linear stability analysis in more detail. Copies of these papers are in the project folder.

¹ W. Peltier and C. Caulfield, Annual review of fluid mechanics **35**, 135 (2003).

² W. Smyth and J. Moum, Oceanography **25**, 140 (2012).

³ W. Smyth, J. Moum, and J. Nash, Journal of Physical Oceanography **41**, 412 (2011).