


The projective to Einstein correspondence and a kink on a wormhole



Alice Claire Waterhouse

Supervisor: Dr. Maciej Dunajski

Department of Applied Mathematics and Theoretical Physics
University of Cambridge

This dissertation is submitted for the degree of
Doctor of Philosophy

Queens' College

November 2019

Abstract

The first part of this thesis is concerned with ϕ^4 field theory on a wormhole spacetime in $3 + 1$ dimensions. This spacetime has two asymptotically flat ends connected by a spherical throat of radius a . We show that the theory possesses a kink solution which is linearly stable, and compare its discrete spectrum to that of the ϕ^4 kink on $\mathbb{R}^{1,1}$. We present some results on the non-linear resonant coupling between the discrete and continuous spectra in the range of a where there is exactly one discrete mode.

The second part of the thesis is based on recent work by Dunajski and Mettler. They show that a class of neutral signature Einstein manifolds M can be canonically constructed as rank n affine bundles over projective structures in dimension n . These have the same symmetry group as the underlying projective manifold, and are also endowed with a natural symplectic form, which is related to the metric by an endomorphism of the tangent bundle that squares to the identity. Consequently, they carry an almost para-Kähler structure.

We show that every metric within the class is a Kaluza–Klein reduction of an Einstein metric on an \mathbb{R}^* bundle over M . We also show that the structures are para- c -projectively compact in the sense of Čap–Gover, and interpret the compactification in terms of the tractor bundle of the projective structure.

In dimension four, the manifolds M have anti-self-dual conformal curvature, and are thus associated with a twistor space. In the presence of a symmetry, they can be reduced to Einstein–Weyl structures in dimension three via the Jones–Tod correspondence. Because M is also Einstein with non-zero scalar curvature, these Einstein–Weyl structures are determined by solutions of the $SU(\infty)$ –Toda equation.

We investigate the Einstein–Weyl structures which can be obtained in this way in terms of the symmetry group of the underlying projective surface. Several examples are considered in detail, resulting in new, explicit solutions of the $SU(\infty)$ –Toda equation. We focus in particular on the case where the projective structure is \mathbb{RP}^n , additionally describing the Jones–Tod reduction from the twistor perspective.

Declaration

I hereby declare that except where specific reference is made to the work of others, the contents of this dissertation are original and have not been submitted in whole or in part for consideration for any other degree or qualification in this, or any other university. This dissertation is my own work and contains nothing which is the outcome of work done in collaboration with others, except as specified in the text.

Alice Claire Waterhouse
November 2019

Acknowledgements

The work in this thesis would not have been possible without the ideas, enthusiasm and sense of humour of my supervisor, Maciej Dunajski. I offer him my sincere thanks for his patience and guidance over the last three years. Additionally, I would like to thank Rod Gover, Piotr Bizoń, Michal Kahl, Nick Manton, Thomas Mettler, Paul Tod and Paul Townsend for interesting discussions and helpful suggestions.

Thanks are also due to the Japesters for Japes; to Jonny, Philip and Chris H for always making B2.12 a good place to be; to Jack for a fantastic adventure; and to the rest of the pav B gang for clarifying the technical difference between a fruit and a vegetable.

I express my deepest gratitude to my parents, Angie and Tim, for their sturdy and ever-present support; to Esther, for making me the person I am today; and to Cherish, without whom I may not have made it to this point in my PhD. Finally, I would like to thank Chris, for laughs, companionship and his unending loyalty.

Table of contents

List of figures	xiii	
1	Introduction	1
2	The ϕ^4 kink on a wormhole spacetime	5
2.1	Background: the ϕ^4 kink on $\mathbb{R}^{1,1}$	5
2.1.1	Topological stability and the kink solution	6
2.1.2	Linear stability	6
2.1.3	Asymptotic stability	8
2.1.4	Derrick's scaling argument	8
2.2	The static kink on a wormhole	9
2.2.1	Large a limit	10
2.3	Linearised perturbations around the kink	13
2.3.1	Finding internal modes numerically	15
2.3.2	Large a limit	16
2.3.3	Critical values of a	18
2.4	Resonant coupling of the internal modes to the continuous spectrum . .	20
2.4.1	Conjectured decay rate in the presence of a single internal mode	21
2.4.2	Numerical investigation of the conjectured decay rate	26
2.4.3	Expected decay rates in other regimes	28
3	The projective to Einstein correspondence $N \rightarrow M$	31
3.1	Projective Geometry	32
3.1.1	The Cartan bundle	35
3.1.2	Tractor bundles	36
3.2	The projective to Einstein correspondence	40
3.2.1	Symmetries of M	42
3.2.2	Tractor perspective	43

3.2.3	The model case	44
4	An Einstein metric on an \mathbb{R}^* bundle over M	45
4.1	The model case	45
4.2	The general case	47
4.2.1	Ricci scalar of g_Λ and divergence of Ω_Λ	50
5	Einstein–Weyl structures and $SU(\infty)$–Toda fields	55
5.1	Background	56
5.1.1	Anti–self–duality, spinors and totally null distributions	56
5.1.2	Einstein–Weyl structures	61
5.1.3	The $SU(\infty)$ –Toda equation	63
5.1.4	Projective structures on a surface	64
5.1.5	From projective surfaces to $SU(\infty)$ –Toda fields	65
5.2	The most general case	66
5.2.1	Solution to the $SU(\infty)$ –Toda equation	67
5.2.2	Two monopoles	69
5.3	The submaximally symmetric case	69
5.4	The model case	73
5.4.1	Conformal Structure on M	73
5.4.2	Twistor space of M	74
5.4.3	Pseudo–hyper–Hermitian structure on M	76
5.4.4	A line bundle over the twistor space of M	78
5.4.5	Factoring the model to Einstein–Weyl	79
5.4.6	An example of the mini–twistor correspondence	81
6	Para–c–projective compactification of M	83
6.1	Background and definitions	83
6.1.1	Almost (para–)complex geometry	83
6.1.2	Almost (para–)CR structures and contact distributions	85
6.1.3	Connections and (para–) c –projective equivalence	86
6.1.4	Para– c –projective compactification	87
6.2	Compactifying the Dunajski–Mettler Class	88
6.2.1	Tractor construction of M	89
6.2.2	The compactification theorem	91
6.3	An alternative approach to Theorem 6.2.4	96
6.3.1	The model case	97

7 Concluding Remarks	101
-----------------------------	------------

References	103
-------------------	------------

List of figures

2.1	The kink solution for several values of a , along with the flat kink $\Phi_0(r)$.	11
2.2	The order ϵ^2 perturbation to the static kink on $\mathbb{R}^{1,1}$	13
2.3	The potential of the 1-dimensional quantum mechanics problem arising from the study of stability of the soliton for values of a between $a = 10$ and $a = 0.3$	14
2.4	Even bound states of the potential $V_0(r)$ and of the potential $V(r)$ for two different values of a	16
2.5	Odd bound states of the potential $V_0(r)$ and of the potential $V(r)$ for two different values of a	17
2.6	The frequencies of the internal modes of the kink plotted against the wormhole radius a	17
2.7	A comparison of the predicted and numerical calculations for the energy of the zero mode as a function of ϵ^2 for small ϵ	19
2.8	A comparison of the predicted and numerical calculations for the energy of the odd vibrational mode as a function of ϵ^2 for small ϵ	19
2.9	The decay of internal mode oscillations for various initial conditions when $a = 0.5$	27

Chapter 1

Introduction

This thesis splits naturally into two parts. In Chapter 2, we discuss ϕ^4 field theory on a wormhole spacetime with two asymptotically flat ends. This theory has two distinct vacua, and permits static, finite energy “kink” solutions which interpolate between the two vacua as one moves from one asymptotically flat end to the other. We study the stability of such solutions from various perspectives, with explicit comparison to the kinks arising in ϕ^4 theory on $\mathbb{R}^{1,1}$.

The remainder of the thesis is concerned with a class of Einstein manifolds which can be canonically constructed from projective structures as shown in recent work by Dunajski and Mettler [28]. A projective structure on a manifold specifies a preferred curve in every direction at every point. It can be understood as an equivalence class of affine connections which share the same geodesics up to parametrisation. Because it is defined at the level of the connection rather than a first order structure such as a metric, it is an intrinsically second order object.

Given a projective structure on a manifold N of dimension n , Dunajski and Mettler [28] canonically construct a neutral signature Einstein metric g with non-zero scalar curvature on a certain rank n affine bundle $M \rightarrow N$. They thus convert a second order object into a first order object on a larger manifold. The $2n$ -dimensional space M also carries a natural symplectic form Ω , and an endomorphism $J : TM \rightarrow TM$ which is such that J^2 is the identity and $g(\cdot, \cdot) = \Omega(\cdot, J\cdot)$. This makes (M, g, Ω) a so-called *almost para-Kähler* structure. We review their construction in Chapter 3.

In Chapter 4 we show that g arises as the Kaluza–Klein reduction of an Einstein metric \hat{g} on an \mathbb{R}^* bundle $\kappa_{\mathcal{Q}} : \mathcal{Q} \rightarrow M$ which has curvature form $\kappa_{\mathcal{Q}}^*(\Omega)$. We will construct \hat{g} explicitly, and give an interpretation of the manifold (\mathcal{Q}, \hat{g}) in terms of the projective geometry on N .

The work in Chapter 5 is based on the fact that for $n = 2$ (so that M has dimension four), the conformal curvature of g is anti-self-dual. Recall that the Hodge operator \star defined by a Euclidean or neutral signature metric in four dimensions is an involution on two-forms (i.e. squares to the identity). It thus has eigenvalues ± 1 , and the space of two-forms splits into the corresponding eigenspaces, which are referred to as self-dual (SD) or anti-self-dual (ASD) respectively. Due to its index symmetries, the Weyl conformal curvature tensor can be thought of as a map from two-forms to two-forms, and therefore has a corresponding decomposition. Since the Weyl tensor encodes the conformal curvature, we say that a conformal or (pseudo-)Riemannian manifold whose Weyl tensor is ASD is equipped with an *ASD conformal structure*.

The field equations corresponding to anti-self-duality of the Weyl tensor in four dimensions can be solved by a twistor construction, and are thus *integrable* [77]. This means that any systems of differential equations which can be obtained from them by symmetry reduction should also be integrable (see [55] for a review). In particular, the class of dispersionless integrable systems in three dimensions arise in this way. The construction [28] provides some examples of ASD conformal structures in neutral signature which, in the presence of a (non-null) symmetry, give rise to solutions of an integrable system called the $SU(\infty)$ -Toda field equation via $(2+1)$ -dimensional Einstein-Weyl structures. In Chapter 5 we discuss the extraction of $SU(\infty)$ -Toda fields obtainable in this way.

In Chapter 6 we return to projective structures of any dimension, and show that the structure (M, g, Ω) can be thought of as compactifiable in a certain sense. Recall that a (pseudo-)Riemannian manifold (M, g) is said to be *conformally compact* if there is a smooth positive function T such that $T^2 g$ smoothly extends to a manifold with boundary $\overline{M} = M \cup \partial M$, and the set $\{m \in \overline{M} : T(m) = 0\}$ is a hypersurface which coincides with the boundary ∂M . This is a useful concept because $(M, T^2 g)$ has the same conformal structure, and hence the same *causal* structure, as (M, g) . It has been used to study causal structures in both general relativity [61] and quantum field theory [83]. It is also useful for formulating the boundary conditions of conformally invariant field equations such as those arising in Yang-Mills theory [75].

Recent work by Čap and Gover [17, 18] has generalised this idea to other geometrical structures which admit some weakening that extends to a manifold with boundary. One example of such a structure is an *almost complex* manifold (M, J) , that is, a manifold carrying a smooth endomorphism J of the tangent bundle which squares to $-Id$. If (M, J) carries a connection ∇ which is correctly adapted to J , one can define the c -projective equivalence class $[\nabla]$ to which ∇ belongs, and ask whether the

c –projective structure $(M, J, [\nabla])$ extends to a manifold with boundary \overline{M} [18]. The main goal of Chapter 6 is to adapt the work of [18] to the *para*– c –projective case, where J instead squares to Id , and to show that the endomorphisms J on the manifolds (M, g, Ω) arising in the projective to Einstein correspondence have correctly adapted connections which admit so–called *para*– c –projective compactification. The result of this is that the manifolds (M, g, Ω) can be thought of as *para*– c –projectively compact.

Chapter 2 has its own notation and conventions; notation and conventions for Chapters 3 to 6 will be introduced at the beginning of Chapter 3.

Chapter 2

The ϕ^4 kink on a wormhole spacetime

The soliton resolution conjecture [71] states that solutions to solitonic equations with generic initial data should, after some non-linear behaviour, eventually resolve into a finite number of solitons plus a radiative term. This conjecture is intimately tied to soliton stability, which has been investigated for a number of solitonic equations, including that of ϕ^4 theory on $\mathbb{R}^{1,1}$. In this chapter, we study a modification of this theory on a $3 + 1$ dimensional wormhole spacetime which has a spherical throat of radius a , with a focus on the stability properties of the modified kink. In particular, we prove that the modified kink is linearly stable, and compare its discrete spectrum to that of the ϕ^4 kink on $\mathbb{R}^{1,1}$. We also study the resonant coupling between the discrete modes and the continuous spectrum for small but non-linear perturbations. Some numerical and analytical evidence for asymptotic stability is presented for the range of a where the kink has exactly one discrete mode. This chapter is almost identical to the preprint [80].

2.1 Background: the ϕ^4 kink on $\mathbb{R}^{1,1}$

One dimensional ϕ^4 theory is well-documented in the literature (see for example [53]). The aim of this section is to introduce some notation and some ideas about stability which will be useful when we come to consider the modified theory.

2.1.1 Topological stability and the kink solution

The action takes the form

$$S = \int_{\mathbb{R}^2} \left(\frac{1}{2} \eta^{ab} \partial_a \phi \partial_b \phi + \frac{1}{2} (1 - \phi^2)^2 \right) dx dt,$$

where $x^a = (t, x)$ are coordinates on $\mathbb{R}^{1,1}$ and η^{ab} is the Minkowski metric with signature $(-, +)$. Note that it has two vacua, given by $\phi = \pm 1$. Finiteness of the associated conserved energy

$$E = \int_{\mathbb{R}} \left(\frac{1}{2} (\phi_t)^2 + \frac{1}{2} (\phi_x)^2 + \frac{1}{2} (1 - \phi^2)^2 \right) dx, \quad (2.1)$$

requires that the field lies in one of these two vacua in the limits $\phi_{\pm} = \lim_{x \rightarrow \pm\infty} [\phi(x)]$. We can thus classify finite energy solutions in terms of their topological charge $N = (\phi_+ - \phi_-)/2$, which takes values in $\{-1, 0, 1\}$.

The equations of motion are

$$\phi_{tt} = \phi_{xx} + 2\phi(1 - \phi^2) \quad (2.2)$$

and we find a static solution $\phi = \tanh(x - c)$ which we call the flat kink. It interpolates between the two vacua and thus has topological charge $N = 1$. The constant of integration c can be thought of as the position of the kink. We will henceforth use Φ_0 to denote the static kink at the origin, that is, $\Phi_0(x) = \tanh(x)$. Since any small departure from $\phi = \pm 1$ in the limits $x \rightarrow \pm\infty$ results in non-convergence of the integral (2.1), no finite energy deformation can affect N . For this reason, we say that the kink is *topologically stable*.

2.1.2 Linear stability

A second notion of stability which will be important to our discussion is linear stability. On discarding non-linear terms, we find that small perturbations $\phi(t, x) = \Phi_0(x) + e^{i\omega_0 t} v_0(x)$ satisfy the Schrödinger equation

$$L_0 v_0 := -v_0'' - 2(1 - 3\Phi_0^2)v_0 = \omega_0^2 v_0. \quad (2.3)$$

The potential $V_0(x) = -2[1 - 3\Phi_0(x)^2]$ exhibits a so-called “mass gap”, meaning that it takes a finite positive value in the limits $x \rightarrow \pm\infty$. In this case, $V_0(\pm\infty) = 4$. For $\omega_0^2 > 4$, (2.3) admits a continuous spectrum of wave-like solutions.

In addition to its continuum states, the Schrödinger operator in (2.3) has two discrete eigenvalues with normalisable solutions given by

$$(v_0(x), \omega_0) = \left(\frac{\sqrt{3}}{2} \operatorname{sech}^2(x), 0 \right) \quad \text{and} \quad (v_0(x), \omega_0) = \left(\frac{\sqrt{3}}{\sqrt{2}} \operatorname{sech}(x) \tanh(x), \sqrt{3} \right), \quad (2.4)$$

where we have chosen the normalisation constant such that $\int_{-\infty}^{\infty} v_0^2(x) dx = 1$.

The first of these is the zero mode of the kink. Its existence is guaranteed by the translation invariance of (2.2), and up to a multiplicative constant it is equal to $\Phi'_0(x)$. Excitation of this state corresponds to performing a Lorentz boost. In the non-relativistic limit, this amounts to replacing $\Phi_0(x)$ with $\Phi_0(x - vt)$ for some $v \ll 1$ [53].

The second normalisable solution, called an *internal* or *discrete* mode, has non-zero frequency ω_0 , and is thus time periodic. In the full non-linear theory, it decays through resonant coupling to the continuous spectrum [52]. This phenomenon is of considerable interest in non-linear PDEs, and was studied in a more general setting in [69]. The corresponding process in the modified theory will be discussed in Section 2.4.

Linear stability of the kink is equivalent to the Schrödinger operator L_0 in (2.3) having no negative eigenvalues, so that linearised perturbations cannot grow exponentially with time. One way to see that the kink is linearly stable is via the Sturm oscillation theorem:

Theorem 2.1.1 (Sturm). *Let L be a differential operator of the form*

$$L = -\frac{d^2}{dx^2} + V(x)$$

on the smooth square integrable functions u on the interval $[0, \infty)$, with the boundary condition $u(0) = 0$ (corresponding to even parity) or $u'(0) = 0$ (corresponding to odd parity). Let ω^2 be an eigenvalue of L with associated eigenfunction $u(x; \omega)$. Then the number of eigenvalues of L (subject to the appropriate boundary conditions) which are strictly below ω^2 is the number of zeros of $u(x; \omega)$ in $(0, \infty)$.

Note that the symmetry of (2.3) under $x \mapsto -x$ means that any solution on the interval $[0, \infty)$ has a corresponding solution on the interval $(-\infty, 0]$, and these solutions can be pieced together to make a smooth solution on $(-\infty, \infty)$ as long as the boundary conditions at $x = 0$ are chosen to ensure parity ± 1 . Thus there is a one-to-one correspondence between solutions on $[0, \infty)$ and solutions on $(-\infty, \infty)$ which are smooth at $x = 0$. Since the eigenfunctions (2.4) have no zeros on the interval $[0, \infty)$, it

follows that there can be no eigenfunctions with $\omega_0^2 < 0$, and thus the kink is linearly stable.

2.1.3 Asymptotic stability

The final notion of stability that we will consider is that of asymptotic stability. Stated simply, asymptotic stability of the kink means that for sufficiently small initial perturbations, solutions of (2.2) will converge locally to $\Phi_0(x)$ or its Lorentz boosted counterpart. This was proved in [45] for odd perturbations, but has not been proved in the general case.

2.1.4 Derrick's scaling argument

Generalisation of the finite energy ϕ^4 kink to higher dimensional Minkowski spacetimes is prohibited by a scaling argument due to Derrick [23]. Suppose $\Phi_n(\mathbf{x})$ is a static, finite energy solution to the equation of motion of the ϕ^4 theory on $\mathbb{R}^{1,n}$. Then it is a minimiser of the (static) energy

$$E(\Phi_n) = \int \left(\nabla \Phi_n(\mathbf{x}) \cdot \nabla \Phi_n(\mathbf{x}) + U(\Phi_n) \right) d^n x =: E_1 + E_2,$$

where we have split E into the two components coming from the two different terms in the integrand. Now consider a spatial rescaling $\mathbf{x} \rightarrow \mu \mathbf{x}$, $\mu > 0$ and define

$$\begin{aligned} e(\mu) &= E(\Phi_n(\mu \mathbf{x})) = \int \left(\nabla(\Phi_n(\mu \mathbf{x})) \cdot \nabla(\Phi_n(\mu \mathbf{x})) + U(\Phi_n(\mu \mathbf{x})) \right) d^d x \\ &= \int \left(\mu^2 \nabla \Phi_n(\mu \mathbf{x}) \cdot \nabla \Phi_n(\mu \mathbf{x}) + U(\Phi_n(\mu \mathbf{x})) \right) d^d x \\ &= \mu^{2-n} E_1 + \mu^{-n} E_2, \end{aligned}$$

where we have obtained the last line by a change of variables from \mathbf{x} to $\mu \mathbf{x}$.

If $\Phi_n(\mathbf{x})$ is a minimiser of E then $\mu = 1$ must also be a stationary point of $e(\mu)$. Evaluating the derivative yields

$$e'(\mu) = \begin{cases} -n\mu^{-n-1} E_2, & \text{if } n = 2 \\ (2-n)\mu^{1-n} E_1 - n\mu^{-n-1} E_2, & \text{otherwise.} \end{cases}$$

Since E_1 , E_2 and μ are all positive, the derivative can only have a zero only when n and $2-n$ have the same sign, which only happens when $n = 1$. We thus conclude that no static, finite energy solutions to the equations of motion exist for $n > 1$.

In order to construct a higher dimensional ϕ^4 kink, we must add curvature. In the next section we introduce a curved background, and show that a modified ϕ^4 kink exists on this background. We will also examine a limit in which the modified kink reduces to the flat kink. In Section 2.3 we consider linearised perturbations around the modified kink, proving that it is linearly stable and comparing its discrete spectrum to that of the flat kink. In Section 2.4 we examine the mode of decay to the modified kink in the full non-linear theory, in particular the resonant coupling of its internal modes to the continuous spectrum.

2.2 The static kink on a wormhole

We now replace the flat $\mathbb{R}^{1,1}$ background with a wormhole spacetime (M, g) , where

$$g = -dt^2 + dr^2 + (r^2 + a^2)(d\vartheta^2 + \sin^2 \vartheta d\varphi^2)$$

for some constant $a > 0$, and $-\infty < r < \infty$. This spacetime was first studied by Ellis [32] and Bronnikov [9], and has featured in a number of recent studies about kinks and their stability [5, 6]. Note the presence of asymptotically flat ends as $r \rightarrow \pm\infty$, connected by a spherical throat of radius a at $r = 0$.

Our action is modified by the presence of a non-flat metric:

$$S = \int \left(\frac{1}{2} g^{ab} \partial_a \phi \partial_b \phi + \frac{1}{2} (1 - \phi^2)^2 \right) \sqrt{-g} dx,$$

where x^a are now local coordinates on M . Variation with respect to ϕ gives

$$\square_g \phi + 2\phi(1 - \phi^2) = 0 \tag{2.5}$$

where $\square_g \phi = \frac{1}{\sqrt{-g}} \partial_a (g^{ab} \sqrt{-g} \partial_b \phi)$. We will always assume ϕ is independent of the angular coordinates (ϑ, φ) , so (2.5) can be written explicitly as

$$\phi_{tt} = \phi_{rr} + \frac{2r}{r^2 + a^2} \phi_r + 2\phi(1 - \phi^2). \tag{2.6}$$

The conserved energy in the theory is given by

$$E = \int_{-\infty}^{+\infty} \left(\frac{1}{2} (\phi_t)^2 + \frac{1}{2} (\phi_r)^2 + \frac{1}{2} (1 - \phi^2)^2 \right) (r^2 + a^2) dr,$$

which we require to be finite. This imposes the condition $\phi^2 \rightarrow 1$ as $r \rightarrow \pm\infty$, so that the field lies at one of the two vacua at both asymptotically flat ends.

Static solutions $\phi(r)$ satisfy

$$\phi'' + \frac{2r}{r^2 + a^2} \phi' = -\frac{d}{d\phi} \left(-\frac{1}{2}(1 - \phi^2)^2 \right), \quad (2.7)$$

which, imagining r as a time coordinate, can be thought of as a Newtonian equation of motion for a particle at position ϕ moving in a potential $\mathcal{U}(\phi) = -(1 - \phi^2)/2$, with a time dependent friction term.

In addition to the two vacuum solutions, we have a single soliton solution which interpolates between the saddle points at $(-1, 0)$ and $(1, 0)$ in the (ϕ, ϕ') plane. Its existence and uniqueness among odd parity solutions follow from a shooting argument: suppose the particle lies at $\phi = 0$ when $r = 0$. If its velocity $\phi'(0)$ is too small, it will never reach the local maximum of the potential at $\phi = 1$, but if $\phi'(0)$ is too large it will overshoot the maximum so that $\mathcal{U}(\phi) \rightarrow -\infty$ as $r \rightarrow \infty$, thus having infinite energy. Continuity ensures that there is some critical velocity $\phi'(0)$ such that the particle reaches $\phi = 1$ in infinite time and has zero velocity upon arrival. This corresponds to the non-trivial kink solution, which we call $\Phi(r)$. Time reversal implies that $\phi \rightarrow -1$ as $r \rightarrow -\infty$, and that the anti-kink $\phi(r) = -\Phi(r)$ is also a solution.

We can find $\Phi(r)$ numerically using a shooting method for the gradient at $r = 0$. Figure 2.1 shows such numerically generated kinks for several values of a . Note that the absolute value of $\Phi(r)$ is always greater than or equal to that of the flat kink $\Phi_0(r)$, and that at fixed non-zero r , the absolute value of Φ decreases as a increases. The reason for this will become clear in Section 2.3. In Section 2.2.1 we examine $\Phi(r)$ in the limit where a is large, finding that it reduces to the flat kink $\Phi_0(r)$, and examining its departure from the flat kink at first order in $1/a^2$.

We again label the values at the boundary as

$$\phi_{\pm} := \lim_{r \rightarrow \pm\infty} \Phi(r) \in \{\pm 1\}.$$

Since no finite energy deformation can change the value of the topological charge $N = (\phi_+ - \phi_-)/2 \in \{-1, 0, 1\}$, we again conclude that $\Phi(r)$ is topologically stable.

2.2.1 Large a limit

As $a \rightarrow \infty$, equation (2.6) becomes the standard equation (2.2) for the flat kink. It is thus helpful to expand the modified kink in $\epsilon^2 := 1/a^2$ for small ϵ^2 , since we can then




Fig. 2.1 The kink solution for several values of a , along with the flat kink $\Phi_0(r)$.

solve both (2.6) and (2.10) analytically up to $\mathcal{O}(\epsilon^4)$. We shall denote the static kink by $\Phi_\epsilon(r)$ in this limit. It satisfies

$$\Phi_\epsilon'' + \frac{2r\epsilon^2}{\epsilon^2 r^2 + 1} \Phi_\epsilon' = -2\Phi_\epsilon(1 - \Phi_\epsilon^2). \quad (2.8)$$

Setting $\Phi_\epsilon(r) = \Phi_0(r) + \epsilon^2 \Phi_1(r) + \mathcal{O}(\epsilon^4)$ we obtain at order zero the equation (2.2) of a static kink on $\mathbb{R}^{1,1}$. This has solution $\Phi_0(r)$, where we have chosen the kink at the origin to restrict to solutions with odd parity.

At order ϵ^2 we find that $\Phi_1(r)$ must satisfy

$$\Phi_1'' + 2r \operatorname{sech}^2 r = 2\Phi_1(2 - 3\operatorname{sech}^2 r).$$

The unique solution which is odd and decays as $r \rightarrow \pm\infty$ is given by

$$\Phi_1(r) = \frac{1}{24} \operatorname{sech}^2 r (f_1(r) + f_2(r) + f_3(r)),$$

where

$$\begin{aligned} f_1(r) &= r[3 - 8\cosh(2r) - \cosh(4r)], \\ f_2(r) &= \sinh(2r)[8\log(2\cosh(r)) - 1] + \sinh(4r)\log(2\cosh(r)), \\ f_3(r) &= \frac{\pi^2}{2} + 6r^2 + 6\text{Li}_2(-e^{-2r}), \end{aligned}$$

and $\text{Li}_2(z)$ is the dilogarithm function, which is the special case $s = 2$ of the polylogarithm

$$\text{Li}_s(z) = \sum_{k=1}^{\infty} \frac{z^k}{k^s}.$$

To show that $\Phi_1(r)$ is odd, note that $\text{sech}^2 r$ is an even function, and that f_1 and f_2 are constructed from products of even and odd functions, and hence are odd. To see that f_3 is also odd, we use Landen identity for the dilogarithm:

$$\begin{aligned} \text{Li}_2(-e^{-2r}) + \text{Li}_2(-e^{2r}) &= -\frac{\pi^2}{6} - \frac{1}{2}[\log(e^{-2r})]^2 \\ &= -\frac{\pi^2}{6} - 2r^2, \end{aligned}$$

thus verifying $f_3(r) + f_3(-r) = 0$.

We now turn to the behaviour of $\Phi_1(r)$ as $r \rightarrow \infty$. Since $\text{sech}^2 r \sim 4e^{-2r}$ for large r , we need only consider terms in the $\{f_i\}$ of order e^{2r} or higher. We first note that

$$\begin{aligned} \log(2\cosh r) &= \log(e^r(1 + e^{-2r})) = r + \log(1 + e^{-2r}) \\ &= r + e^{-2r} + \mathcal{O}(e^{-4r}). \end{aligned}$$

Then

$$\begin{aligned} f_1(r) &= -4re^{2r} - \frac{r}{2}e^{4r} + \mathcal{O}(e^r) \\ f_2(r) &= \frac{1}{2}e^{2r}(8r + 8e^{-2r} - 1) + \frac{1}{2}e^{4r}(r + e^{-2r}) + \mathcal{O}(e^r) \\ &= 4re^{2r} + \frac{r}{2}e^{4r} + \mathcal{O}(e^r), \end{aligned}$$

so $f_1(r) + f_2(r) = \mathcal{O}(e^r)$. Since $f_3(r) = \mathcal{O}(r^2)$ for large r , we see that $\Phi_1(r)$ vanishes as $r \rightarrow \infty$, as we expect. Note that its vanishing as $r \rightarrow -\infty$ then follows using parity. A plot of $\Phi_1(r)$ is shown in Figure 2.2.




Fig. 2.2 The order ϵ^2 perturbation to the static kink on $\mathbb{R}^{1,1}$.

2.3 Linearised perturbations around the kink

To study the linear stability of the kink, we first plug

$$\phi(t, r) = \Phi(r) + w(t, r) \quad (2.9)$$

into equation (2.6), discarding terms non-linear in w . Imposing the fact that $\Phi(r)$ satisfies (2.7), we find

$$w_{tt} = w_{rr} + \frac{2r}{r^2 + a^2} w_r + 2w(1 - 3\Phi^2).$$

For $w(t, r) = e^{i\omega t} (r^2 + a^2)^{-1/2} v(r)$, this becomes a one-dimensional Schrödinger equation

$$Lv := (-\partial_r \partial_r + V(r))v = \omega^2 v, \quad (2.10)$$

where the potential is given by

$$V(r) = \frac{a^2}{(r^2 + a^2)^2} - 2(1 - 3\Phi^2). \quad (2.11)$$




Fig. 2.3 The potential of the 1-dimensional quantum mechanics problem arising from the study of stability of the soliton for values of a between $a = 10$ and $a = 0.3$.

Figure 2.3 shows the potential $V(r)$ for several values of a . Note that for large a it has a single well with a minimum at $r = 0$, close to the potential V_0 corresponding to the flat kink. As a decreases, the critical point at $r = 0$ becomes a maximum with minima on either side, creating a double well. We find numerically that this happens at about $a = 0.55$. Note that those potentials with $a < 1/\sqrt{2}$ are everywhere positive.

Proposition 2.3.1. *The kink solution $\Phi(r)$ is linearly stable with respect to spherically symmetric perturbations.*

Proof. We first decompose the potential $V(r)$ in (2.10) as $V = V_0 + V_1 + V_a$, where

$$V_0 = -2[1 - 3\Phi_0(r)^2], \quad V_1 = 6[\Phi(r)^2 - \Phi_0(r)^2], \quad V_a = \frac{a^2}{(r^2 + a^2)^2}.$$

As discussed above, we know that the operator $L_0 = -\partial_r \partial_r + V_0$ has no negative eigenvalues. It then follows that L itself has no negative eigenvalues as long as the functions $V_1(r)$ and $V_a(r)$ are everywhere non-negative.

The latter is obvious; to prove the former we recall that we can think of $\Phi(r)$ and $\Phi_0(r)$ as the trajectories of particles moving in a potential $\mathcal{U}(\phi)$, where r is imagined as the time coordinate. The particle corresponding to $\Phi(r)$ suffers an increased frictional

force compared to $\Phi_0(r)$, i.e.

$$\Phi_0'' = -\frac{\partial \mathcal{U}}{\partial \phi} \Big|_{\phi=\Phi_0}, \quad \Phi'' + \frac{2r}{r^2 + a^2} \Phi' = -\frac{\partial \mathcal{U}}{\partial \phi} \Big|_{\phi=\Phi}. \quad (2.12)$$

Both Φ and Φ_0 interpolate between the maxima of \mathcal{U} at $\phi = \pm 1$; reaching the minimum ($\phi = 0$) when $r = 0$.

Multiplying the equations (2.12) by Φ_0' and Φ' respectively, then integrating from r to ∞ , we have that at every instant of time

$$\frac{1}{2}(\Phi_0')^2 + \mathcal{U}(\Phi_0) = 0, \quad \frac{1}{2}(\Phi')^2 + \mathcal{U}(\Phi) = \int_r^\infty \frac{2r}{r^2 + a^2} (\Phi')^2 dr. \quad (2.13)$$

These equations are equivalent to conservation of energy for each of the particles. Note that the integral on the right hand side is non-negative for $r \geq 0$, and vanishes only at $r = \infty$. In particular, when $r = 0$ we have $\mathcal{U}(\Phi) = \mathcal{U}(\Phi_0) = -1/2$, so $\Phi'(0) > \Phi_0'(0)$. This means $V_1(r)$ is initially increasing from zero.

For $V_1(r)$ to return to zero at some finite $r = r_0$, we would need that $\Phi(r_0) = \Phi_0(r_0)$ at a point where $\Phi'(r_0) \leq \Phi_0'(r_0)$. However, this is made impossible by equations (2.13), since at such a point $\mathcal{U}(\Phi) = \mathcal{U}(\Phi_0)$ and the integral on the right hand side is positive. Hence $V_1(r)$ remains non-negative for all $r > 0$, and thus for all r since it is even in r .

□

2.3.1 Finding internal modes numerically

Bound states of the potential (2.11) correspond to internal modes of the kink like the odd solution of (2.3) in (2.4). In contrast, for frequencies greater than $\omega = 2$, solutions to (2.10) are interpreted as radiation. It is possible to search for bound states of (2.11) numerically. The method for this is as follows:

1. For the chosen value of a , generate the soliton $\Phi(r)$ as described above.
2. Calculate the potential $V(r)$.
3. For some initial guess of the eigenvalue ω^2 , integrate equation (2.10) numerically, setting $v(0) = 1$ and $v'(0) = 0$ to obtain even bound states and $v(0) = 0, v'(0) = 1$ to obtain odd bound states.
4. Use a bisection method to find the value of ω^2 for which a bound state exists.




Fig. 2.4 Even bound states of the potential $V_0(r)$ and of the potential $V(r)$ for two different values of a .

This procedure will only be effective within the range of r for which $\Phi(r)$ is calculated.

For large a , the potential has both an even and an odd bound state which look qualitatively similar to the internal modes (2.4) of the ϕ^4 kink on $\mathbb{R}^{1,1}$. The bound states for several values of a can be found in Figures 2.4 and 2.5. As a decreases, the eigenvalues ω^2 of the bound states increase, until they disappear into the continuous spectrum ($\omega^2 > 4$). This disappearance will be further discussed in Section 2.3.3. The internal mode frequencies are plotted against a in Figure 2.6. The choice of axis ticks will be motivated in Section 2.4.

2.3.2 Large a limit

We can also perturbatively expand the eigenvalues of the eigenvalue problem (2.10). Consider solutions to (2.6) of the form¹ $\phi_\epsilon(r) = \Phi_\epsilon(r) + e^{i\omega t}v_\epsilon(r)$, where v_ϵ is small. These satisfy

$$v_\epsilon'' + \frac{2r\epsilon^2}{\epsilon^2 r^2 + 1} v_\epsilon' + 2(1 - 3\Phi_\epsilon^2)v_\epsilon = -\omega_\epsilon^2 v_\epsilon. \quad (2.14)$$

¹Note that in Section 2.3 we considered perturbations $v(r)$ which differ from $v_\epsilon(r)$ by a factor of $(r^2 + a^2)^{-1/2}$, since such perturbations are described by a Schrödinger problem. Here it will be simpler to remove this factor; however there is a one-to-one correspondence between $v(r)$ and $v_\epsilon(r)$.




Fig. 2.5 Odd bound states of the potential $V_0(r)$ and of the potential $V(r)$ for two different values of a .




Fig. 2.6 The frequencies of the internal modes of the kink plotted against the wormhole radius a .

Let $(v_\epsilon, \omega_\epsilon^2)$ be a solution to (2.14) with

$$\omega_\epsilon^2 = \omega_0^2 + \epsilon^2 \xi + \mathcal{O}(\epsilon^4) \quad \text{and} \quad v_\epsilon(r) = v_0(r) + \epsilon^2 v_1(r) + \mathcal{O}(\epsilon^4).$$

Our aim will be to find ξ . Substituting into (2.14), at zero order we obtain the equation (2.3) which controls the linear stability analysis of the ϕ^4 kink on $\mathbb{R}^{1,1}$.

The terms of order ϵ^2 in (2.14) give us

$$v_1'' + 2rv_0' + 2(1 - 3\Phi_0^2)v_1 - 12\Phi_0\Phi_1 v_0 = -\omega_0^2 v_1 - \xi v_0. \quad (2.15)$$

We multiply equation (2.15) by v_0 , and subtract from this v_1 multiplied by equation (2.3). Integrating the result from $r = -\infty$ to $r = \infty$, we find

$$\int_{-\infty}^{\infty} (v_1'' v_0 - v_0'' v_1) dr + \int_{-\infty}^{\infty} 2rv_0' v_0 dr - 12 \int_{-\infty}^{\infty} \Phi_0 \Phi_1 v_0^2 dr = -\xi.$$

In the first term the integrand is a total derivative, and the second term is easily found to be -1 using integration by parts. We thus obtain

$$\xi = 1 + 12 \int_{-\infty}^{\infty} \Phi_0 \Phi_1 v_0^2 dr, \quad (2.16)$$

which we can evaluate for each of the solutions (2.4) using the symbolic computation facility in Mathematica. We find $\xi = 2$ in the case of the zero mode and $\xi = \pi^2 - 7$ in the case of the first non-trivial vibrational mode. We can check these values by finding (v, ω) numerically as described in Section 2.3.1 for a range of small values of ϵ and comparing ω^2 to the $\omega_0^2 + \xi\epsilon^2$ predicted here. The corresponding plots are shown in Figures 2.7 and 2.8.

2.3.3 Critical values of a

It is interesting to investigate the values of a at which the internal modes disappear into the continuous spectrum. The larger of these, at which the odd internal mode disappears, we shall call a_1 . The smaller one, at which the even internal mode disappears, we shall call a_0 .

The most convenient method of estimating a_0 and a_1 is based on the Sturm Oscillation Theorem 2.1.1. The points at which the even and odd internal modes disappear into the continuous spectrum are the points at which the zeros of the even and odd eigenfunctions of L with $\omega^2 = 4$ disappear. We can thus examine the number of zeros of the odd eigenfunction with $\omega^2 = 4$ to determine the number of odd bound




Fig. 2.7 A comparison of the predicted and numerical calculations for the energy of the zero mode as a function of ϵ^2 for small ϵ .




Fig. 2.8 A comparison of the predicted and numerical calculations for the energy of the odd vibrational mode as a function of ϵ^2 for small ϵ .

states with $\omega^2 < 4$. The critical value a_1 which we are searching for can then be found using a bisection method. An equivalent method using even bound states will yield an estimate of a_0 .

One problem with this method is that we need the number of zeros in the interval $(0, \infty)$, and the shooting method we use to generate $\Phi(r)$ and $V(r)$ is only accurate up to a finite value of r . Since zeros of the eigenfunction with $\omega^2 = 4$ disappear at $r = \infty$, this limits the accuracy with which we can determine a_0 and a_1 .

For the finite integration range which is accessible based on the shooting method, the odd state disappears at $a_1 \approx 0.8$ and the even state disappears at $a_0 \approx 0.3$.

For a potential $\mathcal{V}(r)$ which decays sufficiently quickly as $r \rightarrow \pm\infty$, it is well known that the condition

$$I := \int_{-\infty}^{\infty} \mathcal{V}(r) dr < 0 \quad (2.17)$$

is sufficient to ensure that the operator $-\partial_r \partial_r + \mathcal{V}(r)$ has at least one bound state. In fact, the condition $I \leq 0$ is sufficient [68]. However, (2.17) is not a necessary condition: there are potentials which have at least one bound state where (2.17) is not satisfied. It is interesting to investigate the disappearance of our ground state in this context.

For us the relevant choice is $\mathcal{V}(r) = V(r) - 4$ so that $\mathcal{V}(r)$ vanishes at the boundaries. We then examine the value of this integral for the critical value $a = a_0$ when the ground state disappears. We find that $I \approx 0$ at the critical value of $a_0 \approx 0.3$ given above. We can also search numerically for the value of a at which $I = 0$; this also occurs at around $a_0 \approx 0.3$. Thus our results would be consistent with the conjecture that (2.10) has no bound states for $I > 0$.

2.4 Resonant coupling of the internal modes to the continuous spectrum

We now move on to consider time dependent perturbations of the form

$$\phi(t, r) = \Phi(r) + (r^2 + a^2)^{-1/2} w(t, r),$$

where we consider non-linear terms in $w(t, r)$. Substituting into (2.6) we find

$$w_{tt} = -Lw + f(w), \quad (2.18)$$

where we have defined

$$f(w) = -\frac{6w^2\Phi}{\sqrt{a^2 + r^2}} - \frac{2w^3}{a^2 + r^2}, \quad (2.19)$$

suppressing the dependence of f on r to simplify the notation. We will not have much need for the expression for f other than to note that it contains terms which are quadratic and cubic in w .

If a is large enough to allow internal modes, then these can only decay through resonant coupling to the continuous spectrum of L . The analogous process of decay to the ϕ^4 kink on $\mathbb{R}^{1,1}$ was discussed in [52], and the general theory was developed in [69]. In the following sections we investigate this decay in the case of a single internal mode, before comparing our result with numerical data.

2.4.1 Conjectured decay rate in the presence of a single internal mode

In this section we follow the analysis in [6]. Looking at Figure 2.6, we note that for $a \in (0.3, 0.8)$ we have

$$\text{spec } L = \{\omega^2\} \cap [m^2, \infty), \quad \omega^2 < m^2 < 4\omega^2 \quad (2.20)$$

where $m^2 = 4$. As above, we denote the unique normalised eigenfunction of L by v , so that $Lv = \omega^2 v$. We will use $\langle \cdot, \cdot \rangle$ to denote the usual inner product on \mathbb{R} .

We decompose the perturbation as

$$w(t, r) = \alpha(t)v(r) + \eta(t, r), \quad (2.21)$$

where $v(r)$ refers to the single even internal mode of the kink and η is a superposition of states from the continuous spectrum of L . Where there is only one internal mode present, its frequency ω always lies in the upper half of the mass gap: $1 < \omega < 2$. This is important because it means that 2ω lies within the continuous spectrum.

We substitute this into (2.18) and project onto and away from the internal mode direction, obtaining the following equations for α and η :

$$\ddot{\alpha} + \omega^2\alpha = \langle v, f(\alpha v + \eta) \rangle \quad (2.22)$$

$$\ddot{\eta} + L\eta = P^\perp f(\alpha v + \eta), \quad (2.23)$$

where P^\perp is the projection onto the space of eigenstates of L which are orthogonal to v , given by

$$P^\perp \psi = \psi - \langle v, \psi \rangle v. \quad (2.24)$$

These equations have initial conditions $\alpha(0)$ and $\eta(0, r)$ such that

$$\begin{aligned} \phi(0, r) &= \Phi(r) + (r^2 + a^2)^{-1/2}(\alpha(0)v(r) + \eta(0, r)), \quad \text{and} \\ \dot{\phi}(0, r) &= (r^2 + a^2)^{-1/2}(\dot{\alpha}(0)v(r) + \dot{\eta}(0, r)). \end{aligned}$$

In the following analysis we investigate decay of $\alpha(t)$. Equation (2.22) has a homogeneous solution consisting of oscillations with frequency ω . Since 2ω lies within the continuous spectrum of L , there will be a resonant interaction between these oscillations and the radiation modes in η with frequencies $\pm 2\omega$, arising from the term of order α^2 in the right hand side of (2.23). Thus, to leading order, (2.23) is a driven wave equation with driving frequency 2ω . This resonant part of η will have a back-reaction on α through (2.22), which will result in decay of the internal mode oscillations.

We now define $\alpha_1 = \alpha$, $\omega\alpha_2 = \dot{\alpha}_1$ so that (2.22) becomes

$$\begin{cases} \dot{\alpha}_1 = \omega\alpha_2, \\ \dot{\alpha}_2 = -\omega\alpha_1 + \frac{1}{\omega}\langle v, f(\alpha_1 v + \eta) \rangle, \end{cases}$$

or equivalently

$$\dot{A} = -i\omega A + \frac{i}{\omega} \left\langle v, f\left(\frac{1}{2}(A + \bar{A})v + \eta\right) \right\rangle, \quad (2.25)$$

where $A = \alpha_1 + i\alpha_2$. Next we write $\eta_1 = \eta$, $\eta_2 = \dot{\eta}_1$, converting (2.23) to

$$\begin{cases} \dot{\eta}_1 = \eta_2, \\ \dot{\eta}_2 = -L\eta_1 + P^\perp f\left(\frac{1}{2}(A + \bar{A})v + \eta_1\right). \end{cases}$$

We will regard the right hand sides of (2.25) and (2.4.1) as power series in A and η . Terms which we expect to be higher order will not be treated rigorously; for this reason, our analysis will produce only a conjecture about the decay rate. Numerical evidence concerning the conjecture will be discussed in Section 2.4.2.

It will be helpful to introduce the notation $\mathcal{O}_p(A, \eta)$ to mean terms of at least order p in $A, \bar{A}, \eta_1, \eta_2$, so that A^2 , η_1^2 and $\bar{A}\eta_1$ are all examples of terms which are $\mathcal{O}_2(A, \eta)$.

Currently, the coupling between (2.25) and (2.4.1) is $\mathcal{O}_2(A, \eta)$. We will write

$$f\left(\frac{1}{2}(A + \bar{A})v + \eta_1\right) = \sum_{k+l \geq 2} f_{kl} A^k \bar{A}^l + \sum_{\substack{k+l \geq 1 \\ n \geq 1}} f_{kln} \eta_1 A^k \bar{A}^l$$

where k, l, n are non-negative, to elucidate the lowest order terms in (2.4.1). Note that f_{kl} and f_{kln} are decaying functions of r defined by (2.19). We can then write

$$P^\perp \left[f\left(\frac{1}{2}(A + \bar{A})v + \eta_1\right) \right] = \sum_{k+l=2} P^\perp[f_{kl}] A^k \bar{A}^l + \sum_{k+l=1} P^\perp[f_{kln} \eta_1] A^k \bar{A}^l + \mathcal{O}_3(A, \eta).$$

Terms in (2.25) with imaginary coefficients correspond to rotation in the complex plane, and thus to oscillatory behaviour in α . At first order, A oscillates with frequency ω . This is exactly the behaviour expected in the linearised theory discussed in Section 2.3. In fact, a priori, all the terms in the power series for \dot{A} have coefficients which are purely imaginary.

The next step in our analysis will be to attempt a change of variable $\eta_i \mapsto \tilde{\eta}_i$ in (2.4.1) so that its right hand side is $\mathcal{O}_3(A, \tilde{\eta})$, meaning $\tilde{\eta}$ is $\mathcal{O}(A^3)$. It will turn out that the required change of variables is complex. The result will be a term in (2.25) which is $\mathcal{O}(A^3)$ and has a real coefficient. This will be the lowest order term with a real coefficient, and thus the key resonant damping term.

We write the change of variables as

$$\eta_1 = \tilde{\eta}_1 + \sum_{k+l=2} b_{kl} A^k \bar{A}^l, \quad \eta_2 = \tilde{\eta}_2 + \sum_{k+l=2} c_{kl} A^k \bar{A}^l, \quad (2.26)$$

where b_{kl} and c_{kl} are functions of r which are so far undetermined. Differentiating with respect to time and using (2.25), we find

$$\begin{aligned} \dot{\eta}_1 &= \dot{\tilde{\eta}}_1 - i\omega \sum_{k+l=2} b_{kl} (k-l) A^k \bar{A}^l + \mathcal{O}_3(A, \tilde{\eta}), \\ \dot{\eta}_2 &= \dot{\tilde{\eta}}_2 - i\omega \sum_{k+l=2} c_{kl} (k-l) A^k \bar{A}^l + \mathcal{O}_3(A, \tilde{\eta}). \end{aligned}$$

We equate these to the right hand sides of (2.4.1), substituting from (2.26) and requiring that

$$\dot{\tilde{\eta}}_1 = \tilde{\eta}_2 + \mathcal{O}_3(A, \tilde{\eta}), \quad \dot{\tilde{\eta}}_2 = -L\tilde{\eta}_1 + \mathcal{O}_3(A, \tilde{\eta}). \quad (2.27)$$

This yields

$$-i\omega b_{kl}(k-l) = c_{kl} \quad \text{and} \quad -i\omega c_{kl}(k-l) = -Lb_{kl} + P^\perp[f_{kl}]$$

for $k+l=2$, where we have discarded

$$\begin{aligned} \sum_{k+l=1} P^\perp[f_{kl1}\eta_1]A^k\bar{A}^l &= \sum_{k+l=1} P^\perp[f_{kl1}\tilde{\eta}_1]A^k\bar{A}^l + \sum_{\substack{k+l=1 \\ p+q=2}} P^\perp[f_{kl1}b_{pq}]A^{k+p}\bar{A}^{l+q} \\ &= \mathcal{O}_3(A) \end{aligned}$$

because $\tilde{\eta}$ is at least third order in A .

The change of variables (2.26) is now given by the solution to

$$(L - \omega^2(k-l)^2)b_{kl} = P^\perp[f_{kl}]. \quad (2.28)$$

Because of the spectrum of L given in (2.20), for $(k,l) \in (2,0) \cup (0,2)$ the solution b_{kl} is in general a complex function of r , whilst for $k=l=1$ the solution is real and decaying. The reason for this can be understood using the variation of parameters method for inhomogeneous ordinary differential equations.

Let $g(r)$ be such that $\langle g, g \rangle$ is finite, and $\lambda \geq 0$ a constant. The general solution of

$$(L - \lambda^2)b(r) = g(r)$$

is given by

$$b(r) = Z_2(r) \int_{-\infty}^r \frac{1}{W(r')} Z_1(r')g(r')dr' + Z_1(r) \int_r^\infty \frac{1}{W(r')} Z_2(r')g(r')dr',$$

where $\{Z_1, Z_2\}$ is a basis for solutions to the homogeneous equation with Wronskian $W(r) = Z_1Z_2' - Z_2Z_1'$. The basis must be chosen so that the above integrals converge.

For $k=l=1$, so that $\lambda^2 = 0$ and hence $\lambda^2 < m^2$, we can choose a basis such that $W=1$ and Z_1, Z_2 are both real, and they decay to zero in the limits $r \rightarrow -\infty$ and $r \rightarrow \infty$ respectively. Then

$$b_{11}(r) = Z_2(r) \int_{-\infty}^r Z_1(r')P^\perp[f_{11}](r')dr' + Z_1(r) \int_r^\infty Z_2(r')P^\perp[f_{11}](r')dr'. \quad (2.29)$$

For $\lambda^2 \geq m^2$, we cannot choose a real solution in general. In the case $(k, l) \in (2, 0) \cup (0, 2)$, we take as a basis the Jost functions $\{j_{\pm}\}$, defined by

$$j_{\pm}(r) \sim e^{\pm i\xi r} \text{ as } r \rightarrow \infty,$$

where $\xi = \sqrt{4\omega^2 - m^2}$. Their Wronskian is then $W(j_+, j_-) = -2i\xi$, and we write the solution as

$$b_{02}(r) = b_{20}(r) = \frac{ij_-(r)}{2\xi} \int_{-\infty}^r j_+(r') P^{\perp}[f_{20}](r') dr' + \frac{ij_+(r)}{2\xi} \int_r^{\infty} j_-(r') P^{\perp}[f_{20}](r') dr'. \quad (2.30)$$

Finally, we use (2.29) and (2.30) to change variable $\eta_i \mapsto \tilde{\eta}_i$ in (2.25), obtaining

$$\dot{A} = -i\omega A + \frac{i}{\omega} \left(\sum_{2 \leq k+l \leq 3} \langle v, f_{kl} \rangle A^k \bar{A}^l + \sum_{\substack{k+l=1 \\ p+q=2}} \langle v, f_{kl1} b_{pq} \rangle A^{k+p} \bar{A}^{l+q} + \mathcal{O}(A^4) \right), \quad (2.31)$$

where we have ignored terms containing $\tilde{\eta}_1$ since these are at least fourth order in A . We can now see that, of the terms which we have written explicitly, the only ones that can give a real contribution to \dot{A} are those containing b_{02} and b_{20} . We thus find

$$\begin{aligned} \frac{d}{dt} |A|^2 &= \dot{A} \bar{A} + A \dot{\bar{A}} = 2\mathbb{R}\text{e}[\dot{A} \bar{A}] \\ &= \frac{2}{\omega} \mathbb{R}\text{e} \left[i \sum_{k+l=1} \left(\langle v, f_{kl1} b_{20} \rangle A^{k+2} \bar{A}^{l+1} + \langle v, f_{kl1} b_{02} \rangle A^k \bar{A}^{l+3} \right) \right] + \mathcal{O}(A^5) \\ &= -\frac{2}{\omega} \mathbb{I}\text{m} \left[\langle v, f_{101} b_{20} \rangle (A^3 \bar{A} + A^2 \bar{A}^2 + A \bar{A}^3 + \bar{A}^4) \right] + \mathcal{O}(A^5). \end{aligned}$$

In particular, the term $A^2 \bar{A}^2 = |A|^4$ is real and non-oscillating, giving a contribution

$$\frac{d}{dt} |A|^2 \sim -\frac{2}{\omega} \mathbb{I}\text{m}[\langle v, f_{101} b_{20} \rangle] |A|^4.$$

The terms $A^3 \bar{A}$, $A \bar{A}^3$ and \bar{A}^4 , on the other hand, would be expected to oscillate at frequencies 2ω and 4ω at first order, and thus time average to zero.

Hence we conclude

$$|A| \sim \left(\Gamma t + \frac{1}{|A(0)|^2} \right)^{-1/2}, \quad \Gamma := \frac{2}{\omega} \mathbb{I}\text{m}[\langle v, f_{101} b_{20} \rangle]. \quad (2.32)$$

The constant Γ is a function of a which can be calculated explicitly. Using (2.19) and (2.30) gives

$$\begin{aligned} \langle v, f_{101}b_{20} \rangle &= \int_{-\infty}^{\infty} dr \, v(r) f_{101}(r) \left(\frac{ij_-(r)}{2\xi} \int_{-\infty}^r j_+(r') P^\perp[f_{20}](r') dr' \right. \\ &\quad \left. + \frac{ij_+(r)}{2\xi} \int_r^{\infty} j_-(r') P^\perp[f_{20}](r') dr' \right). \end{aligned}$$

We now use the facts that $f_{20} = vf_{101}/4$, and $P^\perp[f_{20}] = f_{20} - \langle v, f_{20} \rangle v$. Note that f_{20} is an odd function of r , so in fact $\langle v, f_{20} \rangle = 0$ and so $P^\perp[f_{20}] = f_{20}$. We thus obtain

$$\begin{aligned} \langle v, f_{101}b_{20} \rangle &= \frac{2i}{\xi} \left(\int_{-\infty}^{\infty} dr \int_{-\infty}^r dr' f_{20}(r) j_-(r) f_{20}(r') j_+(r') \right. \\ &\quad \left. + \int_{-\infty}^{\infty} dr \int_r^{\infty} dr' f_{20}(r) j_+(r) f_{20}(r') j_-(r') \right). \end{aligned}$$

The two double integrals are integrals over complementary halves of the (r, r') plane, and thus sum to a single integral over the full plane. Hence

$$\langle v, f_{101}b_{20} \rangle = \frac{2i}{\xi} \int_{-\infty}^{\infty} f_{20}(r) j_+(r) dr \int_{-\infty}^{\infty} f_{20}(r') j_-(r') dr' = \frac{2i}{\xi} |\langle f_{20}, j_+ \rangle|^2,$$

since j_\pm are complex conjugates.

Combining this with (2.32) gives

$$\Gamma = \frac{4}{\omega\xi} |\langle f_{20}, j_+ \rangle|^2.$$

The so-called Fermi Golden Rule then reads

$$|\langle f_{20}, j_+ \rangle| \neq 0.$$

2.4.2 Numerical investigation of the conjectured decay rate

In order to integrate the PDE (2.6) to large times t , we employ the method of hyperboloidal foliations and scri-fixing [86]. This is necessary because surfaces of constant t can never intersect with future null infinity, as they always reach the boundary of the spacetime at spatial infinity. In order to integrate out to large times




Fig. 2.9 The decay of internal mode oscillations for various initial conditions when $a = 0.5$.

at large r , i.e. to reach future null infinity, we define

$$s = \frac{t}{a} - \sqrt{\frac{r^2}{a^2} + 1}, \quad y = \arctan\left(\frac{r}{a}\right),$$

resulting in the hyperbolic equation

$$\partial_s \partial_s F + 2\sin(y) \partial_y \partial_s F + \frac{1 + \sin^2(y)}{\cos(y)} \partial_s F = \cos^2(y) \partial_y \partial_y F + 2a^2 \frac{F(1 - F^2)}{\cos^2(y)} \quad (2.33)$$

for $F(s, y) = \phi(t, r)$. Then surfaces of constant s approach right future null infinity \mathcal{J}_R^+ along outgoing null cones of constant retarded time $t - r$, and left future null infinity \mathcal{J}_L^+ along outgoing null cones of constant advanced time $t + r$. For more details see [6, 86].

We solve the corresponding initial value problem at space-like hypersurfaces of constant s , specifying $\phi(s = 0, y)$ and $\partial_s \phi(s = 0, y)$. No boundary conditions are required, since the principal symbol of (2.33) degenerates to $\partial_s(\partial_s \pm 2\partial_y)$ as $y \rightarrow \pm\pi/2$, so there are no ingoing characteristics. This reflects the fact that no information comes in from future null infinity.

Following [5, 87] we define the auxiliary variables

$$\Psi = \partial_y F, \quad \Pi = \partial_s F + \sin y \partial_y F$$

to obtain the first order symmetric hyperbolic system

$$\partial_s F = \Pi - \Psi \sin y \quad (2.34)$$

$$\partial_s \Psi = \partial_y (\Pi - \Psi \sin y) \quad (2.35)$$

$$\partial_s \Pi = \partial_y (\Psi - \Pi \sin y) + 2 \tan y (\Psi - \Pi \sin y) + 2a^2 \frac{F(1 - F^2)}{\cos^2 y}, \quad (2.36)$$

which we solve numerically using the method of lines. Kreiss–Oliger dissipation is required to reduce unphysical high-frequency noise. We also add the term $-0.1(\Psi - \partial_y F)$ to the right hand side of equation (2.35) to suppress violation of the constraint $\Psi = \partial_y F$.

We are interested in the range of values $a_0 < a < a_1$ for which the kink has exactly one internal mode. We find that, for fixed but arbitrary y , $F(s, y)$ oscillates in s with a frequency close to the internal mode frequency, and that these oscillations tend towards a decay rate of $s^{-1/2}$, as we expect from Section 2.4.1. Plots demonstrating this decay at $y = 0$ for $a = 0.5$ are shown in Figure 2.9. Note that $\phi(0, s)$ is used as a proxy for the internal mode amplitude, and we use a log–log scale to elucidate the dependence on $s^{-1/2}$ in the large s limit. The lines are labelled in the legend by the initial conditions which produced them, with the exception of the gradient line $4.2s^{-1/2}$. The constant 4.2 is related to Γ as defined in (2.32).

2.4.3 Expected decay rates in other regimes

From Figure 2.6, we see that the second internal mode appears before the frequency of the first internal mode moves out of the range $(m/2, m)$. In the presence of more than one internal mode, we expect complicated coupling between their amplitudes, making the behaviour at large times very difficult to predict. However, if we restrict to odd initial data, the solution to (2.18) remains odd. This means that the even internal mode can never be excited, so the system effectively has only one internal mode. In this case, the analysis in Section 2.4.1 still applies, since the frequency of the odd internal mode always lies in the range $(m/2, m)$, so we expect its amplitude to decay like $s^{-1/2}$.

For general initial data and wormhole radii $a > a_1$, we cannot produce a concrete conjecture about the decay rate. However, we expect the behaviour of the system to

depend on the locations of the two internal mode frequencies within the mass gap. The analysis in Section 2.4.1 for a single internal mode suggests that for frequencies less than $m/2$, a real contribution to \dot{A} does not appear until at least $\mathcal{O}(A^4)$. In this case, we require another change of variable in the radiation equation to rule out the contribution of the radiative term at higher order. We then expect to solve an equivalent of (2.28) where $k + l = 3$ to find the required change of variable. If $9\omega^2 < m^2$ so that the solution is still real, we can proceed by induction, changing the variable until the solution is complex. A real contribution to (2.31) will only be obtained for $k + l = N$ such that $N^2\omega^2 > m^2$. This would mean a real contribution to \dot{A} at $\mathcal{O}(A^{N+1})$, and hence result in a decay rate of $s^{-1/N}$. Further detail can be found in [6].

Although the presence of a second internal mode complicates the dynamics, we still expect the smallest N such that the even internal mode frequency ω satisfies $N^2\omega^2 > m^2$ to be an important factor in the behaviour at large s . The axis ticks in Figure 2.6 show the value of N for a range of a .

Chapter 3

The projective to Einstein correspondence $N \rightarrow M$

The purpose of this chapter is to introduce the preliminaries that are required to understand the remainder of the thesis. We will first review projective geometry, including the Cartan and tractor bundles associated with a projective structure, before moving on to the projective to Einstein correspondence of [28]. We begin with some notation and conventions.

Notation and conventions

- We use \mathbb{R}^n to mean the real vector space of dimension n , and \mathbb{R}_n to mean its dual. We think of vectors in \mathbb{R}^n as column vectors, and vectors in \mathbb{R}_n as row vectors.
- We will projectivise vector spaces by *line projectivisation*, that is, we will take the space of *unoriented* lines through the origin, unless stated otherwise.
- When we refer to the projective to Einstein correspondence, the projective manifold will be called N and will have dimension n , whilst the Einstein manifold will be called M and will have dimension $2n$.
- We use the letter π for maps to N , and the letter κ for maps to M . We will attach subscripts to π and κ to give information about the preimage or the significance of the map.
- We use lower case Latin indices $i, j, k, \dots = 1, \dots, n$ for tensorial objects on N and $a, b, c, \dots = 1, \dots, 2n$ for tensorial objects on M .

- We use \odot and \wedge to denote the symmetrised and antisymmetrised tensor product respectively. That is,

$$A \odot B = \frac{1}{2}(A \otimes B + B \otimes A), \quad A \wedge B = \frac{1}{2}(A \otimes B - B \otimes A).$$

We also occasionally write A^2 for $A \odot A$, and in Chapter 5 we will sometimes omit the \odot altogether.

- Where indices have been symmetrised or antisymmetrised over, we will enclose them in round or square brackets respectively.
- Our conventions for differential forms are

$$(d\omega)_{ab\dots c} = \partial_{[a}\omega_{b\dots c]}, \quad (\eta \wedge \omega)_{a\dots d} = \eta_{[a\dots b}\omega_{c\dots d]},$$

$$\omega = \omega_{a\dots b} dx^a \wedge \dots \wedge dx^b, \quad F_{ab} dx^a \wedge dx^b = F_{[ab]} dx^a \otimes dx^b.$$

- We use \lrcorner to denote a contraction between a vector and a form.
- The Riemann curvature tensor $R_{abc}{}^d$ of a connection ∇_a is defined by

$$(\nabla_a \nabla_b - \nabla_b \nabla_a)X^d = R_{abc}{}^d X^c,$$

where X is any vector field.

- The Ricci tensor is defined by the contraction $R_{bc} = R_{abc}{}^a$.

3.1 Projective Geometry

Our discussion follows Eastwood [31].

Definition 3.1.1. *A projective structure $(N, [\nabla])$ on a manifold N is an equivalence class $[\nabla]$ of torsion-free affine connections on N which have the same geodesics as unparametrised curves.*

The following proposition converts definition 3.1.1 to a more operational form.

Proposition 3.1.2. *Two torsion-free connections ∇ and $\bar{\nabla}$ belong to the same projective class if and only if their components Γ_{jk}^i and $\bar{\Gamma}_{jk}^i$ are related by*

$$\bar{\Gamma}_{jk}^i - \Gamma_{jk}^i = \delta_j^i \Upsilon_k + \delta_k^i \Upsilon_j \tag{3.1}$$

for some one-form Υ .

Proof. We denote by \mathcal{V} the vertical sub-bundle of $T(TN)$, where $\pi_T : TN \rightarrow N$ is the tangent bundle to N . A connection defines a splitting of the exact sequence

$$0 \longrightarrow \mathcal{V} \longrightarrow T(TN) \longrightarrow \pi_T^* TN \longrightarrow 0 \quad (3.2)$$

so that each $\xi \in T_x N$ has a unique pull-back in the horizontal sub-bundle complementary to \mathcal{V}_x . The integral curves of these pull-backs, when projected down to N , then define the geodesics of the connection.

Any two connections are related by some $\delta\Gamma_{ij}^k$, which satisfies $\delta\Gamma_{ij}^k = \delta\Gamma_{(ij)}^k$ as long as both connections are torsion-free. A change of connection is equivalent to a change in the splitting of (3.2). At $\xi \in T_x N$, the change is given by the homomorphism from $T_x N$ to $T_x N = \mathcal{V}_x$ defined by the contraction $\xi^i \Gamma_{ij}^k$. Thus the two connections define the same geodesics if and only if $\xi^i \xi^j \Gamma_{ij}^k$ is a multiple of ξ^k for all ξ^i . This is true if and only if there is a one-form Υ_i such that (3.1) is satisfied.¹

□

One can show that the curvature of a connection ∇ in the projective class can be uniquely decomposed as

$$R_{ijk}^l = W_{ijk}^l + 2\delta_{[i}^l P_{j]k} - 2P_{[ij]}\delta_k^l, \quad (3.3)$$

where the Weyl projective curvature tensor, W_{ijk}^l , is trace free, and the Schouten tensor, P_{ij} , is given in terms of the Ricci tensor by

$$P_{ij} = \frac{1}{n-1} R_{(ij)} + \frac{1}{n+1} R_{[ij]}.$$

The objects W_{ijk}^l and P_{ij} transform as

$$\overline{W}_{ijk}^l = W_{ijk}^l, \quad \overline{P}_{ij} = P_{ij} - \nabla_i \Upsilon_j + \Upsilon_i \Upsilon_j \quad (3.4)$$

under a change of representative connection (3.1). Note that for $n = 2$ the Weyl tensor always vanishes.

A projective structure in dimension n is said to be flat if it is diffeomorphic to the real projective space \mathbb{RP}^n with its standard flat projective structure.

¹To see this, take some one-form ω_i and note that $2\xi^i \xi^j \delta_{(i}^k \Upsilon_{j)} \omega_k$ vanishes if and only if $\xi^k \omega_k$ does.

Definition 3.1.3. *The real projective space \mathbb{RP}^n of dimension n is the space of unoriented lines through the origin in \mathbb{R}^{n+1} , thought of as $\mathbb{RP}^n = (\mathbb{R}^{n+1} \setminus \{0\})/\mathbb{R}^*$, where the quotient identifies points $P \in \mathbb{R}^{n+1}$ under the equivalence relation*

$$(P^0, \dots, P^n) \sim (cP^0, \dots, cP^n) \quad \forall c \in \mathbb{R}^*.$$

The geodesics on \mathbb{RP}^n are given by planes through the origin in \mathbb{R}^{n+1} under the projection $\pi_{\mathbb{P}} : \mathbb{R}^{n+1} \rightarrow \mathbb{RP}^n$.

Remark 3.1.4. Let P denote a non-zero point in \mathbb{R}^{n+1} with coordinates $(P^0, \dots, P^n)^T$, and let $[P]$ denote the corresponding point in \mathbb{RP}^n , labelled by homogeneous coordinates. In a patch \mathcal{U}_0 where $P^0 \neq 0$, we can write $[P] = [1, P^1/P^0, \dots, P^n/P^0]^T$ and define inhomogeneous coordinates on \mathbb{RP}^n by

$$(x^1, \dots, x^n) = (P^1/P^0, \dots, P^n/P^0).$$

If we combine this with coordinate patches \mathcal{U}_i where $P^i \neq 0$, $i = 1, \dots, n$, we can build an atlas for \mathbb{RP}^n .

Remark 3.1.5. The flat projective structure on \mathbb{RP}^n has a special duality property which we now discuss. Consider the set of hyperplanes through the origin in \mathbb{R}^{n+1} . These can be specified by their normal vector, which is defined only up to multiplication by \mathbb{R}^* . Let us denote such a hyperplane by a non-zero row vector $L \in \mathbb{R}_{n+1}$. A point $P \in \mathbb{R}^{n+1}$ lies in the hyperplane defined by L if and only if $L \cdot P = 0$.

When we projectivise the \mathbb{R}^{n+1} , any $P \neq 0$ descends to a point $[P] \in \mathbb{RP}^n$, and any hyperplane descends to a hypersurface $[L] \subset \mathbb{RP}^n$. The incidence relation $L \cdot P = 0$ is now equivalent to the point $[P]$ lying in the hypersurface $[L]$. The homogeneous coordinates $[L]$ parametrise a second projective space which we think of as the dual to the \mathbb{RP}^n parametrised by $[P]$, and denote \mathbb{RP}_n .

Remark 3.1.6. Real projective space can be viewed as homogeneous space as follows. The group $SL(n+1, \mathbb{R})$ acts from the left via the fundamental representation on coordinates $(P^0, \dots, P^n)^T$ in \mathbb{R}^{n+1} , and this descends to a transitive action on \mathbb{RP}^n . By the orbit stabiliser theorem, $\mathbb{RP}^n = SL(n+1, \mathbb{R})/S$, where S is a subgroup stabilising a point. If we choose the point $[1, 0, \dots, 0]^T$, the elements of S are matrices of the general form

$$\begin{pmatrix} \det a^{-1} & b \\ 0 & a \end{pmatrix}$$

for some $a \in GL(n, \mathbb{R})$ and $b \in \mathbb{R}_n$.

Remark 3.1.7. The necessary and sufficient condition for flatness of a projective structure depends on the dimension of the manifold on which it is defined. In dimension $n > 2$, a projective structure is flat if and only if its Weyl projective curvature tensor vanishes. However, for $n = 2$ the projective Weyl tensor is always vanishing. It can be shown that a projective structure on a *surface* is flat if and only if the Cotton tensor $\nabla_{[i}P_{j]k}$ vanishes for any choice of representative connection.

3.1.1 The Cartan bundle

One way of understanding the construction in [28] is via the Cartan bundle [20] of the projective structure $(N, [\nabla])$ (see also [43, 66]). Cartan geometries generalise Klein's Erlangen programme [41], a study of homogeneous spaces G/S , to the curved case, in which the total space G is replaced by a principal right S -bundle over a manifold N such that the tangent space to N at every point is isomorphic to the Lie algebra quotient $\mathfrak{g}/\mathfrak{s}$. Since projective structures are modelled on \mathbb{RP}^n , which can be viewed as a homogeneous space, they constitute a type of Cartan geometry.

In the Riemannian case, the model space is $\mathbb{R}^n \cong \text{Euc}(n)/SO(n)$. The corresponding Cartan geometry is a general, curved Riemannian manifold. One has an obvious subclass of frames which are "adapted" to the metric, i.e. those which are orthonormal. We can thus think of a curved Riemannian manifold as a principal $SO(n)$ bundle whose tangent spaces are modelled on $\mathbb{R}^n \cong \mathfrak{Euc}(n)/\mathfrak{so}(n)$. We say that Riemannian manifolds are Cartan geometries of type $(\text{Euc}(n), SO(n))$.

The theory of Cartan geometries was developed as part of Cartan's *method of moving frames*. The idea is to pick out some adapted frames for manifolds equipped with some non-metric structure. The bundle of such frames over a manifold is then a principal bundle $\pi_{\mathcal{G}} : \mathcal{G} \rightarrow N$ with structure group S .

The bundle \mathcal{G} is equipped with a \mathfrak{g} -valued one-form θ called the Cartan connection. It defines an isomorphism $\theta : T_u \mathcal{G} \rightarrow \mathfrak{g}$ at every point $u \in \mathcal{G}$ such that the vertical subspace $\mathcal{V}_u \mathcal{G} \subset T_u \mathcal{G}$ is mapped to \mathfrak{s} and the horizontal subspace $\mathcal{H}_u \mathcal{G} \subset T_u \mathcal{G}$ is defined as the inverse image of $\mathfrak{g}/\mathfrak{s}$. Note that it is not a connection in the usual sense of a principal bundle connection, since it takes value in a Lie algebra larger than that of the structure group. Further details can be found in [66].

In the projective case, if we choose the point which is stabilised by S to be $[1, 0, \dots, 0]$, the Cartan connection can be written as a matrix

$$\theta = \begin{pmatrix} -\text{tr}\phi & \eta \\ \omega & \phi \end{pmatrix}, \quad (3.5)$$

where ω , η and ϕ are one-forms valued in \mathbb{R}^n , \mathbb{R}_n and $\mathfrak{gl}(n, \mathbb{R})$ respectively. We will refer to the components of ω and η with respect to the natural basis of $\mathfrak{sl}(n+1, \mathbb{R})$ as $\{\omega^{(i)}\}$ and $\{\eta_{(i)}\}$, so that $\omega^{(i)}$ and $\eta_{(i)}$ are both one-forms taking values in \mathbb{R} .

Definition 3.1.8. *The Cartan geometry of a projective structure $(N, [\nabla])$ consists of a principal right S -bundle $\pi_{\mathcal{G}} : \mathcal{G} \rightarrow N$, where the right-action of some $s \in S$ on \mathcal{G} is denoted by R_s , and a one-form θ on \mathcal{G} called the Cartan connection, which takes values in $\mathfrak{sl}(n+1, \mathbb{R})$. The Cartan connection can be written in the form (3.5) and has the following properties:*

1. $\theta_u : T_u \mathcal{G} \rightarrow \mathfrak{sl}(n+1, \mathbb{R})$ is an isomorphism for all $u \in \mathcal{G}$;
2. $\theta(\xi_{\mathfrak{v}}) = \mathfrak{v}$ for all fundamental vector fields $\xi_{\mathfrak{v}}$ on \mathcal{G} ;
3. $R_s^* \theta = \text{Ad}(s^{-1})\theta = s^{-1}\theta s$ for all $s \in S$.
4. If ξ is a vector field on \mathcal{G} with the property that $\eta(\xi) = \phi(\xi) = 0$ and $\omega(\xi) \in \mathbb{R}^n \setminus \{0\}$, then the integral curve of ξ projects down to a geodesic on N and conversely every geodesic of $[\nabla]$ arises in this way.
5. The $\mathfrak{sl}(n+1, \mathbb{R})$ -valued curvature two-form Θ satisfies

$$\Theta = d\theta + \theta \wedge \theta = \begin{pmatrix} 0 & L(\omega \wedge \omega) \\ 0 & W(\omega \wedge \omega) \end{pmatrix}, \quad (3.6)$$

where L and W are smooth curvature functions valued in $\text{Hom}(\mathbb{R}^n \wedge \mathbb{R}^n, \mathbb{R}_n)$ and $\text{Hom}(\mathbb{R}^n \wedge \mathbb{R}^n, \mathbb{R}_n \otimes \mathbb{R}^n)$ respectively. The function W represents the Weyl projective curvature tensor appearing in (3.3).

Remark 3.1.9. The Cartan geometry of a projective structure is unique in the sense that for any two Cartan geometries $(\hat{\pi}_{\mathcal{G}} : \hat{\mathcal{G}} \rightarrow N, \hat{\theta})$ and $(\pi_{\mathcal{G}} : \mathcal{G} \rightarrow N, \theta)$ of type $(SL(n+1, \mathbb{R}), S)$ satisfying the above properties there is a S -bundle isomorphism $\nu : \mathcal{G} \rightarrow \hat{\mathcal{G}}$ such that $\nu^* \hat{\theta} = \theta$.

Remark 3.1.10. For every open set $\mathcal{U} \subset N$, projective vector fields on \mathcal{U} are in one-to-one correspondence with vector fields on $\pi_{\mathcal{G}}^{-1}(\mathcal{U})$ which preserve θ and are equivariant under the principal S -action.

3.1.2 Tractor bundles

The Cartan connection also gives us a unique connection on any bundle associated to \mathcal{G} via some S -module. In particular, let \mathcal{B} be a vector space and $\rho_{\mathcal{B}} : S \rightarrow GL(\mathcal{B})$ a

representation of S acting on \mathcal{B} . We can construct an *associated bundle*

$$\pi_{\mathcal{B}} : \mathcal{G} \times_{\rho_{\mathcal{B}}} \mathcal{B} \rightarrow N$$

where points in $\mathcal{G} \times_{\rho_{\mathcal{B}}} \mathcal{B}$ are equivalence classes of pairs $[u, v]$, where $u \in \mathcal{G}$ and $v \in \mathcal{B}$, up to the equivalence relation

$$(u_1, v_1) \sim (u_2, v_2) \Leftrightarrow \exists s \text{ such that } u_2 = u_1 s, v_2 = \rho_{\mathcal{B}}(s^{-1})v_1.$$

We thus obtain a vector bundle over N whose fibres are diffeomorphic to \mathcal{B} . A section $\tilde{\sigma} : N \rightarrow \mathcal{G} \times_{\rho_{\mathcal{B}}} \mathcal{B}$ is represented by a map $\sigma : \mathcal{G} \rightarrow \mathcal{B}$ which is equivariant in the sense that $\sigma(us) = \rho_{\mathcal{B}}(s^{-1})\sigma(u)$ for all $s \in S$. Importantly, any such bundle inherits a connection from the Cartan connection θ on \mathcal{G} . The concept of an associated bundle applies to any principal bundle, but we call vector bundles which are associated to a Cartan bundle *tractor* bundles, and the connections that they inherit from the Cartan connection are called *tractor connections*.

A particularly important example of a vector bundle associated to \mathcal{G} is the *cotractor bundle*, which is defined by the canonical action of S on \mathbb{R}_{n+1} given by $(s, L) \mapsto Ls^{-1}$. We call this bundle $\pi_{\mathcal{T}} : \mathcal{T}^* \rightarrow N$. In order to describe its connection, we consider a section represented by $\sigma : \mathcal{G} \rightarrow \mathbb{R}_{n+1}$ and define the one-form

$$d\sigma - \sigma\theta. \tag{3.7}$$

This turns out to be a *semi-basic*² one-form satisfying

$$R_s^*(d\sigma - \sigma\theta) = (d\sigma - \sigma\theta)s,$$

making $\sigma \mapsto d\sigma - \sigma\theta$ an equivariant connection on \mathcal{T}^* .

Although this construction of \mathcal{T}^* relies on the Cartan bundle, it is possible to construct it independently. In order to do so we need the notion of a *projective density*.

²Recall that a semi-basic form on a fibre bundle $\mathcal{G} \rightarrow N$ is a form which is a linear combination, with coefficients parametrised by the fibres, of basic forms on \mathcal{G} (i.e. forms which are the pull-backs of forms on N).

Projective densities

From the projective change of connection (3.1) we can derive the corresponding change in $\nabla\chi$ for some m -form χ on N :

$$\bar{\nabla}_i \chi_{jk\dots l} = \nabla_i \chi_{jk\dots l} - (m+1)\Upsilon_i \chi_{jk\dots l} - (m+1)\Upsilon_{[i} \chi_{jk\dots l]}. \quad (3.8)$$

In particular, for a volume form ($m = n$) we find

$$\bar{\nabla}_i \chi_{jk\dots l} = \nabla_i \chi_{jk\dots l} - (n+1)\Upsilon_i \chi_{jk\dots l},$$

where the final term in (3.8) has vanished because it contains a symmetrisation over $n+1$ indices. We can write this in a more compact way as

$$\bar{\nabla}_i \chi = \nabla_i \chi - (n+1)\Upsilon_i \chi.$$

Note that for sections τ of the bundle $\mathcal{E}(w) := (\Lambda^n)^{-w/(n+1)}$ we have

$$\bar{\nabla}_i \tau = \nabla_i \tau + w\Upsilon_i \tau. \quad (3.9)$$

We called such sections *projective densities of weight w* , and for any vector bundle $\mathcal{B} \rightarrow N$ we write $\mathcal{B}(w)$ for the tensor product of \mathcal{B} with $\mathcal{E}(w)$. For example, $T^*N(w)$ is the bundle of one-forms with projective weight w , and for sections μ_i of $T^*N(w)$ we have

$$\bar{\nabla}_i \mu_j = \nabla_i \mu_j + (w-1)\Upsilon_i \mu_j - \Upsilon_j \mu_i. \quad (3.10)$$

The cotractor bundle

We can now define the cotractor bundle $\pi_{\mathcal{T}} : \mathcal{T}^* \rightarrow N$. For a choice of connection in the projective class we identify

$$\mathcal{T}^* = \mathcal{E}(1) \oplus T^*N(1), \quad (3.11)$$

so that a section can be represented by a pair

$$\begin{pmatrix} \tau \\ \mu_i \end{pmatrix}. \quad (3.12)$$

Under a change of projective connection (3.1), this splitting changes according to

$$\overline{\begin{pmatrix} \tau \\ \mu_i \end{pmatrix}} = \begin{pmatrix} \tau \\ \mu_i + \Upsilon_i \tau \end{pmatrix}. \quad (3.13)$$

Note the exact sequence

$$0 \longrightarrow T^*N(1) \longrightarrow \mathcal{T}^* \xrightarrow{V} \mathcal{E}(1) \longrightarrow 0, \quad (3.14)$$

where we call the map V the projective *canonical tractor*³. A choice of connection in the projective class defines a splitting (3.11) of (3.14).

The bundle \mathcal{T}^* admits a projectively invariant *tractor connection* given by

$$\nabla_i^{\mathcal{T}} \begin{pmatrix} \tau \\ \mu_j \end{pmatrix} = \begin{pmatrix} \nabla_i \tau - \mu_i \\ \nabla_i \mu_j + P_{ij} \tau \end{pmatrix}, \quad (3.15)$$

where ∇ is the choice of projective connection and P_{ij} is its Schouten tensor. This turns out to agree with (3.7). Under a change of projective connection (3.1), we find

$$\begin{aligned} \overline{\nabla}_i^{\mathcal{T}} \overline{\begin{pmatrix} \tau \\ \mu_j \end{pmatrix}} &= \overline{\nabla}_i^{\mathcal{T}} \begin{pmatrix} \tau \\ \mu_j + \Upsilon_j \tau \end{pmatrix} \\ &= \begin{pmatrix} \overline{\nabla}_i \tau - (\mu_i + \Upsilon_i \tau) \\ \overline{\nabla}_i(\mu_j + \Upsilon_j \tau) + \overline{P}_{ij} \tau \end{pmatrix} \\ &= \begin{pmatrix} \nabla_i \tau + \Upsilon_i \tau - (\mu_i + \Upsilon_i \tau) \\ \nabla_i(\mu_j + \Upsilon_j \tau) - \Upsilon_j(\mu_i + \Upsilon_i \tau) + (P_{ij} - \nabla_i \Upsilon_j + \Upsilon_i \Upsilon_j) \tau \end{pmatrix}, \end{aligned}$$

where we have used (3.13) in the first line, (3.15) in the second and (3.9), (3.10) and (3.4) in the third. After some cancellation, we identify

$$\begin{aligned} \overline{\nabla}_i^{\mathcal{T}} \overline{\begin{pmatrix} \tau \\ \mu_j \end{pmatrix}} &= \begin{pmatrix} \nabla_i \tau - \mu_i \\ \nabla_i \mu_j + \Upsilon_j \nabla_i \tau - \Upsilon_j \mu_i + P_{ij} \tau \end{pmatrix} \\ &= \begin{pmatrix} \nabla_i \tau - \mu_i \\ \nabla_i \mu_j + P_{ij} \tau \end{pmatrix} = \nabla_i^{\mathcal{T}} \begin{pmatrix} \tau \\ \mu_j \end{pmatrix} \end{aligned}$$

³In fact V is a section of a bundle $\mathcal{T}(1)$, where \mathcal{T} can be identified with a direct sum $\mathcal{E}(1) \oplus TN(1)$ given a choice of connection in the projective class. The natural pairing between \mathcal{T} and \mathcal{T}^* defines the map $V : \mathcal{T}^* \rightarrow \mathcal{E}(1)$.

using (3.15) and the change of splitting (3.13) adapted to the tensor product of T^*N and \mathcal{T}^* of which the derivative is a section.

Any tensor product of \mathcal{T} , \mathcal{T}^* and $\mathcal{E}(1)$ is equipped with a tractor connection which is inherited from the connection on the standard cotractor bundle. Equivalently, any such tensor product can be thought of as an associated vector bundle to the Cartan bundle \mathcal{G} , with its connection inherited from the Cartan connection via the corresponding representation of S . It is this connection, with its special equivariance property, that allows us to construct an Einstein metric as an invariant of the projective structure. This construction is the subject of the following section.

3.2 The projective to Einstein correspondence

Consider a quotient of the total space \mathcal{G} of the Cartan bundle by $GL(n, \mathbb{R})$, which is embedded in S in the obvious way:

$$GL(n, \mathbb{R}) \ni a \longmapsto \begin{pmatrix} \det a^{-1} & 0 \\ 0 & a \end{pmatrix} \in S. \quad (3.16)$$

It is easily verified that

$$\begin{pmatrix} \det a^{-1} & 0 \\ 0 & a \end{pmatrix} \begin{pmatrix} 0 & \eta \\ \omega & 0 \end{pmatrix} \begin{pmatrix} \det a^{-1} & 0 \\ 0 & a \end{pmatrix}^{-1} = \begin{pmatrix} 0 & \eta a^{-1} \det a^{-1} \\ (\det a) a \omega & 0 \end{pmatrix},$$

for any $a \in GL(n, \mathbb{R})$, meaning that due to the equivariance property of the Cartan connection, the natural contraction $\eta\omega := \sum_i \eta_{(i)} \otimes \omega^{(i)}$ defined by θ is preserved by the adjoint action of this $GL(n, \mathbb{R})$ subgroup. It thus descends to a naturally defined object on the quotient $M = \mathcal{G}/GL(n, \mathbb{R})$.

Theorem 3.2.1. [28] *There exist a metric and two-form (g, Ω) on $M = \mathcal{G}/GL(n, \mathbb{R})$ such that the quotient map $\kappa_q : P \rightarrow M$ gives*

$$\kappa_q^* g = \text{Sym}(\eta\omega) \quad (3.17)$$

$$\kappa_q^* \Omega = \text{Ant}(\eta\omega), \quad (3.18)$$

where Sym and Ant denote the symmetric and anti-symmetric parts of the $(0, 2)$ tensor $\eta\omega$. Moreover, Ω is closed as a consequence of the Bianchi identity satisfied by the curvature two-form (3.6), g is Einstein with non-zero scalar curvature, and the two

are related by an endomorphism J satisfying $J^2 = \text{Id}$. Hence (g, Ω) is an almost para-Kähler structure on M .

Remark 3.2.2. The full proof of Theorem 3.2.1 only appears explicitly in [28] in the case $n = 2$, although it can be generalised to $n > 2$. This generalisation is discussed in their appendix. They show that the Ricci scalar of g is 24 in the case $n = 2$. In Chapter 4 we will need the Ricci scalar for general n . We will calculate this under the assumption (stated without proof in [28]) that g is Einstein.

Remark 3.2.3. The quotient M turns out to be an affine bundle over N with structure group S , i.e. S acts affinely on the fibres of $\pi_M : M \rightarrow N$, and sections of this bundle are in one-to-one correspondence with representative connections $\nabla \in [\nabla]$. This means that given some choice of connection $\nabla \in [\nabla]$ we have a diffeomorphism $\kappa_A : T^*N \rightarrow M$ with which we can pull back the pair (g, Ω) . In canonical local coordinates (x^i, ζ_i) on the cotangent bundle, we find

$$\begin{aligned}\kappa_A^* g &= d\zeta_i \odot dx^i - (\Gamma_{ij}^k \zeta_k - \zeta_i \zeta_j - P_{ij}) dx^i \odot dx^j, \\ \kappa_A^* \Omega &= d\zeta_i \wedge dx^i + P_{ij} dx^i \wedge dx^j, \quad i, j = 1, \dots, n.\end{aligned}\tag{3.19}$$

Here Γ_{jk}^i are the connection components of the representative connection ∇ that we chose, and its Schouten tensor is denoted P_{ij} . This can be shown to be projectively invariant in the sense that a different choice of $\nabla \in [\nabla]$ corresponds to shifting the fibre coordinates ζ_i , i.e. metrics on T^*N resulting from pulling back g using different representative connections are isometric. Explicitly, a projective transformation (3.1) corresponds to a change

$$\zeta_i \rightarrow \zeta_i + \Upsilon_i. \tag{3.20}$$

Remark 3.2.4. In fact, the metric and symplectic form (3.19) turn out to belong to a one-parameter family $\{(g_\Lambda, \Omega_\Lambda) ; \Lambda \neq 0\}$, which can be written in local coordinates as

$$g_\Lambda = d\zeta_i \odot dx^i - (\Gamma_{ij}^k \zeta_k - \Lambda \zeta_i \zeta_j - \Lambda^{-1} P_{ij}) dx^i \odot dx^j \tag{3.21}$$

$$\Omega_\Lambda = d\zeta_i \wedge dx^i + \frac{1}{\Lambda} P_{ij} dx^i \wedge dx^j, \quad i, j = 1, \dots, n. \tag{3.22}$$

Metrics of the form (3.21) are a subclass of so-called Osserman metrics. More details can be found in [16]. They are all Einstein with non-zero scalar curvature 24Λ , but for $\Lambda \neq 1$ the relation to projective geometry is lost. For the remainder of the thesis we will write g for $g_{\Lambda=1}$ unless stated otherwise. Note that $\{g_\Lambda\}$ will be the subject of Chapter 4, whilst in Chapters 5 and 6 we will restrict our attention to g because the projective geometry is a key aspect of the content of these chapters.

Remark 3.2.5. One could also consider taking a quotient of \mathcal{G} by a different subgroup of S . The \mathbb{R}^* bundle over M which we will discuss in Chapter 4 will turn out to be a quotient of \mathcal{G} by $SL(n, \mathbb{R})$.

Remark 3.2.6. As mentioned above, in the special case where the $(N, [\nabla])$ is a projective surface, M has dimension four, and so anti-self-duality is defined. It turns out that both the symplectic form Ω and the conformal curvature of g are ASD. Both of these facts will play an important role in Chapter 5.

Remark 3.2.7. Note that an endomorphism J which squares to the identity defines two n -dimensional sub-bundles of the tangent bundle TM defined at each $m \in M$ as the vector subspaces of $T_m M$ which have eigenvalues ± 1 with respect to J . These sub-bundles form a pair of smooth distributions D_{\pm} in TM . Further discussion of the endomorphism J will appear in Chapter 6.

3.2.1 Symmetries of M

Recall that a projective vector field on any manifold with a connection generates a one-parameter family of transformations which preserve the geodesics of that connection up to parametrisation. Projective vector fields thus naturally arise as the symmetries of a projective structure. Explicitly, a vector field \widehat{K} is projective if it satisfies

$$\mathcal{L}_{\widehat{K}} \Gamma_{ij}^k = \delta_i^k \Upsilon_j + \delta_j^k \Upsilon_i \quad (3.23)$$

for some 1-form Υ , where Γ_{ij}^k are the connection components, and their Lie derivative is defined by⁴

$$\mathcal{L}_{\widehat{K}} \Gamma_{ij}^k \equiv \frac{\partial^2 \widehat{K}^k}{\partial x^i \partial x^j} + \widehat{K}^m \frac{\partial \Gamma_{ij}^k}{\partial x^m} - \Gamma_{ij}^m \frac{\partial \widehat{K}^k}{\partial x^m} + \Gamma_{im}^k \frac{\partial \widehat{K}^m}{\partial x^j} + \Gamma_{mj}^k \frac{\partial \widehat{K}^m}{\partial x^i}. \quad (3.24)$$

One consequence of the symmetry property of the Cartan connection discussed in remark 3.1.10 is that for every open set $\mathcal{U} \subset N$ we have an isomorphism between the Lie algebra of projective vector fields on \mathcal{U} and the Lie algebra of vector fields on $\pi_{\mathcal{G}}^{-1}(\mathcal{U})$ preserving the natural contraction $\eta\omega$. Such vector fields must descend to vector fields on $\pi_M^{-1}(\mathcal{U})$ preserving (g, Ω) . In fact, it can be shown that every Killing vector field of (M, g_Λ) is also symplectic with respect to Ω_Λ and is therefore the lift of a projective vector field on $(N, [\nabla])$.

⁴Despite the fact that connection components are not tensorial objects, one can still define their Lie derivative with respect to a vector \widehat{K} by considering how they transform when one moves infinitesimally along the curve defined by \widehat{K} . See [85] for further details.

Explicitly, for every projective vector field \widehat{K} of $(N, [\nabla])$ there is a corresponding symmetry K of $(M, g_\Lambda, \Omega_\Lambda)$ given in local coordinates by

$$K = \widehat{K} - \zeta_i \frac{\partial \widehat{K}^j}{\partial x^i} \frac{\partial}{\partial \zeta_j} + \frac{1}{\Lambda} \Upsilon_i \frac{\partial}{\partial \zeta_i}, \quad (3.25)$$

where Υ_i is defined by (3.23).

3.2.2 Tractor perspective

From the tractor perspective, the space M will turn out to be the projectivised cotractor bundle of N with an \mathbb{RP}^{n-1} sub-bundle removed from each fibre. We can understand what this \mathbb{RP}^{n-1} sub-bundle is as follows.

On the total space of \mathcal{T}^* we pull back $\pi_{\mathcal{T}} : \mathcal{T}^* \rightarrow N$ along $\pi_{\mathcal{T}}$ to get $\pi_{\mathcal{T}}^*(\mathcal{T}^*) \rightarrow \mathcal{T}^*$ as a vector bundle over the total space \mathcal{T}^* . By construction this bundle has a tautological section $W \in \Gamma(\pi_{\mathcal{T}}^*(\mathcal{T}^*))$. We also have $\pi_{\mathcal{T}}^*(\mathcal{T}(w))$ for any weight w , and we shall write simply $V \in \Gamma(\pi_{\mathcal{T}}^*(\mathcal{T}(1)))$ for the pull back to \mathcal{T}^* of the canonical tractor V on N .

Now define

$$\kappa_{\mathbb{P}} : \mathcal{T}^* \longrightarrow \mathcal{M} := \mathbb{P}(\mathcal{T}^*) \quad (3.26)$$

by the fibre-wise projectivisation, and use $\pi_{\mathcal{M}}$ for the map

$$\pi_{\mathcal{M}} : \mathcal{M} \rightarrow N.$$

We denote by $\mathcal{E}_{\mathcal{T}^*}(w')$, for $w' \in \mathbb{R}$, the line bundle on $\mathbb{P}(\mathcal{T}^*)$ whose sections correspond to functions $f : \pi_{\mathcal{T}}^*\mathcal{T}^* \rightarrow \mathbb{R}$ that are homogeneous of degree w' in the fibres of $\pi_{\mathcal{T}}^*\mathcal{T}^* \rightarrow \mathbb{P}(\mathcal{T}^*)$. For any weight w we also have $\mathcal{E}(w)$ on N and its pull back to the bundle $\pi_{\mathcal{M}}^*\mathcal{E}(w) \rightarrow \mathbb{P}(\mathcal{T}^*)$. We define the product of these two density bundles on \mathcal{M} as

$$\mathcal{E}(w, w') := \pi_{\mathcal{T}}^*\mathcal{E}(w) \otimes \mathcal{E}_{\mathcal{T}^*}(w').$$

On \mathcal{T}^* there is a canonical density $\tau \in \Gamma(\pi_{\mathcal{T}}^*\mathcal{E}(1))$ given by

$$\tau := V \lrcorner W,$$

Note that τ is homogeneous of degree 1 up the fibres of the map $\kappa_{\mathbb{P}} : \mathcal{T}^* \rightarrow \mathcal{M}$. Thus τ determines, and is equivalent to, a section (that we also denote) τ of the density bundle $\mathcal{E}(1, 1)$. So \mathcal{M} is stratified according to whether or not τ is vanishing, and we

write $\mathcal{Z}(\tau)$ to denote, in particular, the zero locus of τ . We will show in Chapter 6 that M can be identified with $\mathcal{M} \setminus \mathcal{Z}(\tau)$.

3.2.3 The model case

When $(N, [\nabla])$ is the flat projective structure \mathbb{RP}^n , the metric and symplectic form (3.19) reduce to

$$g = d\zeta_i \odot dx^i + \zeta_i \zeta_j dx^i \odot dx^j, \quad \Omega = d\zeta_i \wedge dx^i, \quad i, j = 1, \dots, n. \quad (3.27)$$

In this case the Cartan bundle of N is just $SL(n+1, \mathbb{R})$, and M is simply the Lie group quotient $SL(n+1, \mathbb{R})/GL(n, \mathbb{R})$. The cotractor bundle has zero curvature, and although it is not trivial, the restriction of $\mathbb{P}(\mathcal{T}^*)$ to the set $\mathcal{Z}(\tau) \neq 0$ is. We can thus write M as

$$M = \{([P], [L]) \in \mathbb{RP}^n \times \mathbb{RP}_n \mid P \cdot L \neq 0\}.$$

As discussed in Section 3.1, a point $[L] \in \mathbb{RP}_n$ represents a line $[L] \subset \mathbb{RP}^n$ which passes through $P \in \mathbb{RP}^n$ if and only if $L \cdot P = 0$. In Chapter 5 we will show that for $n = 2$, the conformal structure on M can be obtained by demanding that two pairs $([P], [L])$ and $([\tilde{P}], [\tilde{L}])$ are null-separated if there exists a line which contains the three points $([P], [\tilde{P}], [L] \cap [\tilde{L}])$.

We also find in the model case that the symplectic form Ω is parallel with respect to the Levi–Civita connection $g\nabla$ of g , meaning that the endomorphism J of TM which relates g and Ω is also parallel. As a result of this, the distributions D_{\pm} defined by the two n –dimensional eigen–bundles of J are parallel in the sense that $g\nabla_{\xi_1} \xi_2 \in \Gamma(D_{\pm})$ for all $\xi_1 \in \Gamma(TM)$, $\xi_2 \in \Gamma(D_{\pm})$. This makes the distributions *Frobenius integrable*, meaning that the Lie bracket of any two sections of D_{\pm} is also a section of D_{\pm} , or equivalently (as shown by Frobenius) that each of the two distributions is tangent to a foliation by sub–manifolds of dimension n at every point.

To see that a parallel distribution is necessarily Frobenius integrable, note that the Lie bracket can be written

$$[\xi_1, \xi_2] = \nabla_{\xi_1} \xi_2 - \nabla_{\xi_2} \xi_1 \in \Gamma(D) \quad \text{for all } \xi_1, \xi_2 \in \Gamma(D),$$

where D is a distribution which is parallel with respect to a connection ∇ . The Frobenius integrability of D_{\pm} makes (M, g, Ω) not only almost para–Kähler but also para–Kähler. Further discussion about this distinction can be found in Chapter 6.

Chapter 4

An Einstein metric on an \mathbb{R}^* bundle over M

In this chapter, we show that there is a canonical Einstein metric on an \mathbb{R}^* bundle over M , with a connection whose curvature is the pull-back of the symplectic structure from M . This metric is interesting in the context of Kaluza–Klein theory. The material covered here is based on material appearing in [30].

4.1 The model case

We first note that the flat projective structure on \mathbb{RP}^n gives a one-parameter family of $2n$ -dimensional Einstein spaces M which are so-called Kaluza–Klein reductions of quadrics in \mathbb{R}^{2n+2} endowed with a flat neutral signature metric. When \mathbb{R}^{2n+2} carries such a metric, we call it $\mathbb{R}^{n+1, n+1}$. Note that the phrase Kaluza–Klein refers to a classical unified field theory of gravitation and electromagnetism. We will use it to mean the construction of g from a metric in dimension $2n + 1$ of the form

$$g \pm f^2(A + dt)^2,$$

where g , A and f are a metric, Maxwell potential and function on M .

For $N = \mathbb{RP}^n$, the family $\{(g_\Lambda, \Omega_\Lambda) ; \Lambda \neq 0\}$ is given by

$$\begin{aligned} g_\Lambda &= dx^i \odot d\zeta_i + \Lambda \zeta_i \zeta_j dx^i \odot dx^j \\ \Omega_\Lambda &= d\zeta_i \wedge dx^i. \end{aligned} \tag{4.1}$$

Proposition 4.1.1. *The Einstein spaces M carrying metric and symplectic form (4.1) are projections from the $(2n+1)$ -dimensional quadrics $\mathcal{Q} \subset \mathbb{R}^{n+1, n+1}$ given by $P^\alpha L_\alpha = \frac{1}{\Lambda}$, where $P, L \in \mathbb{R}^{n+1}$ are coordinates on $\mathbb{R}^{n+1, n+1}$ such that the metric is given by*

$$\hat{g} = dP^\alpha dL_\alpha,$$

under the embedding

$$\begin{aligned} P^\alpha &= \begin{cases} x^i e^t, & \alpha = i = 1, \dots, n \\ e^t, & \alpha = n+1 \end{cases} \\ L_\alpha &= \begin{cases} \zeta_i e^{-t}, & \alpha = i = 1, \dots, n \\ e^{-t} \left(\frac{1}{\Lambda} - x^k \zeta_k \right), & \alpha = n+1 \end{cases} \end{aligned} \quad (4.2)$$

following Kaluza–Klein reduction by the vector $\frac{\partial}{\partial t}$.

Proof. We find the basis of coordinate 1-forms $\{dP^\alpha, dL_\alpha\}$ to be

$$\begin{aligned} dP^\alpha &= \begin{cases} e^t(dx^i + x^i dt), & \alpha = i = 1, \dots, n \\ e^t dt, & \alpha = n+1 \end{cases} \\ dL_\alpha &= \begin{cases} e^{-t}(d\zeta_i - \zeta_i dt), & \alpha = i = 1, \dots, n \\ -e^{-t} \left[\left(\frac{1}{\Lambda} - x^k \zeta_k \right) dt + x^k d\zeta_k + \zeta_k dx^k \right], & \alpha = n+1. \end{cases} \end{aligned}$$

The metric is then given by

$$\begin{aligned} \hat{g} &= e^t(dx^i + x^i dt) \odot e^{-t}(d\zeta_i - \zeta_i dt) - e^t dt \odot e^{-t} \left[\left(\frac{1}{\Lambda} - x^k \zeta_k \right) dt + x^k d\zeta_k + \zeta_k dx^k \right] \\ &= dx^i \odot d\zeta_i + (x^i d\zeta_i - \zeta_i dx^i) \odot dt - (x^i \zeta_i) dt^2 \\ &\quad - dt \odot \left[\left(\frac{1}{\Lambda} - x^k \zeta_k \right) dt + x^k d\zeta_k + \zeta_k dx^k \right] \\ &= dx^i \odot d\zeta_i - \frac{1}{\Lambda} dt^2 - 2\zeta_i dx^i \odot dt \\ &= dx^i \odot d\zeta_i + \Lambda \zeta_i \zeta_j dx^i \odot dx^j - \Lambda \left(\frac{dt}{\Lambda} + \zeta_i dx^i \right)^2, \end{aligned}$$

which is clearly going to give g_Λ under Kaluza–Klein reduction by $\frac{\partial}{\partial t}$.

□

Note that the symplectic form Ω is the exterior derivative of the potential term $\zeta_i dx^i$, implying a possible generalisation to the curved case.

4.2 The general case

We now return to a general projective structure $(N, [\nabla])$, corresponding to the metric and symplectic form (3.21,3.22). Since symplectic form picks out the antisymmetric part of the Schouten tensor, it has the fairly simple form

$$\Omega_\Lambda = d\zeta_i \wedge dx^i - \frac{\partial_{[i} \Gamma_{j]k}^k}{\Lambda(n+1)} dx^i \wedge dx^j.$$

By inspection, this can be written $\Omega_\Lambda = d\mathcal{A}$, where

$$\mathcal{A} = \zeta_i dx^i - \frac{\Gamma_{ik}^k}{\Lambda(n+1)} dx^i.$$

This is a trivialisation of the Kaluza–Klein bundle which we are about to construct. Note that for $\Lambda = 1$, under a change of projective connection (3.1) the corresponding change in the fibre coordinates (3.20) ensures that Ω and \mathcal{A} are unchanged.

Motivated by the Kaluza–Klein reduction in the flat case, we consider the following metric.

Theorem 4.2.1. *The metric*

$$\hat{g}_\Lambda = g_\Lambda - \Lambda \left(\frac{dt}{\Lambda} + \mathcal{A} \right)^2 \quad (4.3)$$

on an \mathbb{R}^* bundle $\kappa_Q : Q \rightarrow M$ is Einstein, with Ricci scalar $2n(2n+1)\Lambda$.

Proof. We prove this using the Cartan formalism. Our treatment parallels a calculation by Kobayashi [42], who considered principal circle bundles over Kähler manifolds in order to study the topology of the base. For the remainder of this chapter we will suppress the constant Λ , writing $\hat{g} \equiv \hat{g}_\Lambda$, $g \equiv g_\Lambda$ and $\Omega \equiv \Omega_\Lambda$ since the proof applies to any choice $\Lambda \neq 0$ within this family.

Consider a frame

$$e^a = \begin{cases} dx^i, & a = i = 1, \dots, n \\ d\zeta_i - (\Gamma_{ij}^k \zeta_k - \Lambda \zeta_i \zeta_j - \Lambda^{-1} P_{ij}) dx^j, & a = i + n = n + 1, \dots, 2n. \end{cases} \quad (4.4)$$

In this basis the metric takes the form

$$g = e^1 \odot e^{n+1} + \cdots + e^n \odot e^{2n}. \quad (4.5)$$

We are interested in the metric

$$\hat{g} = -e^0 \odot e^0 + g, \quad (4.6)$$

where

$$e^0 = \sqrt{\Lambda} \left(\frac{dt}{\Lambda} + \mathcal{A} \right).$$

Note that for $\Lambda < 0$ one can instead take the absolute value of Λ inside the square root, and change the sign of $e^0 \odot e^0$ in the definition of \hat{g} .

We reserve Latin indices a, b, \dots for the $2n$ -metric components $1, \dots, 2n$ and allow Greek indices μ, ν, \dots to take values $0, 1, \dots, 2n$. The dual basis to $\{e^\mu\}$ will be denoted $\{E_\mu\}$ and will act on functions as vector fields in the usual way. We wish to find the new connection 1-forms $\hat{\psi}^\mu_\nu$ (defined by $de^\mu = -\hat{\psi}^\mu_\nu \wedge e^\nu$) in terms of the old ones ψ^a_b (defined by $de^a = -\psi^a_b \wedge e^b$). Hence we examine de^0 to find $\hat{\psi}^0_a$.

$$\begin{aligned} de^0 &= \sqrt{\Lambda} d\mathcal{A} = \sqrt{\Lambda} \Omega_{ab} e^a \wedge e^b = -\hat{\psi}^0_a \wedge e^a \implies \\ \hat{\psi}^0_a &= \sqrt{\Lambda} \Omega_{[ab]} e^b = \sqrt{\Lambda} \Omega_{ab} e^b, \quad \hat{\psi}^a_0 = \sqrt{\Lambda} \Omega^a_b e^b. \end{aligned}$$

Since de^a is unchanged, we have that

$$\hat{\psi}^a_0 \wedge e^0 + \hat{\psi}^a_b \wedge e^b = \psi^a_b \wedge e^b,$$

thus

$$\hat{\psi}^a_b \wedge e^b = \psi^a_b \wedge e^b - \sqrt{\Lambda} \Omega^a_b e^b \wedge e^0 \implies \hat{\psi}^a_b = \psi^a_b + \sqrt{\Lambda} \Omega^a_b e^0.$$

We now calculate the curvature 2-forms $\hat{\Psi}^\mu_\nu = d\hat{\psi}^\mu_\nu + \hat{\psi}^\mu_\rho \wedge \hat{\psi}^\rho_\nu = \frac{1}{2} \hat{R}_{\rho\sigma\nu}{}^\mu e^\rho \wedge e^\sigma$ in terms of $\Psi^a_b = d\psi^a_b + \psi^a_c \wedge \psi^c_b$, where $\hat{R}_{\rho\sigma\nu}{}^\mu$ is the Riemann tensor of \mathcal{Q} . Note that

we use the notation $\psi^a_b = \psi^a_{bc}e^c$.

$$\begin{aligned}
\hat{\Psi}^a_b &= d\hat{\psi}^a_b + \hat{\psi}^a_c \wedge \hat{\psi}^c_b + \hat{\psi}^a_0 \wedge \hat{\psi}^0_b \\
&= d\psi^a_b + \sqrt{\Lambda}d(\Omega^a_b e^0) + \psi^a_c \wedge \psi^c_b + \sqrt{\Lambda}\Omega^a_c e^0 \wedge \psi^c_b \\
&\quad + \sqrt{\Lambda}\Omega^c_b \psi^a_c \wedge e^0 + \Lambda\Omega^a_{[c}\Omega_{|b|d]} e^c \wedge e^d \\
&= \Psi^a_b + \sqrt{\Lambda}E_c(\Omega^a_b)e^c \wedge e^0 + \Lambda(\Omega^a_b\Omega_{cd} + \Omega^a_{[c}\Omega_{|b|d]})e^c \wedge e^d \\
&\quad + \sqrt{\Lambda}(\Omega^c_b \psi^a_{cd} - \Omega^a_c \psi^c_{bd})e^d \wedge e^0 \\
&= \Psi^a_b + \sqrt{\Lambda}^g \nabla_c \Omega^a_b e^c \wedge e^0 + \Lambda(\Omega^a_b\Omega_{cd} + \Omega^a_{[c}\Omega_{|b|d]})e^c \wedge e^d. \\
\hat{\Psi}^0_a &= d\hat{\psi}^0_a + \hat{\psi}^0_b \wedge \hat{\psi}^b_a \\
&= \sqrt{\Lambda}E_{[c}(\Omega_{|a|b]})e^c \wedge e^b - \sqrt{\Lambda}\Omega_{ab}\psi^b_c \wedge e^c + \sqrt{\Lambda}\Omega_{bc}e^c \wedge (\psi^b_a + \sqrt{\Lambda}\Omega^b_a e^0) \\
&= \sqrt{\Lambda}(E_{[d}(\Omega_{|a|b]}) - \Omega_{ac}\psi^c_{[bd]} + \Omega_{c[d}\psi^c_{|a|b]})e^d \wedge e^b + \Lambda\Omega_{bc}\Omega^b_a e^c \wedge e^0 \\
&= \sqrt{\Lambda}^g \nabla_{[c}\Omega_{|a|d]}e^c \wedge e^d + \Lambda\Omega_{bc}\Omega^b_a e^c \wedge e^0.
\end{aligned}$$

Hence we have that

$$\begin{aligned}
\hat{R}_{cdb}^a &= R_{cdb}^a + 2\Lambda(\Omega^a_b\Omega_{cd} + \Omega^a_{[c}\Omega_{|b|d]}) \\
\hat{R}_{c0b}^a &= \sqrt{\Lambda}^g \nabla_c \Omega^a_b \\
\hat{R}_{cda}^0 &= 2\sqrt{\Lambda}^g \nabla_{[c}\Omega_{|a|d]} \\
\hat{R}_{c0a}^0 &= \Lambda\Omega_{bc}\Omega^b_a,
\end{aligned}$$

and thus, using $\hat{R}_{\mu\nu} = \hat{R}_{\rho\mu\nu}{}^\rho$,

$$\begin{aligned}
\hat{R}_{00} &= \Lambda\Omega_{bc}\Omega^{bc} = -2n\Lambda = 2n\Lambda\hat{g}_{00} \\
\hat{R}_{b0} &= \sqrt{\Lambda}^g \nabla_c \Omega^c_b = 0 \\
\hat{R}_{db} &= R_{db} + 2\Lambda(\Omega^c_b\Omega_{cd} + \frac{1}{2}\Omega^c_c\Omega_{bd} - \frac{1}{2}\Omega^c_d\Omega_{bc}) - \Lambda\Omega_{cd}\Omega^c_b \\
&= R_{db} + 2\Lambda\Omega^c_b\Omega_{dc} \\
&= 2(n+1)\Lambda g_{db} - 2\Lambda g_{db} = 2n\Lambda g_{db} = 2n\Lambda\hat{g}_{db}.
\end{aligned}$$

Note that we have used the facts that g is Einstein with Ricci scalar $4n(n+1)\Lambda$ and the symplectic form Ω is divergence-free; these are justified in Lemmas 4.2.3 and 4.2.2 below. Since $\hat{g}_{a0} = 0$, we conclude that

$$\hat{R}_{\mu\nu} = 2n\Lambda\hat{g}_{\mu\nu} = \frac{\hat{R}}{2n+1}\hat{g}_{\mu\nu},$$

i.e. \hat{g} is Einstein with Ricci scalar $2n(2n+1)\Lambda$.

□

Physically, this is a Kaluza–Klein reduction with constant dilation field and where the Maxwell two-form is related to the reduced metric by $\Omega_a^c \Omega_{cb} = g_{ab}$. This is what allows both the reduced and lifted metric to be Einstein. A more general discussion can be found in [64].

From the Cartan perspective, $\hat{g}_{\Lambda=1}$ can be thought of as a metric on the $(2n+1)$ -dimensional space obtained by taking a quotient $\hat{\kappa}_q : \mathcal{G} \rightarrow \mathcal{G}/SL(n, \mathbb{R}) = \mathcal{Q}$ of the Cartan bundle, where we embed $SL(n, \mathbb{R}) \subset GL(n, \mathbb{R})$ in S as in (3.16) but with a now denoting an element of $SL(n, \mathbb{R})$ (so that $\det a^{-1} = 1$). This new subgroup acts adjointly on θ as

$$\begin{pmatrix} 1 & 0 \\ 0 & a \end{pmatrix} \begin{pmatrix} -\text{tr}\phi & \eta \\ \omega & \phi \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & a^{-1} \end{pmatrix} = \begin{pmatrix} -\text{tr}\phi & \eta a^{-1} \\ a\omega & \phi \end{pmatrix},$$

so not only is the inner product $\eta\omega$ invariant but also the $(0,0)$ -component $\theta^0_0 = -\text{tr}\phi$, which is a scalar one-form whose exterior derivative is constrained by (3.6) to be $d\theta^0_0 = -\theta^0_i \wedge \theta^i_0 = -\text{Ant}(\eta \wedge \omega)$. Thus, denoting by \mathcal{A} the object on $\mathcal{Q} = \mathcal{G}/SL(n, \mathbb{R})$ which is such that $\hat{\kappa}_q^* \mathcal{A} = \text{tr}\phi$, we have that $d\mathcal{A} = \Omega$ (where we are now taking Ω to be defined on \mathcal{Q} by $\hat{\kappa}_q^* \Omega = \text{Ant}(\eta \wedge \omega)$).

We then have a natural way of constructing a metric \hat{g} on \mathcal{Q} as a linear combination of g and $e^0 \odot e^0$, where e^0 is \mathcal{A} up to addition of some exact one-form. It turns out that there is choice (4.6) of linear combination with $e^0 = \mathcal{A}$ such that \hat{g} is Einstein. The fact that this metric is exactly (4.3) can be verified by constructing the Cartan connection of $(N, [\nabla])$ explicitly in terms of a representative connection $\nabla \in [\nabla]$.

4.2.1 Ricci scalar of g_Λ and divergence of Ω_Λ

Lemma 4.2.2. *The symplectic form $\Omega \equiv \Omega_\Lambda$ on M is divergence-free. In index notation,*

$${}^g \nabla^c \Omega_{cb} = 0.$$

Proof. In the basis (4.4) we have g as above (4.5) and

$$\Omega = \sum_{i=1}^n e^i \wedge e^{i+n} \implies \Omega_{ab} = \sum_{i=1}^n \delta_{[a}^i \delta_{b]}^{i+n}.$$

Note that from now on we will omit the summation sign and use the summation convention regardless of whether i, j indices are up or down. As in Section 4.1, we look for $\psi^a{}_b$ by considering de^a (recall that $i, j = 1, \dots, n$ and $a, b = 1, \dots, 2n$):

$$\begin{aligned}
de^i &= 0 \\
de^{i+n} &= -(E_l(\Gamma_{ij}^k)\zeta_k - \Lambda^{-1}E_l(P_{ij}))dx^l \wedge dx^j - (\Gamma_{ij}^k - 2\Lambda\zeta_{(i}\delta_{j)}^k)d\zeta_k \wedge dx^j \\
&= -(E_l(\Gamma_{ij}^k)\zeta_k - \Lambda^{-1}E_l(P_{ij}))e^l \wedge e^j \\
&\quad - (\Gamma_{ij}^k - 2\Lambda\zeta_{(i}\delta_{j)}^k)(e^{k+n} + (\Gamma_{km}^l\zeta_l - \Lambda\zeta_k\zeta_m - \Lambda^{-1}P_{km})e^m) \wedge e^j \\
&= [\Lambda^{-1}E_m(P_{ij}) - E_m(\Gamma_{ij}^k)\zeta_k + \Lambda^{-1}\Gamma_{ij}^kP_{km} - \Gamma_{ij}^k\Gamma_{km}^l\zeta_l + \Lambda\Gamma_{ij}^k\zeta_m\zeta_k \\
&\quad + 2\Lambda\zeta_{(i}(\Gamma_{j)m}^l\zeta_l - \Lambda\zeta_j)\zeta_m - \Lambda^{-1}P_{j)m}]e^m \wedge e^j + (2\Lambda\zeta_{(i}\delta_{j)}^k - \Gamma_{ij}^k)e^{k+n} \wedge e^j \\
&= [\Lambda^{-1}\nabla_m P_{ij} - (\nabla_m \Gamma_{ij}^k)\zeta_k - 2\zeta_{(i}P_{j)m}]e^m \wedge e^j + (2\Lambda\zeta_{(i}\delta_{j)}^k - \Gamma_{ij}^k)e^{k+n} \wedge e^j.
\end{aligned}$$

Note that we have used ∇ to denote the chosen connection on N with components Γ_{jk}^i ; it is a different object to the Levi–Civita connection ${}^g\nabla$ on M . Next we wish to read off the spin connection $\psi^a{}_b$ such that $de^a = -\psi^a{}_b \wedge e^b$ and the following index symmetries are satisfied:

$$\begin{aligned}
\psi^i{}_j &= \frac{1}{2}\psi_{i+n}{}j = -\frac{1}{2}\psi_{j+n}{}i = -\psi^{j+n}{}_{i+n}, \\
\psi^i{}_{j+n} &= \frac{1}{2}\psi_{i+n}{}_{j+n} = -\frac{1}{2}\psi_{j+n}{}_{i+n} = -\psi^j{}_{i+n}, \\
\psi^{i+n}{}_j &= \frac{1}{2}\psi_{i}{}_{j+n} = -\frac{1}{2}\psi_{j}{}_{i+n} = -\psi^{j+n}{}_i.
\end{aligned}$$

We find that

$$\begin{aligned}
\psi^{i+n}{}_{k+n} &= (2\Lambda\zeta_{(i}\delta_{j)}^k - \Gamma_{ij}^k)e^j = -\psi^k{}_i \\
\psi^{i+n}{}_j &= [2(\nabla_{[i}\Gamma_{j]k}^l)\zeta_l - 2\Lambda^{-1}\nabla_{[i}P_{j]k}^S - \Lambda^{-1}\nabla_k P_{ij}^A + 2\zeta_{(j}P_{k)i} - 2\zeta_{(i}P_{k)j}]e^k =: A_{ijk}e^k \\
\psi^i{}_{j+n} &= 0.
\end{aligned}$$

One can check that these satisfy both the index symmetries above and are such that $de^a = -\psi^a{}_b \wedge e^b$, and we know from theory that there is a unique set of $\psi^a{}_b$ that have both of these properties. Note that we have used P^S and P^A to denote the symmetric and antisymmetric parts of P in order to avoid too much confusion from having multiple symmetrisation brackets in the indices.

We are now ready to calculate the divergence of Ω . Since it is covariantly constant in this basis, we obtain

$$g\nabla_c \Omega_{ab} = -\psi_{ac}^d \Omega_{db} - \psi_{bc}^d \Omega_{ad} = -\psi_{ac}^d \Omega_{db} + \psi_{bc}^d \Omega_{da} = 2\Omega_{d[a} \psi_{b]c}^d.$$

We can split the right hand side into

$$\begin{aligned} \Omega_{da} \psi_{bc}^d &= \Omega_{ka} \psi_{bc}^k + \Omega_{k+n a} \psi_{bc}^{k+n} \\ &= \delta_{[k}^i \delta_{a]}^{i+n} \psi_{bc}^k + \delta_{[k+n}^i \delta_{a]}^{i+n} \psi_{bc}^{k+n} \\ &= \frac{1}{2} \left(-\delta_a^{k+n} \delta_b^i \delta_c^j (2\Lambda \zeta_{(i} \delta_{j)}^k - \Gamma_{ij}^k) - \delta_a^k \delta_b^{l+n} \delta_c^j (2\Lambda \zeta_{(k} \delta_{j)}^l - \Gamma_{kj}^l) - \delta_a^k \delta_b^l \delta_c^m A_{klm} \right). \end{aligned}$$

The first two terms are the same but with $a \leftrightarrow b$, so are lost in the antisymmetrisation. Thus

$$g\nabla_c \Omega_{ab} = -\delta_{[a}^k \delta_{b]}^l \delta_c^m A_{klm}.$$

Tracing amounts to contracting this with g^{ac} :

$$g\nabla^c \Omega_{cb} = -\delta_{[a}^k \delta_{b]}^l g^{ac} \delta_c^m A_{klm} = -\delta_{[a}^k \delta_{b]}^l g^{am} A_{klm},$$

but g^{am} is non-zero only when $a = m + n > n$ and $\delta_{[a}^k \delta_{b]}^l$ is non-zero only when $a = k \leq n$ or $a = l \leq n$. We can therefore conclude that the right hand side is zero and Ω is divergence-free.

□

Lemma 4.2.3. *The metric $g \equiv g_\Lambda$ corresponding to a projective structure $(N, [\nabla])$ in dimension n has Ricci scalar*

$$R = 4n(n+1)\Lambda.$$

Proof. We calculate the Ricci scalar of g (given that it's Einstein, as stated in the appendix of [28]) via the curvature two-forms $\Psi^a_b = d\psi^a_b + \psi^a_c \wedge \psi^c_b = \frac{1}{2} R_{cd}{}^a e^c \wedge e^d$. We are only interested in non-zero components of the Ricci tensor such as $R_{i j+n} = R_{c i j+n}^c$. In fact, we will calculate only $R_{m+n j}$, for which we need to consider $R_{l m+n j}{}^i$ and $R_{k+n m+n j}{}^{l+n}$, i.e. we need only calculate Ψ^i_j and $\Psi^{l+n j}$.

$$\Psi^i_j = d \left((\Gamma_{jk}^i - 2\Lambda \zeta_{(j} \delta_{k)}^i) e^k \right) + \psi^i_k \wedge \psi^k_j + \psi^i_{k+n} \wedge \psi^{k+n j}.$$

The last term vanishes since $\psi^{k+n}{}_j = 0$, and the middle term only has components that look like $\frac{1}{2}R_{lmj}{}^i e^l \wedge e^m$, so the only term we are interested in is

$$-2\Lambda d\zeta_{(j}\delta_k^i e^k = -2\Lambda\delta_{(k}^i(e^{j)+n} + (\Gamma_{j)m}^l\zeta_l - \Lambda\zeta_{j)}\zeta_m - \Lambda^{-1}\mathbf{P}_{j)m})e^m) \wedge e^k.$$

Again, discarding the $e^m \wedge e^k$ term gives

$$-\Lambda(\delta_k^i e^{j+n} \wedge e^k + \delta_j^i e^{k+n} \wedge e^k) = \frac{1}{2}R_{lm+nj}{}^i e^l \wedge e^{m+n} + \frac{1}{2}R_{m+n+lj}{}^i e^{m+n} \wedge e^l,$$

so we conclude

$$R_{lm+nj}{}^i = \Lambda(\delta_j^i\delta_l^m + \delta_l^i\delta_j^m).$$

The other Riemann tensor component we need to know to calculate $R_{m+nj} = R_{cm+nj}{}^c$ is $R_{l+n m+n j}{}^{i+n}$, so we examine

$$\Psi^{i+n}{}_j = d\psi^{i+n}{}_j + \psi^{i+n}{}_k \wedge \psi^k{}_j + \psi^{i+n}{}_{k+n} \wedge \psi^{k+n}{}_j,$$

but none of these terms have $e^{l+n} \wedge e^{m+n}$ components, so $R_{l+n m+n j}{}^{i+n} = 0$. Hence

$$R_{m+nj} = \delta_l^i R_{l m+nj}{}^i = \Lambda(\delta_j^m + n\delta_l^m) = \Lambda(n+1)\delta_j^m.$$

Setting this equal to $\frac{R}{2n}g_{m+nj} = \frac{R}{4n}\delta_j^m$ we find

$$R = 4n(n+1)\Lambda,$$

as required. □

Chapter 5

Einstein–Weyl structures and $SU(\infty)$ –Toda fields

In this chapter, we focus on the four dimensional Einstein manifolds that arise from the projective to Einstein correspondence in the case $n = 2$. As discussed in Chapter 3, it is shown in [28] that the Einstein manifolds in this subclass have ASD Weyl tensor, and are therefore associated with a twistor space [62]. If they also carry a Killing vector field arising from a symmetry of the underlying projective surface, one can extract solutions of the $SU(\infty)$ –Toda equation via symmetry reduction to Lorentzian Einstein–Weyl structures in $2 + 1$ dimensions [40, 74].

The aim of this chapter is to investigate the Einstein–Weyl structures obtainable in this way, resulting in several examples of new, explicit solutions of the Toda equation. We also give an explicit criterion for a vector field that generates a symmetry of a Weyl structure, and prove some results about the Einstein manifold and corresponding twistor space arising from the flat projective surface \mathbb{RP}^2 . The content of this chapter is based on some of the work in [30] and was done in collaboration with Maciej Dunajski.

In the case $n = 2$ we will write the metric and symplectic form as

$$g = dz_{A'} \odot dx^{A'} - (\Gamma_{A'B'}^{C'} z_{C'} - z_{A'} z_{B'} - P_{A'B'}) dx^{A'} \odot dx^{B'}, \quad (5.1)$$

$$\Omega = dz_{A'} \wedge dx^{A'} + P_{A'B'} dx^{A'} \wedge dx^{B'}, \quad A', B', C' = 0, 1. \quad (5.2)$$

where we have replaced $\{\zeta_i\}$ with $\{z_{A'}\}$ and shifted the indices from $i, j = 1, 2$ to $A', B' = 0, 1$. This is helpful for the twistorial calculations because it agrees with the usual notation for two–component spinor indices. Note that a change of projective

connection is now given by

$$\Gamma_{A'B'}^{C'} \rightarrow \Gamma_{A'B'}^{C'} + \delta_{A'}^{C'} \Upsilon_{B'} + \delta_{B'}^{C'} \Upsilon_{A'}, \quad z_{A'} \rightarrow z_{A'} + \Upsilon_{A'}, \quad A', B', C' = 0, 1. \quad (5.3)$$

5.1 Background

5.1.1 Anti–self–duality, spinors and totally null distributions

Let M be an oriented four dimensional manifold with a metric g of signature $(2, 2)$. The Hodge operator $*$ on the space of two forms is an involution, and induces a decomposition [2]

$$\Lambda^2(T^*M) = \Lambda_-^2(T^*M) \oplus \Lambda_+^2(T^*M) \quad (5.4)$$

of two-forms into ASD and SD components, which only depends on the conformal class of g . The Riemann tensor of g can be thought of as a map $\mathcal{R} : \Lambda^2(T^*M) \rightarrow \Lambda^2(T^*M)$ which admits a decomposition under (5.4):

$$\mathcal{R} = \left(\begin{array}{c|c} C_+ + \frac{R}{12} & \phi \\ \hline - & - \\ \phi & C_- + \frac{R}{12} \end{array} \right). \quad (5.5)$$

Here C_{\pm} are the SD and ASD parts of the Weyl conformal curvature tensor¹, ϕ is the trace-free Ricci curvature, and R is the scalar curvature which acts by scalar multiplication. The metric g is Einstein if $\phi = 0$, and the corresponding conformal structure $[g]$ is ASD if $C_+ = 0$. We will call g *conformally ASD* if it has ASD Weyl tensor. If both of these conditions are satisfied then the Riemann tensor is also ASD.

Two–component spinors

The symmetry group of a metric in signature $(2, 2)$ decomposes under the Lie group isomorphism

$$SO(2, 2) \cong SL(2, \mathbb{R}) \times SL(2, \mathbb{R})/\mathbb{Z}_2.$$

¹Note this is a different object to the projective Weyl tensor discussed in Chapter 3.

Locally there exist real rank two vector bundles \mathbb{S}, \mathbb{S}' (spin bundles) over M such that [63]

$$TM \cong \mathbb{S} \otimes \mathbb{S}' \quad (5.6)$$

is a canonical bundle isomorphism. We will use the usual notation $\iota^A \in \Gamma(\mathbb{S})$, $\pi^{A'} \in \Gamma(\mathbb{S}')$, where $A, A' = 0, 1$. There are unique skew-symmetric sections $\varepsilon \in \Gamma(\mathbb{S} \otimes \mathbb{S})$ and $\varepsilon' \in \Gamma(\mathbb{S}' \otimes \mathbb{S}')$, and one can argue that

$$g_{ab} \xi^a \tilde{\xi}^b = \varepsilon_{AB} \varepsilon_{A'B'} \xi^{AA'} \tilde{\xi}^{BB'} \quad (5.7)$$

for vector fields $\xi, \tilde{\xi}$ on M . The bundles \mathbb{S} and \mathbb{S}' inherit connections from $\mathbf{g}\nabla$ for which $\varepsilon, \varepsilon'$ are parallel.

We identify \mathbb{S} with its dual according to

$$\iota_A = \iota^B \varepsilon_{BA}, \quad \iota^A = \varepsilon^{AB} \iota_B,$$

and similarly for \mathbb{S}' . Note that the contraction is always over adjacent indices descending to the right, and $\varepsilon^{AB} \varepsilon_{CB} = \delta_C^A$. In higher valence spinors, the relative order of primed and unprimed indices is unimportant.

A vector $\xi \in \Gamma(TM)$ is called null if $g(\xi, \xi) = 0$. For $\tilde{\xi} = \xi$ the right hand side of (5.7) is just the determinant of $\xi^{AA'}$ viewed as a matrix, so any null vector is of the form $\xi = \iota \otimes \pi$ where ι and π are sections of \mathbb{S} and \mathbb{S}' respectively. The antisymmetry of ε means that $\varepsilon(\iota, \iota) = 0$ for any section $\iota \in \Gamma(\mathbb{S})$ (and similarly for any $\pi \in \mathbb{S}'$), so the converse is also true (i.e. any vector that can be written $V = \iota \otimes \pi$ is null).

Since the symplectic structures $\varepsilon_{AB}, \varepsilon_{A'B'}$ are the unique skew-symmetric two-index spinors up to scale, any spinor of valence n which is skew on a pair of indices can be factorised as the tensor product of a spinor of valence $n - 2$ and either ε or ε' . This leads to the decomposition of a two-form $F_{ab} = F_{AA'BB'} = F_{ABA'B'}$ as

$$F_{ABA'B'} = \varepsilon_{AB} \Phi_{A'B'} + \varepsilon_{A'B'} \Psi_{AB},$$

where $\Phi_{A'B'} = \Phi_{(A'B')}$ and $\Psi_{AB} = \Psi_{(AB)}$. One can show using an analogous decomposition of the volume form that $\Phi_{A'B'}$ and Ψ_{AB} are the SD and ASD parts of F_{ab} respectively.

The nonlinear graviton

Any two dimensional distribution on a four–manifold M can be expressed as the kernel of a two–form, and we define a distribution to be (A)SD if the corresponding two–form is (A)SD. Taking any $\iota^A \in \Gamma(\mathbb{S})$, the two–form $\iota_A \iota_B \varepsilon_{A'B'}$ defines an ASD distribution $D_\beta = \{\iota^A w^{A'}, w^{A'} \in \Gamma(\mathbb{S}')\}$ which is totally null in the sense that $g(\xi, \tilde{\xi}) = 0$ for all $\xi, \tilde{\xi} \in \Gamma(D)$. We call this a β –distribution. Note that it is only defined up to scale, which means that there is an \mathbb{RP}^1 worth of β –planes at every point in M . Given any $\pi^{A'} \in \Gamma(\mathbb{S}')$, one can similarly define a totally null SD distribution called an α –distribution by

$$D_\alpha = \{\iota^A \pi^{A'}, \iota^A \in \Gamma(\mathbb{S})\} = \text{span}\{\pi^{A'} \mathbf{e}_{AA'}\}, \quad (5.8)$$

where $\{\mathbf{e}_{AA'}\}$ is a null tetrad of vector fields.

An α –surface (respectively β –surface) is a two dimensional surface in M which is tangent to an α –(β –)distribution at every point. A foliation by α –surfaces exists if and only if the corresponding distribution (5.8) is Frobenius integrable. Penrose’s nonlinear graviton Theorem [62] states that a maximal, three dimensional family of α –surfaces exists in M if and only if its conformal curvature is ASD, i.e. $C_+ = 0$. Existence of such a family is equivalent to the distribution (5.8) being integrable for any $\pi^{A'}$. We can state this condition as integrability of the lift of the distribution (5.8) to \mathbb{S}' . This is given by

$$\mathcal{D} = \text{span}\{L_A := \pi^{A'} \tilde{\mathbf{e}}_{AA'}\}, \quad (5.9)$$

where the vectors

$$\tilde{\mathbf{e}}_{AA'} = \mathbf{e}_{AA'} - \Gamma_{AA'B'}^{C'} \pi^{B'} \frac{\partial}{\partial \pi^{C'}}$$

are the lifts of the null tetrad $\{\mathbf{e}_{AA'}\}$ to \mathbb{S}' , and $\Gamma_{AA'B'}^{C'}$ are the components of the *spin connection* on \mathbb{S}' , which is inherited from the Levi–Civita connection of g on TM . We call \mathcal{D} the *twistor distribution*.

In fact, Penrose considers four dimensional *complex* manifolds² M carrying a metric which is *holomorphic* in the sense that the metric components depend on the coordinates on M and not on their complex conjugates. Then \mathbb{S}, \mathbb{S}' are complex vector bundles over M and $\varepsilon, \varepsilon'$ are holomorphic symplectic forms. A real conformally ASD metric in a given signature can then be obtained by choosing the correct reality conditions. In neutral signature, complex conjugation is a map from \mathbb{S} to itself (or from \mathbb{S}' to itself)

²Note that familiar facts from real geometry such as a unique Levi–Civita connection and the Frobenius theorem carry over to holomorphic geometry. See [47] for details.

which simply replaces each component of a spinor with its complex conjugate. Thus the reality conditions in neutral signature amount to identifying spinors with their complex conjugates.

The *twistor space* \mathcal{T} of M is then defined as the three dimensional complex manifold comprising the set of all α -surfaces in M . Each point $m \in M$ corresponds to a subset $\mathcal{L}_m \subset \mathcal{T}$ of α -surfaces which pass through m . Since an α -surface at m is defined by a $\pi^{A'} \in \mathbb{S}'|_m$ up to scale, \mathcal{L}_m is an embedding $\mathbb{CP}^1 \subset \mathcal{T}$. The *correspondence space* $\mathcal{F} = M \times \mathbb{CP}^1$ has local coordinates $(x^a, \lambda) := (x^a, \pi_{0'} / \pi_{1'})$, where $\pi^{A'}$ parametrises the set of α -surfaces through the point in M with coordinates x^a . Note that \mathcal{F} can be obtained from the primed spin bundle $\mathbb{S}' \rightarrow M$ by projectivising each fibre, and carries a distribution $\tilde{\mathcal{D}} = \text{span}\{\tilde{L}_A\}$ given by the push forward of the twistor distribution \mathcal{D} to $\mathcal{F} = \mathbb{P}(\mathbb{S}')$. We also call $\tilde{\mathcal{D}}$ the twistor distribution. Note that \mathcal{F} has the alternative definition $\mathcal{F} = \{(Z, m) \in \mathcal{T} \times M : Z \in \mathcal{L}_m\}$, leading to the double fibration

$$M \leftarrow \mathcal{F} \rightarrow \mathcal{T},$$

where the map $\mathcal{F} \rightarrow \mathcal{T}$ is the quotient of \mathcal{F} by the leaves of the distribution $\tilde{\mathcal{D}}$. A twistor function is a function on \mathcal{F} which is constant along $\tilde{\mathcal{D}}$.

The nonlinear graviton allows us to express an ASD conformal structure in terms of the algebraic geometry of \mathcal{T} . First note that if two points $m_1, m_2 \in M$ are null-separated, then the corresponding curves $\mathcal{L}_{m_1}, \mathcal{L}_{m_2}$ intersect at a single point. This is because any null geodesic must have a tangent vector field of the form $\iota^A \pi^{A'}$ for some sections $\iota^A \in \Gamma(\mathbb{S})$ and $\pi^{A'} \in \Gamma(\mathbb{S}')$, and thus the geodesic is contained within the unique α -surface spanned by $\pi^{A'} e_{AA'}$. This unique α -surface corresponds to the point in \mathcal{T} where the curves $\mathcal{L}_{m_1}, \mathcal{L}_{m_2}$ meet.

In order to understand this correspondence at an infinitesimal level and thereby explicitly recover an ASD conformal structure from \mathcal{T} , we need to understand the *normal bundle* $\mathbb{N}(\mathcal{L}_m) := \cup_{Z \in \mathcal{L}_m} \{T_Z \mathcal{T} / T_Z \mathcal{L}_m\}$ over a \mathbb{CP}^1 embedding \mathcal{L}_m . This is evidently a complex vector bundle, and in fact it is a *holomorphic* vector bundle, meaning that the total space is a complex manifold and the projection $\mathbb{N}(\mathcal{L}_m) \rightarrow \mathcal{L}_m$ is holomorphic. It is thus subject to the following theorem due to Birkhoff and Grothendieck (see for example [58] for a proof).

Theorem 5.1.1 (Birkhoff–Grothendieck). *Any rank k holomorphic vector bundle over \mathbb{CP}^1 is isomorphic to a direct sum of k complex line bundles $\mathcal{O}(n_i)$, $1 \leq i \leq k$, each with first Chern class n_i .*

The first Chern class completely classifies complex line bundles topologically. For us, $\mathcal{O}(n)$, $n \in \mathbb{Z}$ will mean a line bundle over $\mathbb{CP}^1 = \mathcal{U}_0 \cup \mathcal{U}_1$, where $\mathcal{U}_i = \{Z_i \neq 0\} \subset \mathbb{CP}^1$ for homogeneous coordinates $[Z_0, Z_1]^T$ on \mathbb{CP}^1 , with transition functions such that local trivialisations $(\lambda_i, v_i) \in \mathcal{U}_i \times \mathbb{C}$ are related on the overlap by $v_1 = \lambda_0^{-n} v_0$, where $\lambda_0 = \lambda_1^{-1} = Z_1/Z_0$. Note that $\mathcal{O}(-n)$ is dual to $\mathcal{O}(n)$, and $\mathcal{O}(n)$ is the tensor product of n copies of $\mathcal{O}(1)$. A section of $\mathcal{O}(n)$ is represented by functions $\sigma_i(\lambda_i)$ such that $(\lambda_0, \sigma_0(\lambda_0))$ and $(\lambda_1, \sigma_1(\lambda_1))$ correspond to the same point, i.e.

$$\sigma_1(\lambda_1) = \lambda_0^{-n} \sigma_0(\lambda_0).$$

If we expand these as power series in the local coordinates and use the fact that $\lambda_1 = \lambda_0^{-1}$, we find by equating coefficients that they are polynomials of degree at most n , making the space of global holomorphic sections $(n+1)$ –dimensional for $n \geq 0$.

The nonlinear graviton Theorem can now be stated as follows.

Theorem 5.1.2 ([62]). *There is a one–to–one correspondence between holomorphic ASD conformal structures and three dimensional complex manifolds containing a four parameter family of \mathbb{CP}^1 embeddings with normal bundle $\mathcal{O}(1) \oplus \mathcal{O}(1)$.*

From the results of Kodaira [44] we have that a vector at a point $m \in M$ corresponds to a holomorphic section of the normal bundle $\mathcal{O}(1) \oplus \mathcal{O}(1)$ of the curve \mathcal{L}_m in \mathcal{T} (which we know from above belongs to a four dimensional space). Penrose shows that we obtain an ASD conformal structure from \mathcal{T} by defining a vector to be null if the corresponding holomorphic section of $\mathcal{O}(1) \oplus \mathcal{O}(1)$ has a zero. This is the infinitesimal version of the intersection condition on \mathcal{L}_{m_1} and \mathcal{L}_{m_2} above.

(Anti–)self–duality in the sense of Calderbank

Although a β –distribution is intrinsically ASD in the sense that it is defined by an ASD two–form, there are two subclasses of β –distribution which we shall call *Calderbank–SD* or *Calderbank–ASD*. In [14], Calderbank associates to any given β –distribution a unique connection. When the curvature of this connection is (A)SD, the β –distribution is called Calderbank–(A)SD. This has nothing to do with the anti–self–duality of the two–form defining the β –distribution; in a sense, a β –distribution which is Calderbank–ASD is doubly ASD. See [14] (also [82]) for further details.

We can express Calderbank (anti–)self–duality in twistor language as follows. Let D_β be a β –distribution defined by an ASD two–form $\Sigma_{ab} = \iota_A \iota_B \epsilon_{A'B'}$, and such that the spinor ι_A satisfies

$$\nabla_{A'(A} \iota_{B)} = \mathcal{A}_{A'(A} \iota_{B)} \tag{5.10}$$

where $d\mathcal{A}$ is an (A)SD Maxwell field. Then D_β is Calderbank–(A)SD.

Local characterisation of the Einstein manifolds (M, g)

A general ASD metric depends, in the real–analytic category, on six arbitrary functions of three variables [27]. Theorem 3.2.1 gives an explicit subclass of such metrics which are additionally constrained by the following local characterisation.

Theorem 5.1.3 ([28]). *Let (M, g) be an real ASD Einstein manifold with scalar curvature 24 admitting a totally null distribution D_β which is Calderbank–ASD and parallel in the sense that ${}^g\nabla_\xi\tilde{\xi}\in\Gamma(D_\beta)$ for all $\xi\in\Gamma(TM)$, $\tilde{\xi}\in\Gamma(D_\beta)$. Then (M, g) is conformally flat, or it is locally isometric to (5.1).*

In our coordinates D_β is the kernel of the two–form $\Sigma = dx^{0'} \wedge dx^{1'}$, and can be written as $D_\beta = \text{span}\{\partial/\partial z_{0'}, \partial/\partial z_{1'}\}$. We find that

$${}^g\nabla\Sigma = 6\mathcal{A}\otimes\Sigma, \quad (5.11)$$

where $d\mathcal{A} = \Omega$, and Ω is the symplectic form on M . Writing $\Sigma_{ab} = \iota_A\iota_B\epsilon_{A'B'}$, (5.11) implies (5.10) for a rescaling of \mathcal{A} , so it is the anti–self–duality of Ω which makes D Calderbank–ASD.

In Section 5.4 we will consider the model case where M is constructed from the flat projective structure on \mathbb{RP}^2 . In this case, we can explicitly describe the ASD Maxwell two–form Ω in terms of the twistor space of M , and we will find that M carries a so–called *pseudo–hyper–Hermitian* structure, in which \mathcal{A} plays an important role.

5.1.2 Einstein–Weyl structures

Definition 5.1.4. *A Weyl Structure $(\mathcal{W}, \mathcal{D}, [h])$ is a conformal equivalence class of metrics $[h]$ on a manifold \mathcal{W} along with a fixed torsion–free affine connection \mathcal{D} which preserves any representative $h\in[h]$ up to conformal class. That is, for some one-form φ ,*

$$\mathcal{D}h = \varphi\otimes h.$$

A pair (h, φ) uniquely defines the connection and hence the Weyl structure, so we can alternatively specify a Weyl structure as a triple $(\mathcal{W}, h, \varphi)$. However, there is an equivalence class of such pairs which define the same Weyl structure. These are related by transformations

$$h \rightarrow \rho^2 h, \quad \varphi \rightarrow \varphi + 2d\ln(\rho), \quad (5.12)$$

where ρ is a smooth, non-zero function on \mathcal{W} . Physically, the Weyl condition in Lorentzian signature corresponds to the statement that null geodesics of the conformal structure $[h]$ are also geodesics of the connection \mathcal{D} .

If additionally the symmetric part of the Ricci tensor of \mathcal{D} is a scalar multiple of h , then \mathcal{W} is said to carry an Einstein–Weyl structure. This condition is invariant under (5.12). A trivial Einstein–Weyl structure is one whose one–form φ is closed, so that it is locally exact and thus may be set to zero by a change of scale (5.12). Then \mathcal{D} is the Levi–Civita connection of some representative $h \in [h]$, and this representative is Einstein.

In three dimensions, the Einstein–Weyl equations give a set of five non–linear partial differential equations on the pair (h, φ) which are integrable by the twistor transform of Hitchin [39].

Theorem 5.1.5 ([39]). *There is a one–to–one correspondence between three dimensional Einstein–Weyl structures and two dimensional complex manifolds containing a three parameter family of \mathbb{CP}^1 embeddings with normal bundle $\mathcal{O}(2)$.*

The conformal structure $[h]$ is obtained by demanding that a vector on \mathcal{W} is null if and only if the corresponding section of $\mathcal{O}(2)$ has a single zero. This condition is equivalent to the quadratic function $\sigma(\lambda)$ which represents the section having vanishing discriminant.

Hitchin’s results can be regarded as a reduction of Penrose’s twistor transform for ASD conformal structures by the following theorem of Jones and Tod.

Theorem 5.1.6. [40]

1. *Let (M, g) be a neutral signature, conformally ASD four–manifold with a conformal Killing vector K . Let*

$$h = |K|^{-2}g - |K|^{-4}\mathbf{K} \odot \mathbf{K}, \quad \varphi = \frac{2}{|K|^2} \star (\mathbf{K} \wedge d\mathbf{K}), \quad (5.13)$$

where $|K|^2 = g(K, K)$, $\mathbf{K} = g(K, \cdot)$ and \star is the Hodge operator defined by g . Then (h, φ) is a solution of the Einstein–Weyl equations on the space of orbits \mathcal{W} of K in M .

2. *Given an Einstein–Weyl structure $(\mathcal{W}, h, \varphi)$ there is a one–to–one correspondence between solutions (\mathcal{V}, α) to the abelian monopole equation*

$$d\mathcal{V} + \frac{1}{2}\varphi\mathcal{V} = \star_h d\alpha. \quad (5.14)$$

on \mathcal{W} , where \mathcal{V} is a function and α is a one-form, and conformally ASD four-metrics

$$g = \mathcal{V}h - \mathcal{V}^{-1}(dx + \alpha)^2 \quad (5.15)$$

over \mathcal{W} with an isometry $K = \partial/\partial x$.

If a metric with ASD Weyl tensor has more than one conformal symmetry, then distinct Einstein–Weyl structures are obtained on the space of orbits of conformal Killing vectors which are not conjugate with respect to an isometry [60].

5.1.3 The $SU(\infty)$ –Toda equation

The $SU(\infty)$ –Toda equation is given by

$$U_{XX} + U_{YY} = \epsilon(e^U)_{ZZ}, \quad \text{where } U = U(X, Y, Z), \quad \text{and } \epsilon = \pm 1 \quad (5.16)$$

Equation (5.16) has originally arisen in the context of complex general relativity [35, 8, 65], and then in Einstein–Weyl [79] and (in Riemannian context, with $\epsilon = -1$) scalar–flat Kähler geometry [48]. It belongs to a class of dispersionless systems integrable by the twistor transform [55, 25, 3], the method of hydrodynamic reduction [34], and the Manakov–Santini approach [51]. The equation is nevertheless not linearisable and most known explicit solutions admit Lie point or other symmetries (there are exceptions - see [11, 12, 54, 67]).

The $SU(\infty)$ –Toda equation is related to a subclass of Einstein–Weyl structures by the following result of Tod which improved the earlier result of Przanowski [65].

Theorem 5.1.7. [74] *Let $(\mathcal{W}, h, \varphi)$ be an Einstein–Weyl structure arising from the first part of Theorem 5.1.6, under the additional assumption that the ASD conformal structure $(M, [g])$ has a representative $g \in [g]$ which is Einstein with non–zero Ricci scalar. Then there exists $h \in [h]$, and coordinates (X, Y, Z) on an open set in \mathcal{W} such that (assuming the signature of h is $(2, 1)$ and the one-form dZ corresponds to a time–like vector)*

$$h = e^U(dX^2 + dY^2) - dZ^2, \quad \varphi = 2U_Z dZ \quad (5.17)$$

and the function $U = U(X, Y, Z)$ satisfies the $SU(\infty)$ –Toda equation (5.16) with $\epsilon = 1$.

Note that the assumptions about the signature of h and the timelike character of dZ in the above theorem are satisfied for all the Einstein–Weyl structures that can be obtained from the projective to Einstein correspondence. An invariant procedure for obtaining the coordinate system (X, Y, Z) is discussed in section 5.2.1.

5.1.4 Projective structures on a surface

Recall (see, for example, [10]) that a projective structure on a surface can be locally specified by a single second order ordinary differential equation: taking coordinates (x, y) on the surface we find that geodesics on which $\dot{x} \neq 0$ can be written as unparametrised curves $y(x)$ such that

$$y'' + a_0(x, y) + 3a_1(x, y)y' + 3a_2(x, y)(y')^2 + a_3(x, y)(y')^3 = 0, \quad (5.18)$$

where the coefficients $\{a_i\}$ are given by the projectively invariant formulae

$$a_0 = \Gamma_{00}^1, \quad 3a_1 = -\Gamma_{00}^0 + 2\Gamma_{01}^1, \quad 3a_2 = -2\Gamma_{01}^0 + \Gamma_{11}^1, \quad a_3 = -\Gamma_{11}^0.$$

Hitchin [39] solves the complexified version of (5.18) by the following twistor transform theorem.

Theorem 5.1.8. *There is a one-to-one correspondence between*

- *equivalence classes under coordinate transformations of complex ordinary differential equations of the form (5.18), where the coefficients a_i are holomorphic functions of x and y , and*
- *complex surfaces containing a two parameter family of \mathbb{CP}^1 embeddings with normal bundle $\mathcal{O}(1)$.*

In the case of the ordinary differential equation resulting from the flat projective structure on \mathbb{CP}^2 , the corresponding twistor space, whose points correspond to projective lines in \mathbb{CP}^2 , is the dual projective surface \mathbb{CP}_2 , and its \mathbb{CP}^1 embeddings are given by projective lines in \mathbb{CP}_2 . In analogy with the nonlinear graviton, we can define the correspondence space \mathcal{F} such that a point in \mathcal{F} is given by a point $p \in \mathbb{CP}^2$ and a projective line (or equivalently a direction) through p . This makes \mathcal{F} the projectivised tangent bundle $\mathbb{P}(T\mathbb{CP}^2)$, or equivalently $\mathbb{P}(T\mathbb{CP}_2)$.

As we saw in Chapter 3, the maximally symmetric projective surface \mathbb{RP}^2 has symmetry group $SL(3, \mathbb{R})$. In fact, the possible symmetry groups of projective surfaces are $SL(3, \mathbb{R})$, $SL(2, \mathbb{R})$, the two dimensional affine group, and \mathbb{R} . A partial classification is given in [46].

1. **On the flat projective surface \mathbb{RP}^2** described in Section 3.1.3, geodesics $y(x)$ are described in inhomogeneous coordinates $(x, y) = (P^0/P^2, P^1/P^2)$ by the ordinary differential equation

$$y'' = 0.$$

2. **The punctured plane** $\mathbb{R}^2 \setminus \{0\}$ has symmetry group $SL(2, \mathbb{R})$ acting via its fundamental representation. In this case there is a one parameter family of projective structures falling into three distinct equivalence classes. For simplicity we will consider only one of the classes, with geodesics $y(x)$ described by the differential equation

$$y'' = -(y - xy')^3, \quad (5.19)$$

where (x, y) are standard Euclidean coordinates on \mathbb{R}^2 .

3. **The two dimensional Lie group of affine transformations on \mathbb{R}** , which we denote $\text{Aff}(1)$, is generated by the unique non-abelian two dimensional Lie algebra $\{\mathfrak{v}_1, \mathfrak{v}_2\}$, where we choose a basis such that $[\mathfrak{v}_1, \mathfrak{v}_2] = \mathfrak{v}_1$. We can choose coordinates on $\text{Aff}(1)$ such that these correspond to vector fields

$$\frac{\partial}{\partial y}, \quad \frac{\partial}{\partial x} + y \frac{\partial}{\partial y},$$

and using invariance under these vector fields, the geodesic equation can be cast in the form [33]

$$y'' = e^{-2x}(y')^3 + A_1 y' + A_2 e^x,$$

where A_1 and A_2 are constants.

4. **The general projective surface with a symmetry**, after a choice of coordinates such that the symmetry is $\frac{\partial}{\partial x}$, corresponds to a set of geodesics $y(x)$ which satisfy an ordinary differential equation that can be written uniquely in the form [33]

$$y'' = A(y)(y')^3 + B(y)(y')^2 + 1. \quad (5.20)$$

Note that each of these classes of projective structures forms a subset of the next, and this can be seen explicitly by some changes of coordinates. For example, the general projective surface with a symmetry is flat when $A(y) = B(y) = 0$.

5.1.5 From projective surfaces to $SU(\infty)$ -Toda fields

The whole construction can now be summarised in the following diagram

$$\begin{array}{ccc} \text{Projective structure with symmetry} & \xrightarrow{\text{Thm 3.2.1}} & \text{ASD Einstein with symmetry} \\ \downarrow & & \downarrow \text{Thm 5.1.6} \\ \text{Solution to } SU(\infty) \text{ Toda} & \xleftarrow{\text{Thm 5.1.7}} & \text{Einstein-Weyl.} \end{array} \quad (5.21)$$

We now consider each of the above projective structures in turn, constructing the corresponding ASD Einstein manifold (M, g) and discussing some examples of Einstein–Weyl structures and $SU(\infty)$ –Toda fields that can be obtained from them.

5.2 The most general case

Consider the most general Einstein–Weyl structure arising from the combination of Theorem 3.2.1 and Theorem 5.1.6. Because of the correspondence (3.25) between symmetries of (M, g) and symmetries of the projective surface $(N, [\nabla])$, the construction must begin with the general projective surface with at least one symmetry.

By trial and error, we chose a representative connection for (5.20) such that the metric (5.1) had the simplest possible form. The choice of connection we took was

$$\Gamma_{11}^0 = A(y), \quad \Gamma_{00}^1 = -1, \quad \Gamma_{11}^1 = -B(y)$$

with all other components vanishing. Note that this choice of connection has a symmetric Ricci tensor, so the Schouten tensor is also symmetric and the symplectic form (5.2) pulls back to just $dz_{A'} \wedge dx^{A'}$. Thus we can write the Maxwell potential \mathcal{A} which is such that $d\mathcal{A} = \Omega$ as $\mathcal{A} = z_{A'} dx^{A'}$. Writing $x^{A'} = (x, y)$, $z_{A'} = (p, q)$, the resulting metric (5.1) is

$$g = (B(y) + p^2 + q)dx^2 + 2(pq + A(y))dxdy + (-A(y)p + B(y)q + q^2)dy^2 + dxdp + dydq. \quad (5.22)$$

Factoring by $K = \frac{\partial}{\partial x}$ following the algorithm of Theorem 5.1.6, equation (5.13) gives the following form for the Einstein–Weyl structure.

Proposition 5.2.1. *The most general Einstein–Weyl structure arising from the procedure (5.21) is locally equivalent to*

$$\begin{aligned} h &= \frac{1}{\mathcal{V}} \left((Bq - Ap + q^2)dy + dq \right) dy - \left((pq + A)dy + \frac{1}{2}dp \right)^2, \\ \varphi &= \mathcal{V}(4dq + 2pdp), \quad \text{where} \quad \mathcal{V} = (B + p^2 + q)^{-1}. \end{aligned} \quad (5.23)$$

Here (p, q, y) are local coordinates on \mathcal{W} , $A(y), B(y)$ are arbitrary functions of y , and the solution to the monopole equation (5.14) arising from the second part of Theorem 5.1.6 is the pair (\mathcal{V}, α) , where

$$\alpha = \mathcal{V}(pq + A)dy + \frac{\mathcal{V}}{2}dp.$$

5.2.1 Solution to the $SU(\infty)$ –Toda equation

The procedure for extracting the corresponding solution to the $SU(\infty)$ –Toda equation is given in [74] (see also [48] and [26]). It involves finding the coordinates (X, Y, Z) that put the metric (5.23) in the form (5.17). Given an ASD Einstein metric (M, g) with a Killing vector K

1. The conformal factor $c : M \rightarrow \mathbb{R}^+$ given by

$$c = |d\mathbf{K} + *_g d\mathbf{K}|_g^{-1/2}$$

has a property that the rescaled self–dual derivative of K

$$\vartheta \equiv c^3 \left(\frac{1}{2} (d\mathbf{K} + *_g d\mathbf{K}) \right)$$

is parallel with respect to $c^2 g$. The metric $c^2 g$ is para–Kähler with self–dual para–Kähler form ϑ , and admits a Killing vector K , as $\mathcal{L}_K(c) = 0$.

2. Define a function $Z : M \rightarrow \mathbb{R}$ to be the moment map:

$$dZ = K \lrcorner \vartheta. \quad (5.24)$$

It is well defined, as the Kähler form is Lie–derived along K .

3. Construct the Einstein–Weyl structure of Theorem 5.1.6 by factoring $(M, c^2 g)$ by K . Restrict the metric h to a surface $Z = Z_0 = \text{const}$, and construct isothermal coordinates (X, Y) on this surface:

$$\gamma \equiv h|_{Z=Z_0} = e^U (dX^2 + dY^2), \quad U = U(X, Y, Z_0).$$

To implement this step chose an orthonormal basis of one–forms such that $\gamma = e_1^2 + e_2^2$. Now (X, Y) are solutions to the linear system of first order partial differential equations

$$(e_1 + ie_2) \wedge (dX + idY) = 0.$$

4. Extend the coordinates (X, Y) from the surface $Z = Z_0$ to \mathcal{W} . This may involve a Z –dependent affine transformation of (X, Y) .

Implementing the Steps 1–4 on MAPLE we find that if $A = 0$, and $B = B(y)$ is arbitrary, then the $SU(\infty)$ –Toda solution is given implicitly by

$$\begin{aligned} X &= -\frac{8e^{-2\int B(y)dy}Z^3p}{(Z^2p^2+4)^2}, \quad Y = \int e^{-2\int B(y)dy}dy + \frac{e^{-2\int B(y)dy}(-2Z^4p^2+8Z^2)}{(Z^2p^2+4)^2}. \\ U &= \ln\left(\frac{(Z^2p^2+4)^3}{64Z^2}\right) + 4\int B(y)dy. \end{aligned} \quad (5.25)$$

We can check that this is indeed a solution using the fact that the $SU(\infty)$ –Toda equation is equivalent to $d\star_h dU = 0$. We have also checked by performing a coordinate transformation of (5.16) to the coordinates (y, p, Z) .

To simplify the form of (5.25) set

$$G = \int \exp\left(-2\int B(y)dy\right), \quad T = \frac{2Z^2}{Z^2p^2+4}.$$

Then (5.25) becomes

$$e^U = \frac{Z^4}{8T^3(G')^2}, \quad Y = G + G'T\left(\frac{4T}{Z^2} - 1\right), \quad X^2 = \frac{4T^4(G')^2}{Z^2}\left(\frac{2}{T} - \frac{4}{Z^2}\right).$$

Eliminating (T, y) between these three equations gives one relation between (X, Y, Z) and U which is our implicit solution. The elimination can be carried over explicitly if $G = y^k$ for any integer k , or if $G = \exp y$. In the latter case the solution is given by

$$4Y^2e^U(e^U X^2 - Z^2)^3 + (2e^{2U}X^4 - 3e^U X^2Z^2 + Z^4 + 2Z^2)^2 = 0.$$

We can also consider the flat projective structure with $A = B = 0$, in which case the coordinate p can be eliminated between

$$e^U = \left(\frac{(Z^2p^2+4)^3}{64Z^2}\right), \quad X = -\frac{8Z^3p}{(Z^2p^2+4)^2}$$

by taking a resultant. This yields

$$e^U(e^U X^2 - Z^2)^3 + Z^4 = 0.$$

Note that even the flat projective surface can yield a non–trivial solution to the Toda equation; further discussion can be found in Section 5.4.5.

5.2.2 Two monopoles

The Einstein–Weyl structures $(\mathcal{W}, h, \varphi)$ in (5.23) that we have constructed in Proposition 5.2.1 are special, as they belong to the $SU(\infty)$ –Toda class. The general solution to the $SU(\infty)$ –Toda equation depends (in the real analytic category) on two arbitrary functions of two variables, but the solutions of the form (5.23) depend on two functions of one variable. The additional constraints on the solutions can be traced back to the four dimensional ASD conformal structures which give rise (by the Jones–Tod construction) to (5.23). In what follows we shall point out how some of the additional structure on \mathcal{W} arises as a couple of solutions to the abelian monopole equation.

Let us call the solution (\mathcal{V}, α) arising in Proposition 5.2.1 the Einstein monopole, as the resulting conformal class contains an Einstein metric (5.22). The second solution $(\mathcal{V}_M, \alpha_M)$ (which we shall call the Maxwell monopole) arises as a symmetry reduction of the ASD Maxwell potential

$$\mathcal{A} = pdx + qdy = -\mathcal{V}_M \mathbf{K} + \alpha_M,$$

where \mathbf{K} is the Killing one–form, and we find

$$\mathcal{V}_M = -p\mathcal{V}, \quad \alpha_M = qdy - p\alpha.$$

5.3 The submaximally symmetric case

Choosing a representative connection from the projective class defined by (5.19), we obtain from (5.1) an Einstein metric

$$g = (p^2 - xy^2p - y^3q + 4y^2)dx^2 + 2(pq + x^2yp + xy^2q - 4xy)dxdy + (q^2 - x^3p - x^2yq + 4x^2)dy^2 + dxdp + dydq \quad (5.26)$$

on M , again with $z_0 =: p$, $z_1 =: q$, having Killing vectors

$$K_1 = x\frac{\partial}{\partial x} - p\frac{\partial}{\partial p} - y\frac{\partial}{\partial y} + q\frac{\partial}{\partial q}, \quad K_2 = x\frac{\partial}{\partial y} - q\frac{\partial}{\partial p}, \quad K_3 = y\frac{\partial}{\partial x} - p\frac{\partial}{\partial q}.$$

These are lifts of the projective vector fields corresponding to the $\mathfrak{sl}(2, \mathbb{R})$ elements

$$\mathfrak{v}_1 = \begin{pmatrix} \epsilon & 0 \\ 0 & -\epsilon \end{pmatrix} \quad \mathfrak{v}_2 = \begin{pmatrix} 0 & 0 \\ \epsilon & 0 \end{pmatrix} \quad \mathfrak{v}_3 = \begin{pmatrix} 0 & \epsilon \\ 0 & 0 \end{pmatrix}.$$

To obtain an example of a Jones–Tod reduction of (5.26), we factor by K_3 . Choosing coordinates

$$r = \frac{p^2}{y^2}, \quad z = 2 \ln(y^2), \quad w = xp + yq,$$

gives an Einstein–Weyl structure

$$\begin{aligned} h &= -dr^2 - 2drdw - w(w^2 + r - 5w + 4)dz^2 + 2(r - w + 4)dzdw, \\ \varphi &= \frac{1}{r - w + 4}dr - \frac{3w}{r - w + 4}dz - \frac{4}{r - w + 4}dw. \end{aligned} \quad (5.27)$$

The solution to the $SU(\infty)$ –Toda equation (5.16) which determines the Einstein–Weyl structure (5.27) is described by an algebraic curve $f(e^U, X, Y, Z) = 0$ of degree six in e^U and degree twelve in the other coordinates. This solution has been found following the Steps 1–4 in Section 5.2.1, and is given by

$$\begin{aligned} &64e^{6U}X^6(X+Y)^3(X-Y)^3 - 92e^{5U}X^4Z^2(X+Y)^3(X-Y)^3 \\ &+ 48e^{4U}X^2Z^2(5X^6Z^2 - 14X^4Y^2Z^2 + 13X^2Y^2Z^2 - 4Y^4Z^2 + 9X^4 + 27X^2) \\ &+ 8e^{3U}Z^4(-20X^6Z^2 + 48X^4Y^2Z^2 - 36X^2Y^4Z^2 + 8Y^6Z^2 - 81X^4 - 243X^2Y^2) \\ &+ 3e^{2U}Z^4(20X^4Z^4 - 36X^2Y^2Z^4 + 16Y^4Z^4 + 108X^2Z^2 + 216Y^2Z^2 + 243) \\ &+ 6e^UZ^8(-2X^2Z^2 + 2Y^2Z^2 - 9) + Z^{12} \\ &= 0. \end{aligned}$$

Note that the formulae (5.27) are independent of the coordinate z , and therefore have a symmetry. This was unexpected because there is no other symmetry of (M, g) that commutes with K_3 . However, it is possible for symmetries to appear in the Einstein–Weyl structure without a corresponding symmetry of the ASD conformal structure. This can be seen from the general formula (5.15); the function \mathcal{V} may depend on the coordinate z so that g depends on z even though h does not. For example, the Gibbons–Hawking metrics [37] give a trivial Einstein–Weyl structure with the maximal symmetry group, but the four-metric is in general not so symmetric. Our discovery of this unexpected symmetry motivated a more concrete description of a symmetry of a Weyl structure.

Definition 5.3.1. An infinitesimal symmetry of a Weyl structure $(\mathcal{W}, \mathcal{D}, [h])$ is a vector field \mathcal{K} which is both an affine vector field with respect to the connection³ \mathcal{D} and a conformal Killing vector with respect to the conformal structure $[h]$.

Proposition 5.3.2. Given an infinitesimal symmetry \mathcal{K} of a Weyl structure $(\mathcal{W}, \mathcal{D}, [h])$ in dimension N , and a representative $h \in [h]$ such that $\mathcal{D}h = \varphi \otimes h$, there exists a smooth function $f : \mathcal{W} \rightarrow \mathbb{R}$ such that

$$\mathcal{L}_{\mathcal{K}}h = fh, \quad \mathcal{L}_{\mathcal{K}}\varphi = \frac{1}{N}d[\mathcal{K} \lrcorner d(\ln(\det(h)))]. \quad (5.28)$$

Proof. The first equation follows immediately from the fact that \mathcal{K} is a conformal Killing vector of h . It remains to evaluate the Lie derivative of the one-form φ along the flow of \mathcal{K} given that $\mathcal{L}_{\mathcal{K}}h = fh$ and $\mathcal{L}_{\mathcal{K}}\Gamma_{jk}^i = 0$, where Γ_{jk}^i are the components of the connection \mathcal{D} . We do this by considering the Lie derivative of $\mathcal{D}h$:

$$\begin{aligned} \mathcal{L}_{\mathcal{K}}(\mathcal{D}_i h_{jk}) &= \mathcal{L}_{\mathcal{K}}(\partial_i h_{jk}) - \mathcal{L}_{\mathcal{K}}(\Gamma_{ji}^l h_{lk} + \Gamma_{ki}^l h_{jl}) \\ &= \mathcal{L}_{\mathcal{K}}(\partial_i h_{jk}) - f(\Gamma_{ji}^l h_{lk} + \Gamma_{ki}^l h_{jl}). \end{aligned}$$

Now

$$\begin{aligned} \mathcal{L}_{\mathcal{K}}(\partial_i h_{jk}) &= \mathcal{K}^l \partial_l \partial_i h_{jk} + (\partial_i \mathcal{K}^l) \partial_l h_{jk} + (\partial_j \mathcal{K}^l) \partial_i h_{lk} + (\partial_k \mathcal{K}^l) \partial_i h_{jl} \\ &= \partial_i [\mathcal{K}^l \partial_l h_{jk} + (\partial_j \mathcal{K}^l) h_{lk} + (\partial_k \mathcal{K}^l) h_{jl}] - (\partial_i \partial_j \mathcal{K}^l) h_{lk} - (\partial_i \partial_k \mathcal{K}^l) h_{jl}. \end{aligned}$$

The term with square brackets is just

$$\partial_i(\mathcal{L}_{\mathcal{K}}h_{jk}) = \partial_i(fh_{jk}) = f\partial_i h_{jk} + \partial_i f h_{jk},$$

so we have

$$\mathcal{L}_{\mathcal{K}}(\mathcal{D}_i h_{jk}) = f\mathcal{D}_i h_{jk} + \partial_i f h_{jk} - (\partial_i \partial_j \mathcal{K}^l) h_{lk} - (\partial_i \partial_k \mathcal{K}^l) h_{jl}.$$

Setting this equal to $\mathcal{L}_{\mathcal{K}}(\varphi_i h_{jk}) = (\mathcal{L}_{\mathcal{K}}\varphi_i) h_{jk} + f\varphi_i h_{jk}$ and cancelling $f\varphi_i h_{jk}$ with $f\mathcal{D}_i h_{jk}$, we find

$$\begin{aligned} (\mathcal{L}_{\mathcal{K}}\varphi_i) h_{jk} &= \partial_i f h_{jk} - (\partial_i \partial_j \mathcal{K}^l) h_{lk} - (\partial_i \partial_k \mathcal{K}^l) h_{jl} \\ \implies \mathcal{L}_{\mathcal{K}}\varphi_i &= \partial_i f - \frac{2}{N} \partial_i \partial_j \mathcal{K}^j. \end{aligned} \quad (5.29)$$

³Recall that an affine vector field of a connection \mathcal{D} is one which preserves its components, i.e. $\mathcal{L}_{\mathcal{K}}\Gamma_{jk}^i = 0$.

Finally, we note that

$$\partial_i \partial_j \mathcal{K}^j = \frac{N}{2} \partial_i f - \frac{1}{2} \partial_i [\mathcal{K} \lrcorner d(\ln(\det(h)))].$$

This follows from tracing the expression $\mathcal{L}_{\mathcal{K}} h_{ij} = f h_{ij}$:

$$\begin{aligned} \mathcal{L}_{\mathcal{K}} h_{ij} &= \mathcal{K}^k \partial_k h_{ij} + (\partial_i \mathcal{K}^k) h_{kj} + (\partial_j \mathcal{K}^k) h_{ik} = f h_{ij} \\ \implies \mathcal{K}^k h^{ij} \partial_k h_{ij} + 2\partial_k \mathcal{K}^k &= Nf \\ \implies 2\partial_i \partial_k \mathcal{K}^k &= N\partial_i f - \partial_i (\mathcal{K}^k h^{jl} \partial_k h_{jl}) \end{aligned}$$

and recalling that $h^{jl} \partial_k h_{jl} = \partial_k \ln(\det(h))$. Substituting into (5.29) then yields the result. \square

We can easily verify the invariance of (5.28) under Weyl transformations. Let $(\bar{h}, \bar{\varphi})$ be a new metric and one-form related to the old ones by (5.12). Then

$$\mathcal{L}_{\mathcal{K}} \bar{\varphi} = \mathcal{L}_{\mathcal{K}} \varphi + 2d[\mathcal{K} \lrcorner d\ln(\rho)]$$

from (5.12), and from (5.28) we have

$$\begin{aligned} \mathcal{L}_{\mathcal{K}} \bar{\varphi} &= \frac{1}{N} d[\mathcal{K} \lrcorner d(\ln(\rho^{2N} \det(h)))] \\ &= \frac{1}{N} d[\mathcal{K} \lrcorner d(\ln(\det(h)))] + \frac{2N}{N} d[\mathcal{K} \lrcorner d\ln(\rho)] \\ &= \mathcal{L}_{\mathcal{K}} \varphi + 2d[\mathcal{K} \lrcorner d\ln(\rho)], \end{aligned}$$

as above. Note that the function f in (5.28) will change according to

$$\bar{f} = f + 2\mathcal{K} \lrcorner d\ln\rho.$$

In the case of the Weyl structure (5.27), the infinitesimal symmetry is

$$\mathcal{K} = \frac{\partial}{\partial z}.$$

Since we have chosen a scale such that \mathcal{K} is in fact a Killing vector of h , we have that $\mathcal{K} \lrcorner d(\ln(\det(h))) = 0$, so the one-form φ is also preserved by \mathcal{K} . This is consistent with the fact that it has no explicit z –dependence.

5.4 The model case

In the following section we discuss the four–manifold (M, g) obtained from the maximally symmetric flat projective surface $N = \mathbb{RP}^2$. In this case, g is not only almost para–Kähler but in fact para–Kähler, since the symplectic form Ω is parallel with respect to the Levi–Civita connection of g . Choosing a representative connection with $\Gamma_{A'B'}^{C'} = 0$ gives g as

$$g = dz_{A'} \odot dx^{A'} + z_{A'} z_{B'} dx^{A'} \odot dx^{B'}. \quad (5.30)$$

We begin by discussing the conformal structure of (5.30), both explicitly and in terms of its twistor space. We then note a pseudo–hyper–Hermiticity property which is unique to the model case, and find some special structure on the twistor space. Finally, we present a classification of the Einstein–Weyl structures which can be obtained from (5.30) by Jones–Tod factorisation, and exhibit an explicit example of such a factorisation from the twistor perspective, reconstructing the conformal structure on \mathcal{W} from minitwistor curves.

5.4.1 Conformal Structure on M

Recall that points $P \in \mathbb{R}^3$ and $L \in \mathbb{R}_3$ respectively define lines and planes in \mathbb{R}^3 which are preserved by multiplication of P, L by members of \mathbb{R}^* , and that these lines and planes respectively descend to points $[P]$ and lines $[L]$ in \mathbb{RP}^2 . In what follows we will drop the square brackets and understand points $[P] \in \mathbb{RP}^2$ and lines $[L] \in \mathbb{RP}_2$ to be represented by vectors $P \in \mathbb{R}^3$ and $L \in \mathbb{R}_3$. Let $M \subset \mathbb{RP}^2 \times \mathbb{RP}_2$ be the set of non–incident pairs (P, L) .

Proposition 5.4.1. *Two pairs (P, L) and (\tilde{P}, \tilde{L}) are null–separated with respect to the conformal structure (5.30) if there exists a line which contains the three points $(P, \tilde{P}, L \cap \tilde{L})$.*

Proof. First note that the null condition of Proposition 5.4.1 defines a co–dimension one cone in TN : generically there is no line through three given points. To make explicit the condition for such a line to exist, consider two pairs (P, L) and (\tilde{P}, \tilde{L}) of non–incident points and lines. By thinking of L, \tilde{L} as normal vectors to planes in \mathbb{R}^3 , we see that $L + t\tilde{L}$ is a plane which intersects L and $t\tilde{L}$ at their intersection, thus defining a line in \mathbb{RP}^2 which intersects the lines $L, \tilde{L} \subset \mathbb{RP}^2$ at their intersection.

If $P, \tilde{P}, L \cap \tilde{L}$ are co–linear then there exists t such that both P and \tilde{P} lie on $L + t\tilde{L}$, i.e.

$$P \cdot (L + t\tilde{L}) = 0, \quad \tilde{P} \cdot (L + t\tilde{L}) = 0. \quad (5.31)$$

Eliminating t from (5.31) gives

$$(P \cdot L)(\tilde{P} \cdot \tilde{L}) - (\tilde{P} \cdot L)(P \cdot \tilde{L}) = 0.$$

Setting $\tilde{P} = P + dP, \tilde{L} = L + dL$ yields a metric g representing the conformal structure

$$g = \frac{dP \cdot dL}{P \cdot L} - \frac{1}{(P \cdot L)^2} (L \cdot dP)(P \cdot dL).$$

We can use the normalisation $P \cdot L = 1$, so that $P \cdot dL = -L \cdot dP$, and

$$g = dP \cdot dL + (L \cdot dP)^2. \quad (5.32)$$

We take affine coordinates

$$P = [x^{A'}, 1], \quad L = [z_{A'}, 1 - x^{A'} z_{A'}] \quad (5.33)$$

with a normalisation $P \cdot L = 1$ to recover the metric (5.30).

□

5.4.2 Twistor space of M

To understand $(M, [g])$ from the twistor perspective, we need to move to the complex picture. In what follows, we will view M as the set of non–incident pairs in $\mathbb{CP}^2 \times \mathbb{CP}_2$. Let $F_{12}(\mathbb{C}^3) \subset \mathbb{CP}^2 \times \mathbb{CP}_2$ be set of incident pairs (p, l) , so that $p \cdot l = 0$. Note that, since l and p correspond to planes and lines in \mathbb{C}^3 respectively, and since $p \cdot l = 0$ is the condition for the line p lying in the plane l , $F_{12}(\mathbb{C}^3)$ coincides with the flag manifold of type $(1, 2)$ in \mathbb{C}^3 , i.e. the collection of one and two dimensional vector subspaces (p, l) in \mathbb{C}^3 such that $p \subset l$. This is the twistor space of (M, g) . A \mathbb{CP}^1 embedding corresponding to a point $(P, L) \in M$ consists of all lines l thorough P , and all points $p = l \cap L$:

$$P \cdot l = 0, \quad p \cdot L = 0, \quad p \cdot l = 0. \quad (5.34)$$

Let (P, L) and (\tilde{P}, \tilde{L}) be points in M . These uniquely define a point $p = L \cap \tilde{L} \in \mathbb{CP}^2$ and line $l \subset \mathbb{CP}^2$ such that $P, \tilde{P} \in l$ given by

$$p = L \wedge \tilde{L}, \quad l = P \wedge \tilde{P},$$

where $[L \wedge \tilde{L}]^\alpha = \epsilon^{\alpha\beta\gamma} L_\alpha \tilde{L}_\beta$ etc. The pair (p, l) lies in F_{12} if p lies on l , i.e. if $p \cdot l = 0$, so that (P, L) and (\tilde{P}, \tilde{L}) are null-separated with respect to the conformal structure (5.31). Then (p, l) is the intersection of the \mathbb{CP}^1 embeddings corresponding to (P, L) and (\tilde{P}, \tilde{L}) .

We shall now give an explicit parametrisation of twistor lines, and show how the metric (5.32) arises from the Penrose condition [62, 78]. Let $P \in \mathbb{CP}^2$. The corresponding $l \in \mathbb{CP}_2$ is represented by some normal vector which is perpendicular to P in \mathbb{C}^3 , i.e.

$$l = P \wedge \pi, \quad \text{where } \pi \sim a\pi + bP, \quad (5.35)$$

where $a \in \mathbb{C}^*, b \in \mathbb{C}$. Thus π parametrises a projective line \mathbb{CP}^1 , and by making a choice of b we can take $\pi = [\pi^0, \pi^1, 0]$, where $\pi^{A'} = [\pi^0, \pi^1] \in \mathbb{CP}^1$. The constraint $P \cdot l = 0$ now holds. To satisfy the remaining constraints in (5.34) we take

$$p = L \wedge l = (L \cdot \pi)P - (L \cdot P)\pi. \quad (5.36)$$

Substituting (5.33) gives the corresponding twistor line parametrised by $[\pi] \in \mathbb{CP}^1$

$$p^\alpha = [(z_{B'} \pi^{B'}) x^{A'} - \pi^{A'}, z_{B'} \pi^{B'}], \quad l_\alpha = [\pi_{A'}, -\pi_{B'} x^{B'}], \quad (5.37)$$

where the spinor indices are raised and lowered with ϵ^{AB} and its inverse.

We shall now derive the expression for the conformal structure using the nonlinear graviton prescription described in Section 5.1.1. To compute the normal bundle, let $([l(\pi, P, L)], [p(\pi, P, L)])$ be the twistor line corresponding to a point $m = (P, L)$ in M . The vector in the direction of a nearby point $(P + \delta P, L + \delta L)$ corresponds to the neighbouring line $([l + \delta l], [p + \delta p])$, where from (5.35) and (5.36) we have

$$\delta l = \delta P \wedge \pi, \quad \delta p = (\delta L \cdot \pi)P + (L \cdot \pi)\delta P - \delta(L \cdot P)\pi.$$

The lines $(l + \delta l, p + \delta p)$ and (l, p) define a section of the normal bundle to (l, p) , which has a zero if and only if this vector is null. Vanishing of the section is equivalent to intersection of the two lines, and this happens if there exists some $[\pi]$ such that $l + \delta l \sim l$ and $p + \delta p \sim p$ (note that the intersection point, if it exists, is unique, since

projective lines in \mathbb{CP}^2 cannot meet more than once). We thus find

$$l + \delta l \sim l \iff \pi \sim \delta P = [\delta x^{0'}, \delta x^{1'}, 0].$$

And $p + \delta p \sim p \iff$

$$0 = p \wedge \delta p = (L \cdot \pi)^2 P \wedge \delta P - (L \cdot P)(\delta L \cdot \pi) \pi \wedge P - (L \cdot \pi) \delta(L \cdot P) P \wedge \pi - (L \cdot P)(L \cdot \pi) \pi \wedge \delta P.$$

Substituting $\pi \sim \delta P$, we find that all terms on the right hand side are proportional to $P \wedge \delta P = [0, 0, x^{B'} dx_{B'}]$, with

$$(L \cdot \delta P)^2 - (L \cdot \delta P) \delta(L \cdot P) + (L \cdot P) (\delta L \cdot \delta P) = 0.$$

Setting $L \cdot P = 1$ this gives the conformal structure (5.32).

5.4.3 Pseudo–hyper–Hermitian structure on M

A pseudo–hyper–complex structure on a four manifold M is a triple of endomorphisms I_1, I_2, I_3 of TM which satisfy

$$I_1^2 = -Id, \quad I_2^2 = I_3^2 = Id, \quad I_1 I_2 I_3 = Id,$$

and such that $c_1 I_1 + c_2 I_2 + c_3 I_3$ is an integrable complex structure for any point on the hyperboloid $c_1^2 - c_2^2 - c_3^2 = 1$. A neutral signature metric g on a pseudo–hyper–complex four–manifold is pseudo–hyper–Hermitian if

$$g(\xi, \xi) = g(I_1 \xi, I_1 \xi) = -g(I_2 \xi, I_2 \xi) = -g(I_3 \xi, I_3 \xi)$$

for any vector field ξ on M .

Given a pseudo–hyper–complex structure $(M, \{I_1, I_2, I_3\})$ and any vector field ξ on M , the frame $(\xi, I_1 \xi, I_2 \xi, I_3 \xi)$ defines a conformal structure on M . With a natural choice of orientation which makes the fundamental two–forms of I_1, I_2, I_3 self–dual, this conformal structure is ASD.

Let $\Sigma^{A'B'}$ be a basis of SD two–forms on M . The following result is proved in [24] (see also [7]) in the Riemannian (i.e. hyper–complex) case.

Theorem 5.4.2 ([24]). *A four-manifold M equipped with a neutral signature metric g is pseudo-hyper-Hermitian if there exists a one-form A depending only on g such that*

$$d\Sigma^{A'B'} + A \wedge \Sigma^{A'B'} = 0.$$

In fact, this condition is necessary and sufficient for hyper-Hermiticity [24, 7]. Given some (M, g) which is conformally ASD, it can also be shown (see Lemma 2 in [24] and Theorem 7.1 in [13]) that a lack of vertical $\partial/\partial\pi$ terms in the twistor distribution (5.9) implies hyper-Hermiticity of (M, g) .

Proposition 5.4.3. *The Einstein metric (5.30) is pseudo-hyper-Hermitian.*

Proof. The null frame for the 4-metric is

$$e^{0A'} = dx^{A'}, \quad e^{1A'} = dz^{A'} + z^{A'}(z_{B'}dx^{B'}), \quad \text{so that} \quad g = \varepsilon_{A'B'}\varepsilon_{AB}e^{AA'}e^{BB'}. \quad (5.38)$$

Thus the forms $\Sigma = dx^{0'} \wedge dx^{1'}$ and $\Omega = dz_{A'} \wedge dx^{A'}$ are ASD. The basis of SD two forms is spanned by

$$dx \wedge dq + q^2 dx \wedge dy, \quad dx \wedge dp - dy \wedge dq + 2pq dx \wedge dy, \quad -dy \wedge dp + p^2 dx \wedge dy$$

or, in a more compact notation, by $\Sigma^{A'B'} = dx^{(A'} \wedge dz^{B')}$ + $z^{A'}z^{B'}\Sigma$. We can verify that

$$d\Sigma^{A'B'} + 2\mathcal{A} \wedge \Sigma^{A'B'} = 0, \quad (5.39)$$

where $\mathcal{A} = z_{A'}dx^{A'}$ is such that $d\mathcal{A} = \Omega$, so from Theorem 5.4.2 we have that M carries a hyper-Hermitian structure, and in fact the corresponding ASD Maxwell field $d\mathcal{A} = \Omega$ coincides with the one arising from the para-Kähler structure on M via (5.11).

Alternatively, note that the twistor distribution (5.9), having chosen the basis (5.38), is given by

$$L_0 = \pi^{A'} \frac{\partial}{\partial x^{A'}} + (z_{B'}\pi^{B'})z_{A'} \frac{\partial}{\partial z_{A'}}, \quad L_1 = \pi_{A'} \cdot \frac{\partial}{\partial z_{A'}}, \quad (5.40)$$

which have no vertical terms. We can easily verify that it is Frobenius integrable, as $[L_0, L_1] = -(\pi^{A'}z_{A'})L_1$. The SD part of the spin connection is given in terms of \mathcal{A} as $\Gamma_{AA'B'C'} = -2\mathcal{A}_{A(B'}\varepsilon_{C')A'}$.

□

In the next section we shall show how to encode \mathcal{A} in the twisted-photon Ward bundle over the twistor space of (M, g) .

5.4.4 A line bundle over the twistor space of M

Ward [76] shows that there is a correspondence between ASD Maxwell potentials on M and \mathbb{C}^* bundles over \mathcal{T} which are trivial on twistor lines⁴. The purpose of this section is to uncover the \mathbb{C}^* bundle over \mathcal{T} corresponding to the ASD Maxwell potential $\mathcal{A} = z_{A'} dx^{A'}$ on M .

The twistor space F_{12} described in Section 5.4.2 can be identified with the projectivised tangent bundle $\mathbb{P}(T\mathbb{CP}_2)$ of the minitwistor space of the flat projective structure, since a point (p, l) in $F_{12} \subset \mathbb{CP}^2 \times \mathbb{CP}_2$ consists of a point $l \in \mathbb{CP}_2$, and a line $p \subset \mathbb{CP}_2$ through l which we can identify with a direction in the tangent space $T_l \mathbb{CP}_2$. Thus the twistor space of M is the correspondence space (in a twistorial sense) of \mathbb{CP}^2 and its twistor space \mathbb{CP}_2 . An obvious \mathbb{C}^* bundle over $\mathbb{P}(T\mathbb{CP}_2)$ is $T\mathbb{CP}_2$.

Proposition 5.4.4. *The \mathbb{C}^* bundle $T\mathbb{CP}_2 \rightarrow \mathbb{P}(T\mathbb{CP}_2) = F_{12}$ is trivial on twistor lines, and corresponds via Ward’s twisted photon construction to the ASD Maxwell potential \mathcal{A} on M .*

Proof. There are many open sets needed to cover $\mathbb{P}(T\mathbb{CP}_2)$, but it is sufficient to consider two: \mathcal{U} , where $(l_1 \neq 0, p^2 \neq 0)$, and $(l_2/l_1, l_3/l_1, p^3/p^2)$ are coordinates, and $\tilde{\mathcal{U}}$ where $(l_1 \neq 0, p^3 \neq 0)$, and $(l_2/l_1, l_3/l_1, p^2/p^3)$ are coordinates. Now consider the total space of $T\mathbb{CP}_2$, and restrict it to the intersection of (pre–images in $T\mathbb{CP}_2$ of) \mathcal{U} and $\tilde{\mathcal{U}}$. The coordinates on $T\mathbb{CP}_2$ in these region are $(l_2/l_1, l_3/l_1, p^2/p^1, p^3/p^1)$, and the fibre coordinates over τ over \mathcal{U} and $\tilde{\tau}$ over $\tilde{\mathcal{U}}$ are related by⁵

$$\tilde{\tau} = \exp(F)\tau, \quad \text{where} \quad F = \ln(p_2/p_3).$$

Now we follow the procedure of [76]: restrict F to a twistor line, and split it. The holomorphic splitting is $F = f - \tilde{f}$, where $f = \ln(p_2)$ is holomorphic in the pre–image of \mathcal{U} in the correspondence space, and $\tilde{f} = \ln(p_3)$ is holomorphic in the pre–image of $\tilde{\mathcal{U}}$. Note that F is a twistor function, but f, \tilde{f} are not. Therefore $L_A F = 0$, where the twistor distribution L_A is given by (5.40). This implies that $L_A f = L_A \tilde{f}$. Since each side of this equation is holomorphic on an open subset of \mathbb{CP}^1 , and since \mathbb{CP}^1 can be covered with such subsets, both sides are globally holomorphic and therefore linear in

⁴Recall that a principal bundle $\mathbb{P} \rightarrow \mathcal{T}$ with structure group G is trivial if there exists a map $\chi : \mathbb{P} \rightarrow \mathcal{T} \times G$. Let $\{\chi_\alpha : \mathcal{U}_\alpha \rightarrow \mathcal{T} \times G\}$ be local trivialisations related by transition functions $F_{\alpha\beta} = \chi_\beta \circ \chi_\alpha^{-1} \in G$. If $F_{\alpha\beta} = f_\beta f_\alpha^{-1}$ for some splitting elements $f_\alpha, f_\beta \in G$ on $\mathcal{U}_\alpha, \mathcal{U}_\beta$ respectively, then there exist $\tilde{\chi}_\alpha = f_\alpha^{-1} \circ \chi_\alpha$ and $\tilde{\chi}_\beta = f_\beta^{-1} \circ \chi_\beta$ such that $\tilde{F}_{\alpha\beta} = f_\beta^{-1} f_\beta f_\alpha^{-1} f_\alpha = Id$, so that the bundle is trivial.

⁵Here we are following Ward [76], and thinking of a \mathbb{C}^* bundle.

$\pi^{A'}$ by the Liouville theorem. Hence

$$L_A f = L_A \tilde{f} = \pi^{A'} \mathcal{A}_{AA'}$$

for some one-form \mathcal{A} on M .

To construct this one-form recall the parametrisation of twistor curves (5.37). This gives

$$f = \ln(z_{A'} \pi^{A'}), \quad \tilde{f} = \ln((z_{A'} \pi^{A'}) x^{1'} - \pi^{1'})$$

and

$$L_1(f) = L_1(\tilde{f}) = 0, \quad L_0(f) = L_0(\tilde{f}) = \pi^{A'} z_{A'}.$$

Therefore $\mathcal{A}_{1A'} = 0$, $\mathcal{A}_{0A'} = z_{A'}$ which gives $\mathcal{A} = z_{A'} dx^{A'}$, and $d\mathcal{A}$ is indeed the ASD para-Kähler structure Ω .

□

5.4.5 Factoring the model to Einstein–Weyl

As stated above, we expect distinct Einstein–Weyl structures if we factor M by conformal Killing vectors which are not conjugate with respect to an isometry [60]. We can thus classify the Einstein–Weyl structures obtainable from the model by first classifying its symmetries up to conjugation.

Proposition 5.4.5. *The non-trivial Einstein–Weyl structures obtainable from the ASD Einstein metric (5.30) by the Jones–Tod correspondence consist of a two parameter family, and two additional cases which do not belong to this family.*

Proof. Since we have an isomorphism between the Lie algebra of projective vector fields on $(N, [\nabla])$ and the Lie algebra of Killing vectors on (M, g) , the problem of classifying the symmetries of (5.30) is reduced to a classification of the infinitesimal projective symmetries of \mathbb{RP}^2 , i.e. the near-identity elements of $SL(3, \mathbb{R})$, up to conjugation.

Non-singular complex matrices are determined up to similarity by their Jordan normal form (JNF). While real matrices do not have such a canonical form, all of the information they contain is determined (up to similarity) by the JNF that they would have if they were considered as complex matrices. Thus we can still discuss the JNF of a real matrix, even if it cannot always be obtained from the real matrix by a real similarity transformation. The possible non-trivial Jordan normal forms of matrices in

$SL(3, \mathbb{R})$ are shown below.

$$\begin{pmatrix} l_1 & 0 & 0 \\ 0 & l_2 & 0 \\ 0 & 0 & 1/l_1 l_2 \end{pmatrix} \quad \begin{pmatrix} l & 0 & 0 \\ 0 & l & 0 \\ 0 & 0 & 1/l^2 \end{pmatrix} \quad \begin{pmatrix} l & 1 & 0 \\ 0 & l & 0 \\ 0 & 0 & 1/l^2 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

It is possible that two matrices in $SL(3, \mathbb{R})$ with the same JNF may be related by a complex similarity transformation, and thus not conjugate in $SL(3, \mathbb{R})$. However, if the JNF is a real matrix, then the required similarity transformation just consists of the eigenvectors and generalised eigenvectors of the matrix, which must also be real since they are defined by real linear simultaneous equations. This means we only have to worry about matrices with complex eigenvalues, and since these occur in complex conjugate pairs, they will only be a problem when we have three distinct eigenvalues.

In this case, we can always make a real similarity transformation such that the matrix is block diagonal, with the real eigenvalue in the bottom right. Then we have limited choice from the 2×2 matrix in the top left. Let us parametrise such a 2×2 matrix by $a, b, c, d \in \mathbb{R}$ as follows:

$$\begin{pmatrix} 1 + a\epsilon & b\epsilon \\ c\epsilon & 1 + d\epsilon \end{pmatrix}.$$

This has characteristic polynomial

$$\chi(l) = l^2 - (2 + \epsilon(a + d))l + 1 + (a + d)\epsilon + (ad - bc)\epsilon^2.$$

Evidently the important degrees of freedom are $a + d$ and $ad - bc$, so we can use these to encode every near-identity element of the class with three distinct eigenvalues. The bottom-right entry will be determined by our choice of $a + d$ and $ad - bc$.

Taking a projective vector field on \mathbb{RP}^2 , we can find the corresponding Killing vector of (5.30) using (3.25), and factor to Einstein–Weyl using (5.13). We find by explicit calculation that vector fields arising from the second and fourth JNFs above give trivial Einstein–Weyl structures, so restricting to the non-trivial cases we have a two parameter family of Einstein–Weyl structures coming from the first class, and two additional Einstein–Weyl structures coming from the third and fifth, as claimed.

□

5.4.6 An example of the mini-twistor correspondence

Below we investigate a one parameter subfamily of the two parameter family. We use the holomorphic vector field on the twistor space F_{12} (see Section 5.4.2) corresponding to the chosen symmetry, and reconstruct the conformal structure $[h]$ on \mathcal{W} using minitwistor curves (in the sense of [39]) on the space of orbits. Take $a \in \mathbb{R}$ and

$$K = P^1 \frac{\partial}{\partial P^1} - L_1 \frac{\partial}{\partial L_1} + a P^2 \frac{\partial}{\partial P^2} - a L_2 \frac{\partial}{\partial L_2}, \quad (5.41)$$

In order to preserve the relations

$$p \cdot L = 0, \quad P \cdot l = 0, \quad p \cdot l = 0,$$

the corresponding holomorphic action on (p, l) must be $p \mapsto gp$, $l \mapsto lg^{-1}$, thus the holomorphic vector field $K_{\mathcal{T}}$ on F_{12} is

$$K_{\mathcal{T}} = p^1 \frac{\partial}{\partial p^1} - l_1 \frac{\partial}{\partial l_1} + a p^2 \frac{\partial}{\partial p^2} - a l_2 \frac{\partial}{\partial l_2}.$$

In order to factor F_{12} by this vector field, we must find invariant minitwistor coordinates (Q, R) . In addition to satisfying $K_{\mathcal{T}}(Q) = K_{\mathcal{T}}(R) = 0$, they must be homogeneous of degree zero in (P, L) . We choose

$$Q = \frac{p^1 l_1}{p^2 l_2}, \quad R = \frac{(l_1)^a}{l_2 (l_3)^{a-1}}.$$

Substituting in our parametrisation (5.37) and using the freedom to perform a Möbius transformation on π , we obtain

$$\begin{aligned} Q &= \frac{(\lambda z - r - 1)\lambda}{w\lambda + \lambda - \frac{rw}{z}} \\ R &= \lambda^a \left(-\lambda - \frac{w}{z} \right)^{1-a}, \end{aligned} \quad (5.42)$$

where we have defined $\lambda = \pi_0 / \pi_1$, and the Einstein–Weyl coordinates

$$r = xp, \quad w = yq, \quad z = x^a q.$$

Note these are invariants of the Killing vector (5.41).

Next we wish to use these minitwistor curves to reconstruct the conformal structure of the Einstein–Weyl space. In doing so we follow [60]. The tangent vector field to a

fixed curve is given by

$$\xi_T = \frac{\partial Q}{\partial \lambda} \frac{\partial}{\partial Q} + \frac{\partial R}{\partial \lambda} \frac{\partial}{\partial R},$$

Hence we can write the normal vector field as

$$\begin{aligned} \xi_N &= dQ \frac{\partial}{\partial Q} + dR \frac{\partial}{\partial R} \bmod \xi_T \\ &= \left(\frac{\partial R}{\partial \lambda} \right)^{-1} \left(dQ \frac{\partial R}{\partial \lambda} - dR \frac{\partial Q}{\partial \lambda} \right) \frac{\partial}{\partial Q}, \end{aligned}$$

where

$$dQ = \frac{\partial Q}{\partial r} dr + \frac{\partial Q}{\partial w} dw + \frac{\partial Q}{\partial z} dz$$

and similarly for dR . Calculating ξ_N using (5.42), we find

$$\xi_N \propto (\eta_1 \lambda^2 + \eta_2 \lambda + \eta_3) \frac{\partial}{\partial Q},$$

where

$$\begin{aligned} \eta_1 &= z^2(w+1)dz - z^3dw, \\ \eta_2 &= -2zrwdz + z^2(a+2r)dw - z^2dr, \\ \eta_3 &= rw(1+r)dz - zr(1+r)dw - azwdr. \end{aligned}$$

The discriminant of this quadratic in λ then gives a representative $h \in [h]$ of our conformal structure:

$$\begin{aligned} h = 4(r^2w + rw^2 + rw)dz^2 - 4zw(a(w+1) + r)dzdr + 4zr(r - aw + 2w + 1)dzdw \\ - z^2dr^2 + 2z^2(2aw + a + 2r)dwdr - z^2(a^2 + 4r(a-1))dw^2. \end{aligned} \tag{5.43}$$

This is the same conformal structure that we obtain by Jones–Tod factorisation of the metric (5.30) by (5.41) using the formula (5.13).

Chapter 6

Para- c -projective compactification of M

In [18] the concept of c -projective compactification was defined. It is based on almost c -projective geometry [15], an analogue of projective geometry defined for almost complex manifolds, i.e., even-dimensional manifolds M carrying a smooth endomorphism J of TM which satisfies $J^2 = -Id$. In c -projective geometry, the equivalence class of torsion-free connections is replaced by an equivalence class of connections which are adapted to the almost complex structure J in a natural way. In this chapter we discuss a notion of compactification which is modified to the “*para*” case, i.e. where the endomorphism J squares to Id rather than $-Id$. We show that the natural almost para-complex structure J on any manifold M arising in the projective to Einstein correspondence admits a type of compactification which we call *para-c*-projective. The content of this chapter is based on material appearing in [29]. It was undertaken in collaboration with Maciej Dunajski and Rod Gover.

6.1 Background and definitions

The purpose of this section is to introduce the definitions which are required to state the main results of [18].

6.1.1 Almost (para-)complex geometry

Definition 6.1.1. *The Nijenhuis tensor of an endomorphism J of TM is defined by*

$$\mathcal{N}(\xi_1, \xi_2) := [\xi_1, \xi_2] - [J\xi_1, J\xi_2] + J([J\xi_1, \xi_2] + [\xi_1, J\xi_2]), \quad (6.1)$$

where ξ_1, ξ_2 are vector fields on M and $[\cdot, \cdot]$ denotes the Lie bracket of vector fields. This is equivalent to

$$\mathcal{N}_{bc}^a = J^d_{[b} \partial_{d]} J^a_{c]} - J^d_{[b} \partial_{c]} J^a_{d]}. \quad (6.2)$$

Let M be a complex manifold of (complex) dimension n , in the sense of having complex coordinates and complex transition functions. Then multiplication of the coordinates by i defines an endomorphism J of TM which squares to $-Id$, so complex manifolds are a subset of almost complex manifolds. In this case, J has eigenvalues $\pm i$, and the corresponding splitting of TM into eigen-bundles is Frobenius integrable. The Newlander–Nirenberg theorem describes complex manifolds in terms of the Nijenhuis tensor (6.1) of J .

Theorem 6.1.2 ([59]). *An almost complex manifold (M, J) is a complex manifold if and only if the Nijenhuis tensor of J vanishes. In this case, we call the almost complex structure J integrable.*

As discussed in Chapter 3, an endomorphism J which squares to Id defines an analogous splitting of the tangent bundle into sub-bundles with eigenvalues ± 1 , and this splitting is also Frobenius integrable if and only if the Nijenhuis tensor of J vanishes. We thus call an almost para-complex structure J with vanishing Nijenhuis tensor a para-complex structure, and say that in this case J is integrable. In all the definitions below, the word *almost* can be removed if the (para-)complex structure J is integrable.

Definition 6.1.3. *A (para-)Hermitian metric on an almost (para-)complex manifold (M, J) is a metric g satisfying*

$$g(J\cdot, J\cdot) = \pm g(\cdot, \cdot),$$

where the minus sign corresponds to the “para” case. The triple (M, J, g) then defines an almost (para-)Hermitian manifold.

Note that every (para-)Hermitian manifold has a naturally defined two-form $\Omega(\cdot, \cdot) = g(\cdot, J\cdot)$ which is (para-)Hermitian in the sense that

$$\Omega(J\cdot, J\cdot) = \pm \Omega(\cdot, \cdot),$$

and can alternatively be specified as (M, J, Ω) or (M, g, Ω) . An almost (para-)Kähler manifold (M, J, g) is a (para-)Hermitian manifold whose associated two-form is closed, meaning M carries compatible complex, pseudo-Riemannian and symplectic structures.

The manifolds M arising in the projective to Einstein correspondence are almost para-Kähler, and para-Kähler when the underlying projective structure is flat [28].

6.1.2 Almost (para-)CR structures and contact distributions

Definition 6.1.4. *An almost (para-)CR structure $(\mathcal{Z}, \mathbb{H}, J)$ on a manifold \mathcal{Z} is a subbundle $\mathbb{H} \subset T\mathcal{Z}$ of the tangent bundle together with a fibre-preserving endomorphism $J : \mathbb{H} \rightarrow \mathbb{H}$ which satisfies $J^2 = Id$ or $J^2 = -Id$ depending on whether or not we are talking about the “para” case.*

We will be interested in the case where \mathbb{H} is a hyperplane distribution on \mathcal{Z} ; then $(\mathcal{Z}, \mathbb{H}, J)$ is called an almost (para-)CR structure of *hypersurface type*. An almost (para-)complex structure (M, J) of dimension $2n$ defines an almost (para-)CR structure of hypersurface type on any hypersurface $\mathcal{Z} \subset M$ given by the restriction of J to the hyperplane distribution $\mathbb{H} := T\mathcal{Z} \cap J(T\mathcal{Z})$ on \mathcal{Z} . Note that this distribution must have dimension $2n - 2$. An almost (para-)CR structure is a (para-)CR structure if and only if the splitting of \mathbb{H} into eigen-bundles induced by J is Frobenius integrable.

We can define the notion of non-degeneracy for an almost (para-)CR structure as follows. The Lie bracket of vector fields induces an antisymmetric \mathbb{R} -bilinear operator $\Gamma(\mathbb{H}) \times \Gamma(\mathbb{H}) \rightarrow \Gamma(T\mathcal{Z}/\mathbb{H})$ which in fact is also bilinear over smooth functions on \mathcal{Z} . This means it is induced by a bundle map $\mathcal{L} : \mathbb{H} \times \mathbb{H} \rightarrow T\mathcal{Z}/\mathbb{H}$ which is called the *Levi bracket*. Since it takes values in a line bundle it can be thought of as an antisymmetric bilinear form called the *Levi form*. Degeneracy (or not) of the almost (para-)CR structure is defined as degeneracy (or not) of the Levi form. Note that the Levi form also defines a *symmetric* bilinear form $h_{\mathbb{H}}(\cdot, \cdot) = \mathcal{L}(\cdot, J\cdot)$ as long as \mathcal{L} is (para-)Hermitian with respect to J , and that this symmetric bilinear form is non-degenerate if and only if \mathcal{L} is.

Definition 6.1.5. *A contact structure on a manifold \mathcal{Z} of dimension $2n - 1$ is a hyperplane distribution $\mathbb{H} \subset T\mathcal{Z}$ specified as the kernel of a one-form β on \mathcal{Z} which satisfies the complete non-integrability condition*

$$\beta \wedge \underbrace{(d\beta \wedge \cdots \wedge d\beta)}_{n-1 \text{ times}} \neq 0. \quad (6.3)$$

The complete non-integrability condition can be thought of as the opposite of Frobenius integrability of the hyperplane distribution, see for example [1].

6.1.3 Connections and (para-) c -projective equivalence

Definition 6.1.6. *A connection on an almost (para-)complex manifold (M, J) is called complex if it preserves J .*

Note that, in contrast to a metric connection, it is not always possible to define a complex connection which is torsion-free. In fact, this is possible if and only if the Nijenhuis tensor (6.1) of J vanishes. However, one can always define a complex connection whose torsion is equal to the Nijenhuis tensor of J up to a constant multiplicative factor [15]. Such connections are called *minimal*.

Definition 6.1.7. *Two affine connections ∇ and $\bar{\nabla}$ on an almost (para-)complex manifold (M, J) are called (para-) c -projectively equivalent if there is a one-form Υ_a on M such that their components Γ_{bc}^a and $\bar{\Gamma}_{bc}^a$ are related by*

$$\bar{\Gamma}_{bc}^a - \Gamma_{bc}^a = \delta_b^a \Upsilon_c + \delta_c^a \Upsilon_b \pm (\Upsilon_d J^d_b J^a_c + \Upsilon_d J^d_c J^a_b), \quad (6.4)$$

where the plus corresponds to the case $J^2 = Id$ and the minus corresponds to the case $J^2 = -Id$.

Note that the para- c -projective change of connection differs from the c -projective case in the signs of some of the terms, to account for the fact that J squares to the Id rather than $-Id$. It is easy to show that if ∇ is complex then so is $\bar{\nabla}$, and the index symmetry of the right hand side of (6.4) means that if ∇ is minimal then so is $\bar{\nabla}$. An almost (para-) c -projective structure on a manifold M comprises an almost (para-)complex structure J and a (para-) c -projective equivalence class $[\nabla]$ of complex minimal connections.

We note here for later use that c -projective geometry in $2n$ dimensions can be expressed as a Cartan geometry. The model Lie group quotient is G/S , where

$$G = \{g \in SL(2n+2, \mathbb{R}) : g\mathbb{J} = \mathbb{J}g\},$$

and \mathbb{J} is an endomorphism of \mathbb{R}^{2n+2} which squares to $-Id$. This can be identified with $SL(n+1, \mathbb{C})$. The subgroup S is the stabiliser subgroup of a complex line in \mathbb{C}^{n+1} , or equivalently a real plane in \mathbb{R}^{2n+2} . Since a complex line in \mathbb{C}^{n+1} projects to a point in \mathbb{CP}^n , \mathbb{CP}^n can be realised as G/S . More details can be found in [15]. Although the para- c -projective case has not been studied in detail, we expect the construction to be analogous, with \mathbb{J} instead squaring to the identity on \mathbb{R}^{2n+2} .

6.1.4 Para- c -projective compactification

We now specialise to the “para” case, where $J^2 = Id$. Note that all the corresponding results for $J^2 = -Id$ can be found in [18].

Definition 6.1.8. *Let (M, J) be an almost para-complex manifold, and let ∇ be a complex minimal connection. The structure (M, J) admits a para- c -projective compactification to a manifold with boundary $\overline{M} = M \cup \partial M$ if there exists a function $T : \overline{M} \rightarrow \mathbb{R}$ such that the zero locus $\mathcal{Z}(T)$ is the boundary $\partial M \subset \overline{M}$, the differential dT does not vanish on ∂M , and the connection $\overline{\nabla}$, related to ∇ by (6.4) with $\Upsilon = dT/(2T)$, extends to \overline{M} .*

It follows easily from this definition that the endomorphism J on M naturally extends to all of \overline{M} by parallel transport with respect to $\overline{\nabla}$. It thus defines an almost para-CR structure on the hyperplane distribution \mathbb{H} defined by $\mathbb{H}_m := T_m \partial M \cap J(T_m \partial M)$ for all $m \in \partial M$. It can be shown that this almost para-CR structure is non-degenerate if and only if for any local defining function T the one-form $\beta = dT \circ J$, whose restriction to ∂M has kernel \mathbb{H} , satisfies the complete non-integrability condition (6.3) making \mathbb{H} a contact distribution on ∂M .

To see this, first note that $\beta(\xi) = 0 \forall \xi \in \Gamma(\mathbb{H})$ implies $d\beta(\cdot, \cdot) = -\beta([\cdot, \cdot])$, so the restriction of $d\beta$ to $\mathbb{H} \times \mathbb{H}$ represents the Levi form \mathcal{L} . This means that the almost para-CR structure on ∂M is non-degenerate if and only if the restriction of $d\beta(\xi, \cdot)$ to \mathbb{H} is non-zero for all non-zero $\xi \in \Gamma(\mathbb{H})$. But this is equivalent to the non-integrability condition (6.3).

Another result of lemma 5 of [18] is that $d\beta$ is Hermitian on ∂M if and only if the Nijenhuis tensor (6.1) of J takes so-called *asymptotically tangential values*. This is equivalent to the following statement in index notation:

$$\left(\mathcal{N}^a_{bc} \nabla_a T \right) \Big|_{T=0} = 0. \quad (6.5)$$

Note in particular that Hermiticity of $d\beta$ on ∂M implies Hermiticity of $d\beta$ on \mathbb{H} , and hence the existence of a non-degenerate metric $h_{\mathbb{H}}(\cdot, \cdot) = d\beta(\cdot, J\cdot)|_{\mathbb{H}}$ on \mathbb{H} . Both of these facts also apply in the “para” case.

Although c -projective compactification is defined for any almost complex manifold, the definition can be applied to pseudo-Riemannian metrics g which are Hermitian with respect to the almost complex structure so long as there exists a connection which preserves both g and J and has minimal torsion. Such Hermitian metrics are said to be *admissible*. Note that such a connection, if it exists, is uniquely defined, since the

conditions that it be complex and minimal determine its torsion. It is thus given by the Levi–Civita connection of g plus a constant multiple of the Nijenhuis tensor (6.1) of J .

The first main result of [18] is Theorem 8 in this reference, which gives a local form for an admissible Hermitian metric which is sufficient for the corresponding c -projective structure to be c -projectively compact. The theorem is stated below, adapted to the para- c -projective case. The proof can be obtained by a trivial adaptation of the arguments in [18], and so further details may be obtained from that source.

Theorem 6.1.9 ([18]). *Let \overline{M} be a smooth manifold with boundary ∂M and interior M . Let J be an almost para-complex structure on \overline{M} , such that ∂M is non-degenerate and the Nijenhuis tensor \mathcal{N} of J has asymptotically tangential values. Let g be an admissible pseudo-Riemannian Hermitian metric on M . For a local defining function T for the boundary defined on an open subset $\mathcal{U} \subset \overline{M}$, put $\beta = dT \circ J$ and, given a non-zero real constant C , define a Hermitian tensor field $h_{T,C}$ on $\mathcal{U} \cap M$ by*

$$h_{T,C} := Tg + \frac{C}{T}(dT^2 - \beta^2).$$

Suppose that for each $x \in \partial M$ there is an open neighbourhood \mathcal{U} of x in \overline{M} , a local defining function T defined on \mathcal{U} , and a non-zero constant C such that

- $h_{T,C}$ admits a smooth extension to all of \mathcal{U}
- for all vector fields ξ_1, ξ_2 on \mathcal{U} with $dT(\xi_2) = \beta(\xi_2) = 0$, the function $h_{T,C}(\xi_1, J\xi_2)$ approaches $Cd\beta(\xi_1, \xi_2)$ at the boundary.

Then g is c -projectively compact.

Note that the statement in Theorem 6.1.9 does not depend on the choice of T . Different choices of T result in rescalings of the one-form β on the boundary by a nowhere vanishing function.

6.2 Compactifying the Dunajski–Mettler Class

In order to construct the para- c -projective compactification of the manifolds M arising in the projective to Einstein correspondence, we will need to understand them from a tractor perspective. This is the goal of the following subsection.

6.2.1 Tractor construction of M

In Section 3.2.2 it was shown that the projectivised cotractor bundle of N is stratified by the canonical density $\tau = V \lrcorner W$, where V is the pull back of the canonical tractor along $\pi_{\mathcal{T}} : \mathcal{T}^* \rightarrow N$ and W is the tautological section of $\pi_{\mathcal{T}}^*(\mathcal{T}^*)$. It is easily verified that the zero locus $\mathcal{Z}(\tau)$ of τ is a smoothly embedded hypersurface in $\mathcal{M} := \mathbb{P}(\mathcal{T}^*)$. In the following theorem, we show that $\mathcal{M} \setminus \mathcal{Z}(\tau)$ can be identified with M .

Theorem 6.2.1. [29] *There is a metric g and two-form Ω on $\mathcal{M} \setminus \mathcal{Z}(\tau)$ determined by the canonical pairing of the horizontal and vertical subspaces of $T(\mathcal{T}^*)$. The pair (g, Ω) agrees with (3.19).*

Proof. Considering first the total space \mathcal{T}^* and then its tangent bundle, note that there is an exact sequence

$$0 \rightarrow \pi_{\mathcal{T}}^* \mathcal{T}^* \rightarrow T(\mathcal{T}^*) \rightarrow \pi_{\mathcal{T}}^* TN \rightarrow 0, \quad (6.6)$$

where we have identified $\pi_{\mathcal{T}}^* \mathcal{T}^*$ as the vertical sub-bundle of $T(\mathcal{T}^*)$. The tractor connection on the vector bundle $\mathcal{T}^* \rightarrow N$ is equivalent to a splitting of this sequence, identifying $\pi_{\mathcal{T}}^* TN$ with a distinguished sub-bundle of horizontal subspaces in $T(\mathcal{T}^*)$ so that we have

$$T(\mathcal{T}^*) = \pi_{\mathcal{T}}^* TN \oplus \pi_{\mathcal{T}}^* \mathcal{T}^*. \quad (6.7)$$

We move now to the total space of $\mathcal{M} := \mathbb{P}(\mathcal{T}^*)$, and we note that again the tractor (equivalently, Cartan) connection determines a splitting of the tangent bundle $T(\mathbb{P}\mathcal{T}^*)$ in which the second term of the display (6.7) is replaced by a quotient of $\pi_{\mathcal{T}}^* \mathcal{T}^*(0, 1)$ [19]. Indeed, if we work at a point $m \in \mathbb{P}(\mathcal{T}^*)$, observe that $\pi_{\mathcal{T}}^* \mathcal{T}^*(0, 1)$ has a filtration

$$0 \longrightarrow \mathcal{E}(0, 0)_m \xrightarrow{W_m} \pi_{\mathcal{M}}^* \mathcal{T}^*(0, 1)|_m \longrightarrow \pi_{\mathcal{M}}^* \mathcal{T}^*(0, 1)|_m / \langle W_m \rangle \longrightarrow 0 \quad (6.8)$$

where, as usual, W is the canonical section. But away from $\mathcal{Z}(\tau)$, we have that W canonically splits the appropriately re-weighted pull back of the sequence (3.14)

$$0 \longrightarrow \pi_{\mathcal{M}}^* T^* N(1, 1) \longrightarrow \pi_{\mathcal{M}}^* \mathcal{T}^*(0, 1) \xrightarrow{V/\tau} \mathcal{E}(0, 0) \longrightarrow 0.$$

This identifies the quotient in (6.8), and thus we have canonically

$$T(\mathbb{P}(\mathcal{T}^*) \setminus \mathcal{Z}(\tau)) = \pi_{\mathcal{M}}^* TN \oplus \pi_{\mathcal{M}}^* T^* N(1, 1).$$

It follows that on $M := \mathcal{M} \setminus \mathcal{Z}(\tau)$ there is canonically a metric \mathbf{g} and symplectic form Ω taking values in $\mathcal{E}(1, 1)$, given by

$$\begin{aligned}\mathbf{g}(w_1, w_2) &= \frac{1}{2} \left(\Pi_H(w_1) \lrcorner \Pi_V(w_2) + \Pi_H(w_2) \lrcorner \Pi_V(w_1) \right) \quad \text{and} \\ \Omega(w_1, w_2) &= \frac{1}{2} \left(\Pi_H(w_1) \lrcorner \Pi_V(w_2) - \Pi_H(w_2) \lrcorner \Pi_V(w_1) \right)\end{aligned}$$

where

$$\Pi_H : TM \rightarrow \pi_{\mathcal{M}}^* TN \quad \text{and} \quad \Pi_V : TM \rightarrow \pi_{\mathcal{M}}^* T^* N(1, 1)$$

are the projections. Then we obtain the metric and symplectic form by

$$g := \frac{1}{\tau} \mathbf{g} \quad \text{and} \quad \Omega := \frac{1}{\tau} \Omega. \quad (6.9)$$

What remains to be done, is to show that (6.9) agrees with the form obtained in [28] once a trivialisation of $\mathcal{T}^* \rightarrow N$ has been chosen.

Let $x \in N$ and let $\mathcal{U} \subset N$ be an open neighbourhood of x with local coordinates (x^1, \dots, x^n) such that $T_x N = \text{span}(\partial/\partial x^1, \dots, \partial/\partial x^n)$. The connection (3.15) gives a splitting of $T(\mathcal{T}^*)$ into the horizontal and vertical sub-bundles

$$T(\mathcal{T}^*) = \mathcal{H}(\mathcal{T}^*) \oplus \mathcal{V}(\mathcal{T}^*),$$

as in (6.7). To obtain the explicit form of this splitting, let σ_α , $\alpha = 0, 1, \dots, n$ be components of a local section of \mathcal{T}^* in the trivialisation over \mathcal{U} . Then

$$\nabla^{\mathcal{T}} \sigma_\beta = d\sigma_\beta - \gamma_\beta^\alpha \sigma_\alpha,$$

where $\gamma_\alpha^\beta = \gamma_{i\alpha}^\beta dx^i$, and the components of the co-tractor connection $\gamma_{i\alpha}^\beta$ are given in terms of the connection ∇ on N , and its Schouten tensor, and can be read-off from (3.15):

$$\gamma_{i0}^0 = 0, \quad \gamma_{i0}^j = \delta_i^j, \quad \gamma_{ij}^k = \Gamma_{ij}^k, \quad \gamma_{ij}^0 = -P_{ij}.$$

In terms of these components we can write

$$\begin{aligned}\mathcal{H}(\mathcal{T}^*) &= \text{span} \left(\frac{\partial}{\partial x^i} + \gamma_{i\alpha}^\beta \sigma_\beta \frac{\partial}{\partial \sigma_\alpha}, i = 1, \dots, n \right), \\ \mathcal{V}(\mathcal{T}^*) &= \text{span} \left(\frac{\partial}{\partial \sigma_\alpha}, \alpha = 0, 1, \dots, n \right).\end{aligned}$$

Setting $\zeta_i = \sigma_i/\sigma_0$, where $\tau = \sigma_0 \neq 0$ on the complement of $\mathcal{Z}(\tau)$, we can compute the push forwards of these subspaces to $\mathbb{P}(\mathcal{T}^*) \setminus \mathcal{Z}(\tau)$:

$$\kappa_* \mathcal{H}(\mathcal{T}^*) = \text{span} \left(h_i \equiv \frac{\partial}{\partial x^i} - (P_{ij} + \zeta_i \zeta_j - \Gamma_{ij}^k \zeta_k) \frac{\partial}{\partial \zeta_j} \right), \quad \kappa_* \mathcal{V}(\mathcal{T}^*) = \text{span} \left(v^i \equiv \frac{\partial}{\partial \zeta_i} \right).$$

The non-zero components of the metric (6.9) are given by

$$g(v^i, h_j) = \delta^i_j.$$

This is identical to the form appearing in [28].

□

Remark 6.2.2. Note that the shift (3.20) in the fibre coordinates ζ_i corresponding to a change of projective connection can be motivated from the change (3.13) in the splitting of \mathcal{T}^* and the definitions of σ_i and ζ_i .

Remark 6.2.3. We can also understand $\mathbb{P}(\mathcal{T}^*) \setminus \mathcal{Z}(\tau)$ as an affine bundle modelled on T^*N . Given a connection in the projective class and hence a decomposition (3.11), there is a smooth fibre bundle isomorphism

$$\kappa_A : T^*N \rightarrow \mathbb{P}(\mathcal{T}^*) \setminus \mathcal{Z}(\tau). \quad (6.10)$$

given by

$$T_x^*N \ni \zeta_i \mapsto [(1, \zeta_i)] = [(\tau, \tau \zeta_i)] \in \mathbb{P}(\mathcal{T}_x^*) \setminus \mathcal{Z}(\tau). \quad (6.11)$$

6.2.2 The compactification theorem

As noted in Chapter 3, in the model case where $N = \mathbb{RP}^n$ and $[\nabla]$ is projectively flat, the manifold $M = SL(n+1, \mathbb{R})/GL(n, \mathbb{R})$ can be identified with the projectivisation of $\mathbb{R}^{n+1} \times \mathbb{R}_{n+1} \setminus \mathcal{Z}$, where \mathcal{Z} denotes the set of incident pairs (point, hyperplane). The compactification procedure described in the Theorem 6.2.4 below will, for the model, attach these incident pairs back to M , and more generally (in case of a curved projective structure $(N, [\nabla])$) will attach the zero locus of τ back into $\mathbb{P}(\mathcal{T}^*)$. The boundary $\partial M \cong \mathcal{Z}(\tau)$ from definition 6.1.8 will play the role of a submanifold separating two open sets in $\mathbb{P}(\mathcal{T}^*)$ which have $\tau > 0$ and $\tau < 0$ respectively. The method of the proof will be to show that near the boundary $\mathcal{Z}(\tau) = 0$ of \overline{M} the metric (3.19) can be put in the local normal form of Theorem 6.1.9.

Theorem 6.2.4. *The Einstein almost para-Kähler structure (M, g, Ω) given by (3.19) admits a para- c -projective compactification \overline{M} . The $(2n - 1)$ -dimensional boundary $\partial M \cong \mathcal{Z}(\tau)$ of \overline{M} carries a contact structure together with a conformal structure and an almost para-CR structure defined on the contact distribution.*

Proof. In the proof below we shall explicitly construct the boundary ∂M together with the contact structure and the associated conformal structure on the contact distribution. We shall first deal with the model case (3.27), and then explain how the addition of non-vanishing projective curvature modifies the compactification.

Now consider an open set $\mathcal{U} \subset M$ given by $\zeta_i x^i > 0$, and define the function T on \mathcal{U} by

$$T = \frac{1}{\zeta_i x^i}. \quad (6.12)$$

We shall attach a boundary $\partial\mathcal{U}$ to the open set \mathcal{U} such that T extends to a function \overline{T} on $\mathcal{U} \cup \partial\mathcal{U}$, and \overline{T} is the defining function for this boundary. We then investigate the geometry on M in the limit $T \rightarrow 0$. It is clear from above that the zero locus of \overline{T} will be contained in the zero locus $\mathcal{Z}(\tau)$ of τ , and therefore belongs to the boundary of \overline{M} . We will use \overline{T} as a defining function for \overline{M} in an open set $\overline{\mathcal{U}} \subset \overline{M}$. The strategy of the proof is to extend T to a coordinate system on \mathcal{U} , such that near the boundary the metric g takes a form as in Theorem 6.1.9.

First define $\beta \in \Lambda^1(\overline{M})$ by

$$\xi \lrcorner \beta = J(\xi) \lrcorner dT, \quad \text{or equivalently} \quad \beta_a = \Omega_{ac} g^{bc} (\mathbf{g} \nabla_b T), \quad a, b, c = 1, \dots, 2n \quad (6.13)$$

where J is the para-complex structure of (g, Ω) and ξ is a vector field on M . Using (3.27) this gives

$$\beta = 2T(1 - T)\zeta_i dx^i - dT.$$

We need n open sets $\mathcal{U}_1, \dots, \mathcal{U}_n$ such that $\zeta_k \neq 0$ on \mathcal{U}_k to cover the zero locus of T . Here we chose $k = n$, and use a coordinate system given by

$$(T, Z_1, \dots, Z_{n-1}, X^1, \dots, X^{n-1}, Y),$$

where T is given by (6.12) and

$$Z_A = \frac{\zeta_A}{\zeta_n}, \quad X^A = x^A, \quad Y = x^n, \quad \text{where} \quad A = 1, \dots, n-1.$$

We compute

$$\beta = 2(1-T) \frac{dY + Z_A dX^A}{K} - dT, \quad \zeta_n = \frac{1}{KT}, \quad \text{where} \quad K \equiv Y + Z_A X^A,$$

and substitute

$$\zeta_i dx^i = \frac{1}{KT} (dY + Z_A dX^A)$$

into (3.19). This gives

$$g = \frac{\beta^2 - dT^2}{4T^2} + \frac{1}{T} h_T, \quad (6.14)$$

where

$$h_T = \frac{1}{4(1-T)} (\beta^2 - dT^2) + \frac{1}{K} \left(dZ_A \odot dX^A - \frac{1}{2(1-T)} X^A dZ_A \odot (\beta + dT) \right)$$

is regular at the boundary $T = 0$. This is in agreement with the asymptotic form in Theorem 6.1.9 (see [18] for further details).

The restriction of h_T to ∂M gives a metric on the distribution $\mathbb{H} = \text{Ker}(\beta|_{T=0})$

$$\begin{aligned} \beta|_{T=0} &= 2 \frac{dY + Z_A dX^A}{Y + Z_A X^A}, \\ h_T|_{T=0} &= \frac{1}{4} (\beta|_{T=0})^2 + \frac{1}{2(Y + Z_A X^A)} (2dZ_A \odot dX^A - X^A dZ_A \odot (\beta|_{T=0})). \end{aligned} \quad (6.15)$$

Note that T is only defined up to multiplication by a positive function. Changing the defining function in this way results in a conformal rescaling of $\beta|_{T=0}$, thus the metric on the contact distribution is also defined up to an overall conformal scale. We shall choose the scale so that the contact form is given by $\beta_0 \equiv K\beta|_{T=0}$ on $T(\partial M)$, with the metric on \mathbb{H} given by

$$h_{\mathbb{H}} = dZ_A \odot dX^A. \quad (6.16)$$

We now move on to deal with the curved case where the metric on M is given by (3.19). The coordinate system (T, Z_A, X^A, Y) is as above, and the one-form β in (6.13) is given by

$$\beta = 2T(1-T)\zeta_i dx^i - dT + 2T^2(P_{ij} - \Gamma_{ij}^k \zeta_k) x^i dx^j,$$

or in the (T, Z_A, X^A, Y) coordinates,

$$\begin{aligned}\beta = & 2(1-T) \frac{Z_A dX^A + dY}{K} - dT \\ & + 2T^2 \left[\left(P_{AB} - \frac{\Gamma_{AB}^C Z_C + \Gamma_{AB}^n}{TK} \right) X^A dX^B + \left(P_{nB} - \frac{\Gamma_{nB}^C Z_C + \Gamma_{nB}^n}{TK} \right) Y dX^B \right. \\ & \left. + \left(P_{An} - \frac{\Gamma_{An}^C Z_C + \Gamma_{An}^n}{TK} \right) X^A dY + \left(P_{nn} - \frac{\Gamma_{nn}^C Z_C + \Gamma_{nn}^n}{TK} \right) Y dY \right].\end{aligned}$$

Guided by the formula (6.14) we define

$$h_T = Tg - \frac{1}{4T}(\beta^2 - dT^2),$$

which we find to be

$$\begin{aligned}h_T = & \frac{1}{4(1-T)}(\beta^2 - dT^2) + \frac{1}{K} \left(dZ_A \odot dX^A - \frac{1}{2(1-T)} X^A dZ_A \odot (\beta + dT) \right) \\ & - \frac{1}{K} \left((\Gamma_{AB}^C Z_C + \Gamma_{AB}^n) dX^A \odot dX^B + (\Gamma_{nn}^C Z_C + \Gamma_{nn}^n) dY \odot dY \right. \\ & \left. + 2(\Gamma_{An}^C Z_C + \Gamma_{An}^n) dX^A \odot dY \right) \\ & + T(P_{AB} dX^A \odot dX^B + 2P_{An} dX^A \odot dY + P_{nn} dY \odot dY).\end{aligned}$$

This is smooth as $T \rightarrow 0$.

Restricting h_T to $T = 0$ yields a metric which differs from (6.15) by the curved contribution given by the components of the connection, but not the Schouten tensor. Substituting $dY = K\beta|_{T=0}/2 - Z_A dX^A$, disregarding the terms involving $\beta|_{T=0}$ in h_T , and conformally rescaling by K yields the metric

$$\begin{aligned}h_{\mathbb{H}} &= (dZ_A - \Xi_{AB} dX^B) \odot dX^A, \quad \text{where} \\ \Xi_{AB} &= \Gamma_{AB}^C Z_C + \Gamma_{AB}^n + (\Gamma_{nn}^C Z_C + \Gamma_{nn}^n) Z_A Z_B - 2(\Gamma_{An}^C Z_C + \Gamma_{An}^n) Z_B\end{aligned}\tag{6.17}$$

defined on the contact distribution $\mathbb{H} = \text{Ker}(\beta_0)$, where $\beta_0 = 2(dY + Z_A dX^A)$.

We now invoke Theorem 6.1.9, verifying by explicit computation that the remaining two conditions are satisfied. The first of these conditions is that the metric h_T is compatible with the Levi-form of the almost para-CR structure on the boundary. The second is that the Nijenhuis tensor takes asymptotically tangential values, i.e. that (6.5) is satisfied.

Both of these can be checked by computing the almost para–complex structure J in the (T, Z_A, X^A, Y) coordinates. We find

$$\begin{aligned} \mathcal{J} := J|_{T=0} = & -\frac{\partial}{\partial X^A} \otimes dX^A + \frac{\partial}{\partial Y} \otimes dY + \frac{\partial}{\partial Z_A} \otimes dZ_A + \frac{\partial}{\partial T} \otimes dT \\ & - \frac{Z_B}{K} \frac{\partial}{\partial T} \otimes dX^B - \frac{1}{K} \frac{\partial}{\partial T} \otimes dY \\ & - (\Gamma_{AB}^D Z_D + \Gamma_{AB}^n) \frac{\partial}{\partial Z_A} \otimes dX^B + (\Gamma_{nB}^D Z_D + \Gamma_{nB}^n) Z_C \frac{\partial}{\partial Z_C} \otimes dX^B \\ & - (\Gamma_{An}^D Z_D + \Gamma_{An}^n) \frac{\partial}{\partial Z_A} \otimes dY + (\Gamma_{nn}^D Z_D + \Gamma_{nn}^n) Z_C \frac{\partial}{\partial Z_C} \otimes dY. \end{aligned} \quad (6.18)$$

Restricting to vectors in \mathbb{H} amounts to substituting $dY = \beta_0/2 - Z_A dX^A$ and disregarding the terms involving β_0 as above, so that

$$\begin{aligned} J|_{\mathbb{H}} = & -\frac{\partial}{\partial X^A} \otimes dX^A + Z_A \frac{\partial}{\partial Y} \otimes dX^A + \frac{\partial}{\partial Z_A} \otimes dZ_A + \frac{\partial}{\partial T} \otimes dT \\ & - \frac{2Z_B}{K} \frac{\partial}{\partial T} \otimes dX^B - \Xi_{AB} \frac{\partial}{\partial Z_A} \otimes dX^B \end{aligned}$$

and the boundary compatibility condition is satisfied.

For the Nijenhuis condition, note that we need only consider components of \mathcal{N} with $a = T$ to verify (6.5). Let us use the notation $J^{(T)}$ for the one–form comprising the $\partial/\partial T$ components of J . We find this to be

$$\begin{aligned} J^{(T)} = & \left(-\frac{Z_B}{K} + \frac{T[2Z_B + (\Gamma_{AB}^D Z_D + \Gamma_{AB}^n)X^A + (\Gamma_{nB}^D Z_D + \Gamma_{nB}^n)Y]}{K} \right. \\ & \left. - T^2[P_{AB}X^A + P_{nB}Y] \right) dX^B \\ & + \left(-\frac{1}{K} + \frac{T[2 + (\Gamma_{An}^D Z_D + \Gamma_{An}^n)X^A + (\Gamma_{nn}^D Z_D + \Gamma_{nn}^n)Y]}{K} \right. \\ & \left. - T^2[P_{An}X^A + P_{nn}Y] \right) dY. \end{aligned}$$

Note that this agrees with (6.18) when $T = 0$. Using the formula (6.2), we now calculate

$$\mathcal{N}^a{}_{bc} \mathbf{g} \nabla_a T|_{T=0} = \left(\mathcal{J}^d_{[b} \partial_{|d|} J^{(T)}_{c]} - \mathcal{J}^d_{[b} \partial_{c]} J^{(T)}_{d]} \right) \Big|_{T=0}$$

to verify (6.5).

□

Remark 6.2.5. In the case if $n = 2$ let us use coordinates $(x, y, z) = (X^1, Y, Z_1)$ on ∂M , then (6.17) yields

$$h_{\mathbb{H}} = dz \odot dx - [\Gamma_{11}^2 + (\Gamma_{11}^1 - 2\Gamma_{12}^2)z + (\Gamma_{22}^2 - 2\Gamma_{12}^1)z^2 + \Gamma_{22}^1 z^3]dx \odot dx,$$

which is transparently invariant under the projective changes (3.1) of ∇ . In the two-dimensional case the projective structures $(N, [\nabla])$ are equivalent to second order ordinary differential equations (5.18) whose integral curves C are the unparametrised geodesics of ∇ . The curves C are integral submanifolds of a differential ideal $\mathcal{I} = < \beta_0, \beta_1 >$, where

$$\beta_0 = dy + zdx, \quad \beta_1 = dz - \left(\Gamma_{11}^2 + (\Gamma_{11}^1 - 2\Gamma_{12}^2)z + (\Gamma_{22}^2 - 2\Gamma_{12}^1)z^2 + \Gamma_{22}^1 z^3 \right)dx$$

are one-forms on a three-dimensional manifold $\mathcal{Z} = \mathbb{P}(T^*N)$ with local coordinates (x, y, z) . If $f : C \rightarrow \mathcal{Z}$ is an immersion, then $f^*(\beta_0) = 0, f^*(\beta_1) = 0$ is equivalent to (5.18) as long as $\beta_2 \equiv dx$ does not vanish. In terms of these three one-forms the contact structure, and the metric on the contact distribution are given by $\beta_0, h_{\mathbb{H}} = \beta_1 \odot \beta_2$.

6.3 An alternative approach to Theorem 6.2.4

It would be possible to show that the structures (M, g, Ω) arising in the projective to Einstein correspondence are para- c -projectively compact using a purely tractor-based approach, without relying on Theorem 6.1.9 and the local form (3.19). The basis for this alternative method is the *curved orbit decompositions* appearing in [19], which arise from *holonomy reductions* of Cartan geometries.

Recall first that a connection θ on a principal S -bundle $\pi : \mathcal{G} \rightarrow M$ defines a unique horizontal lift of any smooth curve on M , and we can define the *holonomy group* of θ based at $u \in \mathcal{G}$ as

$$\begin{aligned} \text{Hol}_u(\theta) = \{s \in S \mid u \text{ can be joined to } us \text{ by the} \\ \text{horizontal lift of a loop in } M \text{ based at } \pi(u)\}. \end{aligned}$$

Then $\text{Hol}_u(\theta)$ is a subgroup of S , and if M is connected then holonomy groups at different basepoints are related by conjugation in S . We can thus forget about the basepoint u by defining $\text{Hol}(\theta)$ as a conjugacy class of subgroups of S . Any subgroup

H such that $\text{Hol}(\theta) \subset H \subset S$ defines a reduced bundle $\mathcal{G} \times_S G/H$ with structure group H on which θ induces a connection.

Although the Cartan connection is not a principal bundle connection in the usual sense, one can still define a notion of holonomy for Cartan connections, and it turns out [19] that a parallel section of an associated tractor bundle defines a Cartan holonomy reduction by a subgroup $H \subset S$. The subgroup H decomposes the homogeneous model G/S into H -orbits, and it turns out that there is a corresponding decomposition of G/S in the curved case, with *reduced Cartan geometries* arising on the orbits. This is the origin of the name *curved orbit decomposition*. In the c -projective case, the model is decomposed into a pair of open orbits separated by a closed submanifold. The open orbits carry almost Kähler metrics and the closed orbit carries an almost CR structure.

By our construction above it follows that \mathcal{M} has a canonical para- c -projective geometry. In the following section, we realise the model in terms of an orbit decomposition of a Lie group quotient, which we expect to be the homogeneous model for a para- c -projective Cartan bundle. We conjecture that a full description of the corresponding Cartan connection would lead to a proof of Theorem 6.2.4 using the general Cartan holonomy theory in [19].

6.3.1 The model case

Recall from Section 3.2.3 that the flat projective structure on $N = \mathbb{RP}^n$ gives rise to the neutral signature para-Kähler Einstein metric (3.27) on the manifold

$$\begin{aligned} M &= \{([P], [L]) \in \mathbb{RP}^n \times \mathbb{RP}_n \mid P \cdot L \neq 0\} \\ &= \mathcal{M} \setminus \mathcal{Z}(\tau), \end{aligned}$$

where \mathcal{M} was the projectivised cotractor bundle $\mathbb{P}(\mathcal{T}^*)$ of \mathbb{RP}^n and \mathcal{Z} was the zero locus of the density τ , or equivalently the set of incident pairs in $\mathbb{RP}^n \times \mathbb{RP}_n$.

Here we shall instead take N to be the sphere S^n with its standard flat projective structure where the geodesics are great circles, so that N is orientable in all dimensions and the cotractor bundle is trivial. Note that S^n is a double cover of \mathbb{RP}^n which consists of the set of *oriented* lines in \mathbb{R}^{n+1} . We obtain it by taking an analogous quotient of \mathbb{R}^{n+1} where points are considered equivalent only up to multiplication by a *positive* number. We call this ray projectivisation and denote it $\mathbb{P}_+(\mathbb{R}^{n+1})$.

Replacing \mathbb{RP}^n with S^n allows us to write the cotractor bundle of N as $\mathcal{T}^* = S^n \times \mathbb{R}_3$, and if we also projectivise the fibres by ray projectivisation, we obtain a larger manifold

$$\widetilde{\mathcal{M}} = \mathbb{P}_+(\mathcal{T}^*) = S^n \times S_n,$$

where S_n is the dual to S^n in the same sense that \mathbb{RP}_n is dual to \mathbb{RP}^n . Then $\widetilde{\mathcal{M}}$ contains two copies of M which are separated by the hypersurface $\mathcal{Z}(\tau)$. We now express this decomposition of $\widetilde{\mathcal{M}}$ as an orbit decomposition.

Consider first two vector spaces \mathbb{V}, \mathbb{W} each isomorphic to \mathbb{R}^{n+1} , and view each as a representation space for an $SL(n+1, \mathbb{R})$ action. Define $G := SL(\mathbb{V}) \times SL(\mathbb{W})$ with its action on $\mathbb{V} \times \mathbb{W}$. We can write

$$\mathbb{P}_+(\mathbb{V}) \times \mathbb{P}_+(\mathbb{W}) = G/S = (SL(\mathbb{V})/P_V) \times (SL(\mathbb{W})/P_W)$$

where P_V (respectively P_W) is the parabolic subgroup in $SL(\mathbb{V})$ that stabilises a point $[V]$ in $\mathbb{P}_+(\mathbb{V})$ (respectively $[W] \in \mathbb{P}_+(\mathbb{W})$), and S is the group product $P_V \times P_W$ which itself is a parabolic subgroup of the semisimple group G . Since the action of $SL(\mathbb{V})$ descends to a transitive action on the ray projectivisation $\mathbb{P}_+(\mathbb{V})$ and similarly $SL(\mathbb{W})$ acts transitively on $\mathbb{P}_+(\mathbb{W})$, we have that $G := SL(\mathbb{V}) \times SL(\mathbb{W})$ acts transitively on the manifold $\mathbb{P}_+(\mathbb{V}) \times \mathbb{P}_+(\mathbb{W})$.

Note that we may consider \mathbb{V} and \mathbb{W} as the ± 1 eigenspaces of the single vector space $\mathbb{V} \oplus \mathbb{W}$ equipped with an endomorphism \mathbb{J} such that $\mathbb{J}^2 = 1$. Then the quotient G/S is exactly analogous to the model for c -projective geometry discussed in Section 6.1.3. We can therefore expect G/S to be the model for para- c -projective geometry.

Now introduce an additional structure which breaks the G symmetry. Namely we fix an isomorphism

$$I : \mathbb{W} \rightarrow \mathbb{V}^*$$

where \mathbb{V}^* denotes the dual space to \mathbb{V} . The subgroup $H \cong SL(n+1, \mathbb{R})$ of G that fixes this may be identified with $SL(\mathbb{V})$ which acts on a pair $(V, W) \in \mathbb{V} \times \mathbb{V}^*$ by the defining representation and on the first factor and by the dual representation on the second factor. Note in particular that this action preserves $W(V)$.

Given this structure we may now (suppress I and) write

$$\widetilde{\mathcal{M}} = \mathbb{P}_+(\mathbb{V}) \times \mathbb{P}_+(\mathbb{V}^*).$$

The H action on $\widetilde{\mathcal{M}}$ has two open orbits and a closed orbit. The last is the incidence space

$$\mathcal{Z} = \{([V], [W]) \in \mathcal{M} \mid W(V) = 0\}$$

which sits as smooth orientable separating hypersurface in $\widetilde{\mathcal{M}}$. Then there are the open orbits

$$M_+ = \{([V], [W]) \in \mathcal{M} \mid W(V) > 0\} \quad \text{and} \quad M_- = \{([V], [W]) \in \mathcal{M} \mid W(V) < 0\}.$$

We may think of \mathcal{Z} as the ‘boundary’ (at infinity) for the open orbits M_{\pm} , each of which is a copy of our para–Kähler Einstein manifold M .

We therefore have an orbit decomposition of the homogeneous space G/S , where the additional structure I has induced a reduction of the holonomy group by H . Based on the results of [19] for the analogous c –projective case, we expect the closed orbit \mathcal{Z} to carry a para–CR structure and the open orbits M_{\pm} to carry para–Kähler metrics which are induced by the dual pairing between \mathbb{V} and \mathbb{V}^* just as we saw in Section 6.2.1.

Chapter 7

Concluding Remarks

We use this chapter to discuss some of the open questions which remain unanswered by the work presented here.

We began by discussing a modified ϕ^4 theory on a wormhole spacetime, finding that there is a kink solution and that it is topologically and linearly stable. We investigated its asymptotic stability for the range of a where exactly one discrete mode is present. It would be interesting to expand the investigation in Section 2.4 to the case when both discrete modes are present. This problem is much more complicated because of the extra terms which arise from the amplitude of the second internal mode. Similar problems have been discussed in [81], although no such analysis has been done for non-linear Klein–Gordon equation of this type with two discrete modes. The ϕ^4 theory on the wormhole presents a useful setting to undertake such analysis because the kink has exactly two discrete modes for any $a > a_1$, and because their frequencies can be controlled by the parameter a .

The modified ϕ^4 model shares an interesting property with the modified sine–Gordon theory on the same wormhole spacetime [6]. In both cases, we expect a discontinuous change in decay behaviour when a moves out of the range $a_0 < a < a_1$. Insight from the ϕ^4 case may help to elucidate the character of such discontinuous changes.

In this thesis, we have considered the manifolds M arising in the projective to Einstein correspondence from a number of different perspectives. However, there remain several open questions. In Chapter 4 we showed that a second Einstein metric can be canonically constructed on an \mathbb{R}^* –bundle over M , and interpreted this space in terms of the Cartan bundle of the projective structure. It would be interesting to understand what this larger manifold is from the tractor perspective.

In Chapter 5 we constructed several examples of Einstein–Weyl structures arising as Jones–Tod reductions of the Einstein manifolds M in the special case $n = 2$, where

M has ASD conformal curvature, and extracted corresponding solutions to the $SU(\infty)$ –Toda equation. Although we produced many local expressions for Einstein–Weyl structures, we were unable to provide a coordinate and scale invariant local characterisation of the Einstein–Weyl structures which are obtainable from the projective to Einstein correspondence. We know that they belong to the $SU(\infty)$ –Toda class, and we showed in Section 5.2.2 that they carry two solutions to the Abelian monopole equation, but it would be interesting to try and extend this result to a complete local characterisation.

Finally, in Chapter 6 we showed that the manifolds M are *para-c-projectively compact* using a local form of the metric. A more fundamental understanding of this result could be obtained by realising M in terms of the curved orbit decomposition of a para- c –projective structure, viewed as a Cartan holonomy reduction.

References

- [1] Arnold, V. I. *Mathematical Methods of Classical Mechanics*. Springer, 1989.
- [2] Atiyah, M. F. Hitchin, N. J. and Singer, I. M. (1978) Self-duality in four-dimensional Riemannian geometry. *Proc. Roy. Soc. London Ser. A* **362**, 425–461.
- [3] Atiyah, M., Dunajski, M. and Mason, L. (2017) Twistor theory at fifty: from contour integrals to twistor strings. *Proc. Roy. Soc.* **473**.
- [4] Bailey, T. N., Eastwood M. G. and Gover, A. R. (1994) Thomas's structure bundle for conformal, projective and related structures. *Rocky Mt. J. Math.* **24**, 1–27.
- [5] Bizoń, P. and Kahl, M. (2015) Wave maps on a wormhole. *Phys. Rev. D* **91**, 065003
- [6] Bizoń, P., Dunajski, M., Gibbons, G., Kahl, M. and Kowalczyk, M. (2018) Sine-Gordon on a Wormhole (preprint).
- [7] Boyer, C. (1988) A note on hyperhermitian four-manifolds *Proc. Amer. Math. Soc.* **102**, 157–164.
- [8] Boyer, C. and Finley, D. (1982) Killing vectors in self-dual, Euclidean Einstein spaces. *J. Math. Phys.* **23**, 1126.
- [9] Bronnikov, K. A. (1973) Scalar-tensor theory and scalar charge. *Acta Phys. Polonica* **b4**, 251–266
- [10] Bryant, R. L., Dunajski, M. and Eastwood, M. (2009) Metrisability of two-dimensional projective structures. *J. Differ. Geom.* **83**, 465–499.
- [11] Calderbank, D. M. J. (2000) The geometry of the Toda equation. *J. Geom. Phys.* **36**, 152–162.
- [12] Calderbank, D. M. J. and Tod, K. P. (2001) Einstein metrics, hypercomplex structures and the Toda field equation. *Differ. Geom. Appl.* **14**, 199–208.
- [13] Calderbank, D. M. J. (2014) Integrable Background Geometries. *SIGMA* **10**, 034.
- [14] Calderbank, D. M. J. (2014) Selfdual 4-manifolds, projective structures, and the Dunajski-West construction. *SIGMA* **10**, 035.
- [15] Calderbank, D. M. J., Eastwood, M. G., Matveev, V. S. and Neusser, K. (2016) C-projective geometry. [arXiv:1512.04516](https://arxiv.org/abs/1512.04516)

[16] Calvino-Louzao, E., García-Río, E., Gilkey, P. and Vazquez-Lorenzo, R. (2009) The geometry of modified Riemannian extensions. *Proc. Roy. Soc. A* **465**, 2107

[17] Čap, A. and Gover, A. R. (2016) Projective Compactifications and Einstein metrics. *J. Reine Angew. Math.* **717**, 47–75.

[18] Čap, A. and Gover, A. R. (2019) C-Projective Compactification; (quasi-)Kaehler Metrics and CR boundaries. *Am. J. Math.* **141**, 813–856.

[19] Čap, A., Gover, A. R. and Hammerl, M. (2014) Holonomy reductions of Cartan geometries and curved orbit decompositions. *Duke Math. J.* **163**, 1035–1070.

[20] Cartan, E. (1924) Sur les variétés à connexion projective. *Bull. Soc. Math. France* **52**, 205–241.

[21] Casey, S., Dunajski, M. and Tod, K. P. (2013) Twistor geometry of a pair of second order ODEs. *Comm. Math. Phys.* **321**, 681–701.

[22] Cherkis, S. A. and Kapustin, A. (1999) Singular monopoles and gravitational instantons. *Comm. Math. Phys.* **203**, 713–728.

[23] Derrick, G. H. (1964) Comments on nonlinear wave equations as models from elementary particles. *J. Math. Phys.* **5**, 1252–1254.

[24] Dunajski, M. (1999) The twisted photon associated to hyper-Hermitian four-manifolds. *J. Geom. Phys.* **30**, 266–281.

[25] Dunajski, M. *Solitons, Instantons, and Twistors*. Oxford Graduate Texts in Mathematics, Oxford University Press, 2009.

[26] Dunajski, M. and Tod, K. P. (2010) Four Dimensional Metrics Conformal to Kähler. *Math. Proc. Camb. Phil. Soc.* **148**, 485–503.

[27] Dunajski, M., Ferapontov, E. and Kruglikov, B. (2015) On the Einstein–Weyl and conformal self-duality equations. *J. Math. Phys.* **56**, 8.

[28] Dunajski, M. and Mettler, T. (2018) Gauge theory on Projective Surfaces and Anti-self-dual Einstein Metrics in Dimension Four. *J. Geom. Anal.* **28**, 2780–2811.

[29] Dunajski M., Gover, A. R. and Waterhouse, A. (2018) Some examples of projective and c-projective compactifications of Einstein metrics. [arXiv:1807.06105](https://arxiv.org/abs/1807.06105)

[30] Dunajski, M. and Waterhouse, A. (2019) Einstein Metrics, Projective Structures and the $SU(\infty)$ Toda Equation. *J. Geom. Phys.* **147**, 103523.

[31] Eastwood, M. (2008) Notes on Projective Differential Geometry, in *Symmetries and Overdetermined Systems of Partial Differential Equations*, The IMA Volumes in Mathematics and its Applications vol **144** Springer, New York, NY.

[32] Ellis, H. G. (1973) Ether flow through a drainhole: A particle model in general relativity. *J. Math. Phys.* **14**, 104–118.

[33] Falla Luza, H. M. and Loray, F. (2017) Projective structures and neighbourhoods of rational curves. *arXiv:1707.07868v1*

[34] Ferapontov, E. V., Korotkin D. A. and Shramchenko, V. A. (2002) Boyer–Finley equation and systems of hydrodynamic type. *Class. Quantum Gravity* **19**, 24.

[35] Finley, J.D. and Plebański, J.F. (1979) The classification of all H spaces admitting a Killing vector. *J. Math. Phys.* **20**, 1938.

[36] Gibbons, G. W. and Hawking, S. W. (1978) Gravitational Multi - Instantons. *Phys. Lett.* **B78**, 430.

[37] Gibbons, G. W. and Hawking, S. W. (1979) Classification of Gravitational Instanton Symmetries. *Comm. Math. Phys.* **66**, 291–310.

[38] Grossman, D. (2000) Torsion-free path geometries and integrable second order ODE systems. *Selecta Math. (N.S.)* **6**, 399–442.

[39] Hitchin, N. J. (1982) in *Twistor Geometry and Non-linear Systems*, Lecture Notes in Mathematics vol **970** Springer, Berlin.

[40] Jones, P. E. and Tod, K. P (1985) Minitwistor spaces and Einstein-Weyl spaces. *Class. Quantum Gravity* **15**, 565–577.

[41] Klein, F. (1872) Vergleichende Betrachtungen über neuere geometrische Forschungen. *Math. Ann.* **43**, 63–100.

[42] Kobayashi, S. (1963) Topology of positively pinched Kähler manifolds. *Tohoku Math. J.* **2**, 121–139.

[43] Kobayashi, S. and Nagano, T. (1964) On projective connections. *J. Math. Mech.* **13**, 215–235.

[44] Kodaira, K. (1963) On stability of compact submanifolds of complex manifolds. *Am. J. Math.* **85**, 79–94.

[45] Kowalczyk, M., Martel, Y. and Muñoz, C. (2017) Kink Dynamics in the ϕ^4 Model: Asymptotic Stability for Odd Perturbations in the Energy Space. *J. Amer. Math. Soc.* **30**, 769–798.

[46] Kowalski, O. and Vlášek, Z. (2007) Classification of locally projectively homogeneous torsion-less affine connections in the plane domains. *Beitr. Algebra Geom.* **48**, 11–26.

[47] LeBrun, C. (1983) Spaces of Complex Null Geodesics in Complex–Riemannian geometry. *Trans. Am. Math. Soc.* **278**, 209–231.

[48] LeBrun, C. (1991) Explicit self–dual metrics on $\mathbb{CP}_2 \# \dots \# \mathbb{CP}_2$. *J. Differ. Geom.* **34**, 223–253.

[49] Levi–Civita, T. (1896) On the transformations of the dynamical equations. *Ann. Mat. Pura Appl.* **24**, 255–300.

[50] Libermann, P. (1954) Sur le probleme d'équivalence de certaines structures infinitesimales. *Ann. Mat. Pura Appl.* **36**, 27–120.

[51] Manakov, S. V. and Santini, P. M. (2009) The dispersionless 2D Toda equation: dressing, Cauchy problem, longtime behaviour, implicit solutions and wave breaking. *J. Phys.* **A42**.

[52] Manton, N. S. and Merabet, H. (1997) ϕ^4 Kinks - Gradient Flow and Dynamics. *Nonlinearity* **10**, 3–18.

[53] Manton, N. S. and Sutcliffe, P. (2004) *Topological Solitons*. Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge.

[54] Martina, L. Sheftel, M. and Winternitz, P. (2001) Group foliation and non-invariant solutions of the heavenly equation. *J. Phys.* **A34**.

[55] Mason, L. J. and Woodhouse, N. M. J. *Integrability, self-duality and twistor theory*. London Mathematical Society Monographs, New Series, Vol. 15, The Clarendon Press, Oxford University Press, New York, 1996.

[56] Matveev, V. S. and Rosemann, S. (2015) The degree of mobility of Einstein metrics. *J. Geom. Phys.* **99**, 42–56.

[57] Melrose, R. B. (1995) *Geometric scattering theory*. Stanford Lectures. Cambridge University Press, Cambridge.

[58] Morrow, J. and Kodaira, K. (2006) *Complex Manifolds*. AMS bookstore, Providence, RI.

[59] Newlander, A. and Nirenberg, L. (1957) Complex analytic coordinates in almost complex manifolds, *Ann. Math.* **65**, 391–404.

[60] Pedersen, H. and Tod, P. (1993) Three-Dimensional Einstein-Weyl Geometry, *Adv. Math.* **97**, 74–109.

[61] Penrose, R. (1965) Zero rest-mass fields including gravitation: asymptotic behaviour. *Proc. R. Soc. London, Ser. A* **284**, 159–203.

[62] Penrose, R. (1976) Nonlinear gravitons and curved twistor theory. *Gen. Rel. Grav.* **7**, 31–52.

[63] Penrose, R. and Rindler, W. *Spinors and Spacetime, Volumes I and II*. Cambridge University Press 1984.

[64] Pope, C. Kaluza–Klein Theory.
<http://people.physics.tamu.edu/pope/ihplec.pdf>

[65] Przanowski, M. (1991) Killing vector fields in self-dual, Euclidean Einstein spaces with $\Lambda \neq 0$. *J. Math. Phys.* **32**, 1004–1010.

[66] Sharpe, R. W. *Differential Geometry: Cartan's Generalization of Klein's Erlangen Program*. Springer, New York, 1997.

[67] Sheftel, M. (2003) Method of group foliation, hodograph transformation and non-invariant solutions of the Boyer-Finley equation. *Theor. Math. Phys.* **137**, 1743–1752.

[68] Simon, B. (1975) The Bound State of Weakly Coupled Schrödinger Operators in One and Two Dimensions. *Ann. Phys.* **97**, 279–288.

[69] Soffer, A. and Weinstein, M. I. (1998) Time Dependent Resonance Theory. *Geom. Funct. Anal.* **8**, 1086–1128.

[70] Soffer, A. and Weinstein, M. I. (1999) Resonances, radiation damping and instability in Hamiltonian nonlinear wave equations. *Invent. math.* **136**, 9–74.

[71] Tao, T. (2009) Why are Solitons Stable? *Am. Math. Soc.* **46**, 1–33.

[72] Tod, P. (1995) Scalar–flat Kähler and hyper–Kähler metrics from Painlevé –III. *Class. Quantum Grav.* **12**, 1535–1547.

[73] Tod, P. (1997) The $SU(\infty)$ –Toda field equation and special four–dimensional metrics, in *Geometry and Physics* (Aarhus, 1995), Lecture Notes in Pure and Applied Mathematics **184**, 307–312 Dekker, New York.

[74] Tod, P. (2006) A note on Riemannian anti–self–dual Einstein metrics with symmetry [arXiv:hep-th/0609071](https://arxiv.org/abs/hep-th/0609071)

[75] Uhlenbeck, K. (1982) Removable singularities in Yang–Mills fields. *Commun. Math. Phys.* **83**, 11–29.

[76] Ward, R. S. (1976) The Twisted Photon. *Twistor Newsletter* **1**, 7–8.

[77] Ward, R. S. (1977) On Self–Dual Gauge Fields. *Phys. Lett. A* **61**, 81–82.

[78] Ward, R. S. (1980) Self–dual space–times with cosmological constant, *Comm. Math. Phys.* **78**, 1–17.

[79] Ward, R. S. (1990) Einstein–Weyl spaces and $SU(\infty)$ Toda fields. *Class. Quantum Gravity* **7**, L95–L98.

[80] Waterhouse, A. (2019) The ϕ^4 kink on a wormhole spacetime. [arXiv: 1908.09650](https://arxiv.org/abs/1908.09650)

[81] Weinstein, M. I. (2002) Extended Hamiltonian Systems. *Handbook of Dynamical Systems* IB, 1135–1153.

[82] West, S. *Neutral ASD Conformal Structures with Null Killing Vectors*. Thesis. University of Cambridge, 2007.

[83] Witten, E. (1998) Anti de Sitter space and holography. *Adv. Theor. Math. Phys.* **2**, 253–291.

[84] Xinliang, An. and Soffer, A. (2017) Fermi’s golden rule and H^1 scattering for nonlinear Klein–Gordon equations with metastable states. [arXiv:1510.06485](https://arxiv.org/abs/1510.06485)

- [85] Yano, K. *The Theory Of Lie Derivatives And Its Applications*. North Holland, Amsterdam, 1955.
- [86] Zenginoglu, A. (2008) Hyperboloidal foliations and scri-fixing. *Class. Quantum Gravity* **25**, 145002.
- [87] Zenginoglu, A. (2008) A hyperboloidal study of tail decay rates for scalar and Yang-Mills fields. *Class. Quantum Gravity* **25**, 175013.