

# Koopman Spectra from Data: Guarantees, Limitations, and Implications for Prediction

Matthew Colbrook 11/06/2026



Papers and talks:

[http://www.damtp.cam.ac.uk/  
user/mjc249/home.html](http://www.damtp.cam.ac.uk/user/mjc249/home.html)

# Amazing collaborators and friends!



**Nicolas  
Boullé**  
(Imperial)



**Steve  
Brunton**  
(UW)



**Gustav  
Conradie**  
(Cambridge)



**JC Loiseau**  
(Arts et  
Métiers)



**Igor Mezić**  
(UC Santa  
Barbara)



**Alexei  
Stepanenko**  
(Industry)



**Alex  
Townsend**  
(Cornell)

- C., Townsend. *"Rigorous data-driven computation of spectral properties of Koopman operators for dynamical systems."* **Communications on Pure and Applied Mathematics**, 2024.
- C., Mezić, Stepanenko, *"Adversarial dynamical systems characterize when data-driven learning succeeds or fails."* **Nature Communications**, 2026.
- Boullé, C., Conradie, *"Convergent Methods for Koopman Operators on Reproducing Kernel Hilbert Spaces."* <https://arxiv.org/abs/2506.15782>
- Conradie, Boullé, Loiseau, Brunton, C., *"Trustworthy Koopman Operator Learning: Invariance Diagnostics and Error Bounds."* <https://arxiv.org/abs/2603.15091>

# What is a Koopman operator?

- $\mathcal{X}$  – *the state space*
- $\mathcal{X} \ni x$  – *the state*

cts  $F: \mathcal{X} \rightarrow \mathcal{X}$  – *the dynamics*:  $x_{n+1} = F(x_n)$

Henri Poincaré  
(Sorbonne)



# What is a Koopman operator?

- $\mathcal{X}$  – the state space
- $\mathcal{X} \ni x$  – the state


cts  $F: \mathcal{X} \rightarrow \mathcal{X}$  – the dynamics:  $x_{n+1} = F(x_n)$

- Functions  $g: \mathcal{X} \rightarrow \mathbb{C}$  a.k.a “observables”,  $g \in L^2(\mathcal{X}, \omega)$
- Koopman operator  $\mathcal{K}_F: [\mathcal{K}_F g](x) = g(F(x))$

**LINEAR!**

Observe  $g$  one time step forward

Bernard Koopman  
(Columbia)



John von Neumann  
(IAS)



- Koopman, “Hamiltonian systems and transformation in Hilbert space,” *Proc. Natl. Acad. Sci. USA*, 1931.
- Koopman, v. Neumann, “Dynamical systems of continuous spectra,” *Proc. Natl. Acad. Sci. USA*, 1932.

# What is a Koopman operator?

- $\mathcal{X}$  – the state space
- $\mathcal{X} \ni x$  – the state
- Unknown cts  $F: \mathcal{X} \rightarrow \mathcal{X}$  – the dynamics:  $x_{n+1} = F(x_n)$
- Functions  $g: \mathcal{X} \rightarrow \mathbb{C}$  a.k.a “observables”,  $g \in L^2(\mathcal{X}, \omega)$
- Koopman operator  $\mathcal{K}_F: [\mathcal{K}_F g](x) = g(F(x))$  **LINEAR!**
- Available snapshot data:  $\left\{ \left( x^{(m)}, y^{(m)} = F(x^{(m)}) \right) : m = 1, \dots, M \right\}$

**Can we compute spectral properties from trajectory data?**

$$g(x_n) = [\mathcal{K}^n g](x_0)$$


Why?

If  $\|\mathcal{K}g - \lambda g\| \leq \varepsilon$ , then  $g(x_n) = [\mathcal{K}^n g](x_0) = \lambda^n g(x_0) + \mathcal{O}(n\varepsilon)$

$$g(x_n) = [\mathcal{K}^n g](x_0)$$

Why?

If  $\|\mathcal{K}g - \lambda g\| \leq \varepsilon$ , then  $g(x_n) = [\mathcal{K}^n g](x_0) = \lambda^n g(x_0) + \mathcal{O}(n\varepsilon)$



**Coherent features!**

$$\text{Sp}_\varepsilon(\mathcal{K}) = \{z \in \mathbb{C} : \exists g, \|g\| = 1, \|\mathcal{K}g - zg\| \leq \varepsilon\}$$

**Trades:** Nonlinear, finite-dimensional  $\Rightarrow$  Linear, infinite-dimensional.

# Koopman Mode Decomposition

- Find  $(g_j, \lambda_j)$  with  $\|\mathcal{K}g_j - \lambda_j g_j\| \leq \varepsilon$
- Expand observable:

$$h(x) \approx \sum_j c_j g_j(x)$$

Verified Eigenfunctions

coefficients, called  
"Koopman modes"

- Forecasts:

$$h(x_n) = \sum_j \lambda_j^n c_j g_j(x) + \mathcal{O}(n\varepsilon)$$

**Intuition:** A nonlinear separation of variables through a linear operator!

# Perils of discretization: Warmup on $\ell^2(\mathbb{Z})$

$$\begin{pmatrix} \ddots & & & & & & \\ & \ddots & & & & & \\ & & 0 & 1 & & & \\ & & & 0 & 1 & & \\ & & & & 0 & 1 & \\ & & & & & 0 & 1 \\ & & & & & & 0 & \ddots \\ & & & & & & & \ddots \end{pmatrix} \xrightarrow{\text{Two-way infinite}} \begin{pmatrix} 0 & 1 & & & & \\ & \ddots & \ddots & & & \\ & & \ddots & \ddots & & \\ & & & \ddots & 1 & \\ & & & & 1 & 0 \end{pmatrix} \in \mathbb{C}^{N \times N}$$

- Spectrum is unit circle.
- Spectrum is stable.
- Continuous spectra.
- Unitary evolution.

- Spectrum is  $\{0\}$ .
- Spectrum is unstable.
- Discrete spectra.
- Nilpotent evolution.

**Lots of Koopman operators are built up from operators like these!**

# Explicit example: Matrix approximation of $\mathcal{K}$ (EDMD)

Observables  $\psi_j: \mathcal{X} \rightarrow \mathbb{C}, j = 1, \dots, N$

$$\{x^{(m)}, y^{(m)} = F(x^{(m)})\}_{m=1}^M$$

quadrature points

$$\langle \psi_k, \psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(x^{(m)})} \psi_k(x^{(m)}) = \left[ \underbrace{\begin{pmatrix} \psi_1(x^{(1)}) & \dots & \psi_N(x^{(1)}) \\ \vdots & \ddots & \vdots \\ \psi_1(x^{(M)}) & \dots & \psi_N(x^{(M)}) \end{pmatrix}}_{\Psi_X} \right]^* \underbrace{\begin{pmatrix} w_1 & & \\ & \ddots & \\ & & w_M \end{pmatrix}}_W \underbrace{\begin{pmatrix} \psi_1(x^{(1)}) & \dots & \psi_N(x^{(1)}) \\ \vdots & \ddots & \vdots \\ \psi_1(x^{(M)}) & \dots & \psi_N(x^{(M)}) \end{pmatrix}}_{\Psi_X} \right]_{jk}$$

quadrature weights

$$\langle \mathcal{K}\psi_k, \psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(x^{(m)})} \underbrace{\psi_k(y^{(m)})}_{[\mathcal{K}\psi_k](x^{(m)})} = \left[ \underbrace{\begin{pmatrix} \psi_1(x^{(1)}) & \dots & \psi_N(x^{(1)}) \\ \vdots & \ddots & \vdots \\ \psi_1(x^{(M)}) & \dots & \psi_N(x^{(M)}) \end{pmatrix}}_{\Psi_X} \right]^* \underbrace{\begin{pmatrix} w_1 & & \\ & \ddots & \\ & & w_M \end{pmatrix}}_W \underbrace{\begin{pmatrix} \psi_1(y^{(1)}) & \dots & \psi_N(y^{(1)}) \\ \vdots & \ddots & \vdots \\ \psi_1(y^{(M)}) & \dots & \psi_N(y^{(M)}) \end{pmatrix}}_{\Psi_Y} \right]_{jk}$$


Galerkin  
Approximation

$$\mathcal{K} \rightarrow (\Psi_X^* W \Psi_X)^{-1} \Psi_X^* W \Psi_Y \in \mathbb{C}^{N \times N}$$

- Schmid, "Dynamic mode decomposition of numerical and experimental data," **J. Fluid Mech.**, 2010.
- Rowley, Mezić, Bagheri, Schlatter, Henningson, "Spectral analysis of nonlinear flows," **J. Fluid Mech.**, 2009.
- Williams, Kevrekidis, Rowley "A data-driven approximation of the Koopman operator: Extending dynamic mode decomposition," **J. Nonlinear Sci.**, 2015.

# EDMD doesn't converge!

- Duffing oscillator:  $\dot{x} = y$ ,  $\dot{y} = -\alpha y + x(1 - x^2)$ , sampled  $\Delta t = 0.3$ .
- Gaussian radial basis functions, Monte Carlo integration ( $M = 50000$ )



# The fix: Residual DMD (ResDMD)

$$\langle \psi_k, \psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(x^{(m)})} \psi_k(x^{(m)}) = \left[ \underbrace{\Psi_X^* W \Psi_X}_G \right]_{jk}$$


$$\langle \mathcal{K}\psi_k, \psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(x^{(m)})} \underbrace{\psi_k(y^{(m)})}_{[\mathcal{K}\psi_k](x^{(m)})} = \left[ \underbrace{\Psi_X^* W \Psi_Y}_A \right]_{jk}$$

- C., Townsend, "Rigorous data-driven computation of spectral properties of Koopman operators for dynamical systems," **Commun. Pure Appl. Math.**, 2023.
- C., Ayton, Szóke, "Residual Dynamic Mode Decomposition," **J. Fluid Mech.**, 2023.
- Code: <https://github.com/MColbrook/Residual-Dynamic-Mode-Decomposition>

# The fix: Residual DMD (ResDMD)

$$\langle \psi_k, \psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(x^{(m)})} \psi_k(x^{(m)}) = \left[ \underbrace{\Psi_X^* W \Psi_X}_G \right]_{jk}$$

$$\langle \mathcal{K}\psi_k, \psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(x^{(m)})} \underbrace{\psi_k(y^{(m)})}_{[\mathcal{K}\psi_k](x^{(m)})} = \left[ \underbrace{\Psi_X^* W \Psi_Y}_A \right]_{jk}$$




- C., Townsend, "Rigorous data-driven computation of spectral properties of Koopman operators for dynamical systems," **Commun. Pure Appl. Math.**, 2023.
- C., Ayton, Szóke, "Residual Dynamic Mode Decomposition," **J. Fluid Mech.**, 2023.
- Code: <https://github.com/MColbrook/Residual-Dynamic-Mode-Decomposition>

# The fix: Residual DMD (ResDMD)

$$\langle \psi_k, \psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(x^{(m)})} \psi_k(x^{(m)}) = \underbrace{[\Psi_X^* W \Psi_X]}_G]_{jk}$$

$$\langle \mathcal{K}\psi_k, \psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(x^{(m)})} \underbrace{\psi_k(y^{(m)})}_{[\mathcal{K}\psi_k](x^{(m)})} = \underbrace{[\Psi_X^* W \Psi_Y]}_A]_{jk}$$

$$\langle \mathcal{K}\psi_k, \mathcal{K}\psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(y^{(m)})} \psi_k(y^{(m)}) = \underbrace{[\Psi_Y^* W \Psi_Y]}_L]_{jk}$$



- C., Townsend, "Rigorous data-driven computation of spectral properties of Koopman operators for dynamical systems," **Commun. Pure Appl. Math.**, 2023.
- C., Ayton, Szóke, "Residual Dynamic Mode Decomposition," **J. Fluid Mech.**, 2023.
- Code: <https://github.com/MColbrook/Residual-Dynamic-Mode-Decomposition>

# The fix: Residual DMD (ResDMD)

$$\langle \psi_k, \psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(x^{(m)})} \psi_k(x^{(m)}) = \left[ \underbrace{\Psi_X^* W \Psi_X}_G \right]_{jk}$$

$$\langle \mathcal{K}\psi_k, \psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(x^{(m)})} \underbrace{\psi_k(y^{(m)})}_{[\mathcal{K}\psi_k](x^{(m)})} = \left[ \underbrace{\Psi_X^* W \Psi_Y}_A \right]_{jk}$$

$$\langle \mathcal{K}\psi_k, \mathcal{K}\psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(y^{(m)})} \psi_k(y^{(m)}) = \left[ \underbrace{\Psi_Y^* W \Psi_Y}_L \right]_{jk}$$

**Residuals:**  $g = \sum_{j=1}^N \mathbf{g}_j \psi_j$ ,  $\|\mathcal{K}g - \lambda g\|^2 = \langle \mathcal{K}g - \lambda g, \mathcal{K}g - \lambda g \rangle$

- C., Townsend, "Rigorous data-driven computation of spectral properties of Koopman operators for dynamical systems," *Commun. Pure Appl. Math.*, 2023.
- C., Ayton, Szóke, "Residual Dynamic Mode Decomposition," *J. Fluid Mech.*, 2023.
- Code: <https://github.com/MColbrook/Residual-Dynamic-Mode-Decomposition>

# The fix: Residual DMD (ResDMD)

$$\langle \psi_k, \psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(x^{(m)})} \psi_k(x^{(m)}) = \left[ \underbrace{\Psi_X^* W \Psi_X}_G \right]_{jk}$$

$$\langle \mathcal{K}\psi_k, \psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(x^{(m)})} \underbrace{\psi_k(y^{(m)})}_{[\mathcal{K}\psi_k](x^{(m)})} = \left[ \underbrace{\Psi_X^* W \Psi_Y}_A \right]_{jk}$$

$$\langle \mathcal{K}\psi_k, \mathcal{K}\psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(y^{(m)})} \psi_k(y^{(m)}) = \left[ \underbrace{\Psi_Y^* W \Psi_Y}_L \right]_{jk}$$

**Residuals:**  $g = \sum_{j=1}^N \mathbf{g}_j \psi_j$ ,  $\|\mathcal{K}g - \lambda g\|^2 = \sum_{k,j=1}^N \mathbf{g}_k \overline{\mathbf{g}_j} \langle \mathcal{K}\psi_k - \lambda \psi_k, \mathcal{K}\psi_j - \lambda \psi_j \rangle$

- C., Townsend, "Rigorous data-driven computation of spectral properties of Koopman operators for dynamical systems," **Commun. Pure Appl. Math.**, 2023.
- C., Ayton, Szóke, "Residual Dynamic Mode Decomposition," **J. Fluid Mech.**, 2023.
- Code: <https://github.com/MColbrook/Residual-Dynamic-Mode-Decomposition>

# The fix: Residual DMD (ResDMD)

$$\langle \psi_k, \psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(x^{(m)})} \psi_k(x^{(m)}) = \left[ \underbrace{\Psi_X^* W \Psi_X}_G \right]_{jk}$$

$$\langle \mathcal{K}\psi_k, \psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(x^{(m)})} \underbrace{\psi_k(y^{(m)})}_{[\mathcal{K}\psi_k](x^{(m)})} = \left[ \underbrace{\Psi_X^* W \Psi_Y}_A \right]_{jk}$$

$$\langle \mathcal{K}\psi_k, \mathcal{K}\psi_j \rangle \approx \sum_{m=1}^M w_m \overline{\psi_j(y^{(m)})} \psi_k(y^{(m)}) = \left[ \underbrace{\Psi_Y^* W \Psi_Y}_L \right]_{jk}$$




**Residuals:**  $g = \sum_{j=1}^N \mathbf{g}_j \psi_j$ ,  $\|\mathcal{K}g - \lambda g\|^2 = \lim_{M \rightarrow \infty} \mathbf{g}^* [L - \lambda A^* - \bar{\lambda} A + |\lambda|^2 G] \mathbf{g}$

- C., Townsend, "Rigorous data-driven computation of spectral properties of Koopman operators for dynamical systems," *Commun. Pure Appl. Math.*, 2023.
- C., Ayton, Szóke, "Residual Dynamic Mode Decomposition," *J. Fluid Mech.*, 2023.
- Code: <https://github.com/MColbrook/Residual-Dynamic-Mode-Decomposition>

# ResDMD does converge!

- Duffing oscillator:  $\dot{x} = y, \dot{y} = -\alpha y + x(1 - x^2)$ , sampled  $\Delta t = 0.3$ .
- Gaussian radial basis functions, Monte Carlo integration ( $M = 50000$ )

Compute  $\text{Sp}_\varepsilon(\mathcal{K})$ , local adaptive control on  $\varepsilon \downarrow 0$



# What can we do?

Consider space of observables with finite energy:  $L^2(\mathcal{X}, \omega)$

**Theorem:** There **exists** algorithms  $\Gamma_{N,M}$  using snapshots such that

$$\lim_{N \rightarrow \infty} \lim_{M \rightarrow \infty} \Gamma_{N,M}(F) = \text{Sp}_\varepsilon(\mathcal{K}_F)$$

for all systems.



$N$  = size of basis,     $M$  = amount of data (quadrature)

$$\text{Sp}_\varepsilon(\mathcal{K}) = \{z \in \mathbb{C} : \exists g, \|g\| = 1, \|\mathcal{K}g - zg\| \leq \varepsilon\}$$

# What can we do?

Consider space of observables with finite energy:  $L^2(\mathcal{X}, \omega)$

**Theorem:** There **exists** algorithms  $\Gamma_{N,M}$  using snapshots such that

$$\lim_{N \rightarrow \infty} \lim_{M \rightarrow \infty} \Gamma_{N,M}(F) = \text{Sp}_\varepsilon(\mathcal{K}_F)$$

for all systems.

$N$  = size of basis,     $M$  = amount of data (quadrature)

Double limit  $\lim_{N \rightarrow \infty} \lim_{M \rightarrow \infty}$

**Can we do better?**

# Adversaries: Double limit is necessary!

Implies  $\mathcal{K}$  is unitary

*Class of systems:*  $\Omega_{\mathbb{D}} = \{F: \bar{\mathbb{D}} \rightarrow \bar{\mathbb{D}} \mid F \text{ cts, measure preserving, invertible}\}$ .

*Data an algorithm can use:*  $\mathcal{T}_F = \{(x, y_m) \mid x \in \bar{\mathbb{D}}, \|F(x) - y_m\| \leq 2^{-m}\}$ .

**Theorem:** There **does not exist** any sequence of deterministic algorithms  $\{\Gamma_n\}$  using  $\mathcal{T}_F$  such that  $\lim_{n \rightarrow \infty} \Gamma_n(F) = \text{Sp}_\varepsilon(\mathcal{K}_F) \forall F \in \Omega_{\mathbb{D}}$ .

**NB:**

- $n$  can index anything.
- Universal - any type of algorithm or computational model.
- Similarly, no random algorithms converging with probability  $> 1/2$ .

# Proof idea: Constructing an adversary

$$F_0: \text{rotation by } \pi, \text{Sp}(\mathcal{K}_{F_0}) = \{\pm 1\}$$

**Phase transition lemma:** Let  $X = \{x_1, \dots, x_N\}, Y = \{y_1, \dots, y_N\}$  be distinct points in annulus  $\mathcal{A} = \{x \in \mathbb{D} \mid 0 < R < \|x\| < r < 1\}$  with  $X \cap Y = \emptyset$ . There exists a measure-preserving homeomorphism  $H$  such that  $H$  acts as the identity on  $\mathbb{D} \setminus \mathcal{A}$  and  $H(y_j) = F_0(H(x_j)), j = 1, \dots, N$ .

*Conjugacy of data ( $x_j \rightarrow y_j$ ) with  $F_0$*

**Idea:** Use lemma to trick any algorithm into oscillating between spectra.

# Proof idea: Constructing an adversary

Suppose (for contradiction)  $\{\Gamma_n\}$  uses  $\mathcal{J}_F$ ,  $\lim_{n \rightarrow \infty} \Gamma_n(F) = \text{Sp}(\mathcal{K}_F) \quad \forall F \in \Omega_{\mathbb{D}}$ .

Build an **adversarial**  $F$  ...

$$\mathcal{J}_F = \{(x, y_m) \mid \|F(x) - y_m\| \leq 2^{-m}\}$$


# Proof idea: Constructing an adversary

Suppose (for contradiction)  $\{\Gamma_n\}$  uses  $\mathcal{T}_F$ ,  $\lim_{n \rightarrow \infty} \Gamma_n(F) = \text{Sp}(\mathcal{K}_F) \quad \forall F \in \Omega_{\mathbb{D}}$ .

Build an **adversarial**  $F$ ...

$$\widetilde{F}_1(r, \theta) = (r, \theta + \pi + \phi(r)), \text{supp}(\phi) \subset [1/4, 3/4]$$

$$\text{Sp}(\mathcal{K}_{\widetilde{F}_1}) = \mathbb{T} \text{ (unit circle).}$$



$$\mathcal{T}_F = \{(x, y_m) \mid \|F(x) - y_m\| \leq 2^{-m}\}$$

# Proof idea: Constructing an adversary

Suppose (for contradiction)  $\{\Gamma_n\}$  uses  $\mathcal{T}_F$ ,  $\lim_{n \rightarrow \infty} \Gamma_n(F) = \text{Sp}(\mathcal{K}_F) \quad \forall F \in \Omega_{\mathbb{D}}$ .

Build an **adversarial**  $F$ ...


$$\widetilde{F}_1(r, \theta) = (r, \theta + \pi + \phi(r)), \text{supp}(\phi) \subset [1/4, 3/4]$$

$$\text{Sp}(\mathcal{K}_{\widetilde{F}_1}) = \mathbb{T} \text{ (unit circle).}$$

$$\lim_{n \rightarrow \infty} \Gamma_n(\widetilde{F}_1) = \text{Sp}(\mathcal{K}_{\widetilde{F}_1}) \Rightarrow \exists n_1 \text{ s.t. } \text{dist}(i, \Gamma_{n_1}(\widetilde{F}_1)) \leq 1.$$

**BUT**  $\Gamma_{n_1}$  uses finite amount of info to output  $\Gamma_{n_1}(\widetilde{F}_1)$ .

Let  $X, Y$  correspond to these snapshots.



$$\mathcal{T}_F = \{(x, y_m) \mid \|F(x) - y_m\| \leq 2^{-m}\}$$

# Proof idea: Constructing an adversary

Suppose (for contradiction)  $\{\Gamma_n\}$  uses  $\mathcal{T}_F$ ,  $\lim_{n \rightarrow \infty} \Gamma_n(F) = \text{Sp}(\mathcal{K}_F) \quad \forall F \in \Omega_{\mathbb{D}}$ .

Build an **adversarial**  $F$ ...

$$\widetilde{F}_1(r, \theta) = (r, \theta + \pi + \phi(r)), \text{supp}(\phi) \subset [1/4, 3/4]$$

$$\text{Sp}(\mathcal{K}_{\widetilde{F}_1}) = \mathbb{T} \text{ (unit circle).}$$

$$\lim_{n \rightarrow \infty} \Gamma_n(\widetilde{F}_1) = \text{Sp}(\mathcal{K}_{\widetilde{F}_1}) \Rightarrow \exists n_1 \text{ s.t. } \text{dist}(i, \Gamma_{n_1}(\widetilde{F}_1)) \leq 1.$$


**BUT**  $\Gamma_{n_1}$  uses finite amount of info to output  $\Gamma_{n_1}(\widetilde{F}_1)$ .

Let  $X, Y$  correspond to these snapshots.

Lemma:  $F_1 = H_1^{-1} \circ F_0 \circ H_1$  on annulus  $\mathcal{A}_1$ .

Consistent data  $\Rightarrow \Gamma_{n_1}(F_1) = \Gamma_{n_1}(\widetilde{F}_1)$ ,  $\text{dist}(i, \Gamma_{n_1}(F_1)) \leq 1$

**BUT**  $\text{Sp}(\mathcal{K}_{F_1}) = \text{Sp}(\mathcal{K}_{F_0}) = \{\pm 1\}$




# Proof idea: Constructing an adversary

**Inductive step:** Repeat on annuli,  $F_k = H_k^{-1} \circ F_0 \circ H_k$  on  $\mathcal{A}_k$ .  $F = \lim_{k \rightarrow \infty} F_k$

Consistent data  $\Rightarrow \Gamma_{n_k}(F) = \Gamma_{n_k}(\widetilde{F}_k)$ ,  $\text{dist}(i, \Gamma_{n_k}(F)) \leq 1$ ,  $n_k \rightarrow \infty$

**BUT**  $\text{Sp}(\mathcal{K}_F) = \text{Sp}(\mathcal{K}_{F_0}) = \{\pm 1\}$

**CANNOT CONVERGE**



Cascade of disks


# Proof idea: Constructing an adversary

**Inductive step:** Repeat on annuli,  $F_k = H_k^{-1} \circ F_0 \circ H_k$  on  $\mathcal{A}_k$ .  $F = \lim_{k \rightarrow \infty} F_k$

Consistent data  $\Rightarrow \Gamma_{n_k}(F) = \Gamma_{n_k}(\widetilde{F}_k)$ ,  $\text{dist}(i, \Gamma_{n_k}(F)) \leq 1$ ,  $n_k \rightarrow \infty$

**BUT**  $\text{Sp}(\mathcal{K}_F) = \text{Sp}(\mathcal{K}_{F_0}) = \{\pm 1\}$


**CANNOT CONVERGE**




# Classifications: *Solvability Complexity Index (SCI)*


**SCI:** Fewest number of limits needed to solve a computational problem.

- $\Delta_1$ : One limit, full error control. E.g.,  $d(\Gamma_n(F), \text{Sp}(\mathcal{K}_F)) \leq 2^{-n}$ .
- $\Delta_{m+1}$ :  $\text{SCI} \leq m$ .
- $\Sigma_m$ :  $\text{SCI} \leq m$ , final limit from below.

**trust output**  E.g.,  $\Sigma_1: \sup_{z \in \Gamma_n(F)} \text{dist}(z, \text{Sp}(\mathcal{K}_F)) \leq 2^{-n}$ .

•  $\Pi_m$ :  $\text{SCI} \leq m$ , final limit from above.  
E.g.,  $\Pi_1: \sup_{z \in \text{Sp}(\mathcal{K}_F)} \text{dist}(z, \Gamma_n(F)) \leq 2^{-n}$ .

 **verification**


 **covers spectrum**

- Hansen, "On the solvability complexity index, the n-pseudospectrum and approximations of spectra of operators." *J. Am. Math. Soc.*, 2011.
- C., "The foundations of infinite-dimensional spectral computations," **PhD diss.**, University of Cambridge, 2020.
- C., Hansen, "The foundations of spectral computations via the solvability complexity index hierarchy," *J. Eur. Math. Soc.*, 2022.
- C., Antun, Hansen, "The difficulty of computing stable and accurate neural networks," *Proc. Natl. Acad. Sci. USA*, 2022.
- Ben-Artzi, C., Hansen, Nevanlinna, Seidel, "On the solvability complexity index hierarchy and towers of algorithms," arXiv, 2020.

# Classification for Koopman

Lower + upper bounds

3 limits needed in general!



Different classes:

$$\Omega_{\mathcal{X}} = \{F: \mathcal{X} \rightarrow \mathcal{X} \mid F \text{ cts}\}$$

$$\Omega_{\mathcal{X}}^m = \{F: \mathcal{X} \rightarrow \mathcal{X} \mid F \text{ cts, m. p.}\}$$

$$\Omega_{\mathcal{X}}^{\alpha} = \{F: \mathcal{X} \rightarrow \mathcal{X} \mid F \text{ mod. cty. } \alpha\}$$

$$[d_{\mathcal{X}}(F(x), F(y)) \leq \alpha(d_{\mathcal{X}}(x, y))]$$

Optimal algorithms and classifications of dynamical systems.

## **Johann Wolfgang von Goethe:**

“Mathematicians are like Frenchmen: whatever you say to them they translate into their own language and forthwith it is something entirely different.”

Let's change the space...

# Reproducing kernel Hilbert space (RKHS)


Hilbert space of functions on  $\mathcal{X}$  s.t.  $g \mapsto g(x)$  bounded  $\forall x \in \mathcal{X}$ .

Generated by a kernel  $\mathfrak{K}: \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{C}$

$$g(x) = \langle g, \mathfrak{K}_x \rangle, \quad \mathfrak{K}(x, y) = \langle \mathfrak{K}_x, \mathfrak{K}_y \rangle = \mathfrak{K}_x(y)$$

## Advantages over $L^2(\mathcal{X}, \omega)$ :

- Forecasts: space bounds  $\Rightarrow$  pointwise bounds.
- High-dimensional systems practical through kernel trick.
- Fast methods for evaluating  $\mathfrak{K}$ .
- Different  $\mathfrak{K} \Rightarrow$  different  $\mathcal{K}$ ! Can be tailored to application. (This is where the community is currently heading.)
- Couples  $M$  and  $N$ ...



# SpecRKHS: Avoiding large data limit $M \rightarrow \infty$

Look at “Left eigenpairs” through  $\mathcal{K}^*$ :

$$\mathcal{K}^* \mathfrak{K}_x = \mathfrak{K}_{F(x)}$$

Evolution of functionals.  
 $g(x) = \langle g, \mathfrak{K}_x \rangle_{\mathcal{H}}$

No quadrature needed:

$$G_{jk} = \langle \mathfrak{K}_{x^{(k)}}, \mathfrak{K}_{x^{(j)}} \rangle = \mathfrak{K}(x^{(k)}, x^{(j)})$$

$$A_{jk} = \langle \mathcal{K}^* \mathfrak{K}_{x^{(k)}}, \mathfrak{K}_{x^{(j)}} \rangle = \langle \mathfrak{K}_{y^{(k)}}, \mathfrak{K}_{x^{(j)}} \rangle = \mathfrak{K}(y^{(k)}, x^{(j)})$$

$$L_{jk} = \langle \mathcal{K}^* \mathfrak{K}_{x^{(k)}}, \mathcal{K}^* \mathfrak{K}_{x^{(j)}} \rangle = \langle \mathfrak{K}_{y^{(k)}}, \mathfrak{K}_{y^{(j)}} \rangle = \mathfrak{K}(y^{(k)}, y^{(j)})$$

$$g = \sum_{m=1}^M \mathbf{g}_m \mathfrak{K}_{x^{(m)}}, \quad \|\mathcal{K}^* g - \lambda g\|_{\mathcal{H}}^2 = \mathbf{g}^* (L - \lambda A^* - \bar{\lambda} A + G) \mathbf{g}$$

# SpecRKHS: Example algorithm

$$\text{res}^*(\lambda, \mathbf{g})^2 = \frac{\|\mathcal{K}^* g - \lambda g\|_{\mathcal{H}}^2}{\|g\|_{\mathcal{H}}^2} = \frac{\mathbf{g}^* [L - \lambda A^* - \bar{\lambda} A + |\lambda|^2 G] \mathbf{g}}{\mathbf{g}^* G \mathbf{g}}$$


1. Compute  $G, A, L \in \mathbb{C}^{N \times N}$  ( $N = M$ )
2. For  $z_k$  in grid, compute  $\tau_k = \min_{g = \sum_{m=1}^N \mathbf{g}_m \mathfrak{K}_x(m)}$   $\text{res}^*(z_k, \mathbf{g})$ , corresponding  $g_k$  (gen. SVD).
3. **Output:**  $\{z_k: \tau_k < \varepsilon\}, \{g_k: \tau_k < \varepsilon\}$  ( $\varepsilon$ -pseudoeigenfunctions).

## Theorem:


- **Error control:**  $\{z_k: \tau_k < \varepsilon\} \subseteq \text{Sp}_\varepsilon(\mathcal{K}^*)$
- **Convergence:** Converges locally uniformly to  $\text{Sp}_\varepsilon(\mathcal{K}^*)$  (as  $N \rightarrow \infty$ )

$$\text{Sp}_\varepsilon(\mathcal{K}^*) = \{z \in \mathbb{C}: \exists g, \|g\|_{\mathcal{H}} = 1, \|\mathcal{K}^* g - z g\|_{\mathcal{H}} \leq \varepsilon\}$$

# Optimal algorithms and classifications of systems



# Optimal algorithms and classifications of systems

$\varepsilon = 0$  

increasing difficulty 

**Error control**

**1 limit**

**2 limits**

**3 limits**

$L_2(x, \omega)$




$$\Omega_x^\alpha = \{F: \text{mod. cty. } \alpha\}$$

$$d(F(x), F(y)) \leq \alpha(d(x, y))$$

$$\Omega_x^m \cap \Omega_x^\alpha$$


$$\Omega_x^m = \{F : F \text{ cts, m. p.}\}$$

$$\Omega_x = \{F : F \text{ cts}\}$$





# Optimal algorithms and classifications of systems




# Practical gains: Arctic sea ice forecasting



Satellite data




- Problems:**
1. Very hard to locate geographical significant regions.
  2. Very hard to predict more than two months in advance.



- C., Mezić, Stepanenko, "Adversarial dynamical systems characterize when data-driven learning succeeds or fails," *Nature Com.*, 2026.

# Avoid spurious evals $\Rightarrow$ State-of-the-art forecasts



$$g(x_n) = [\mathcal{K}^n g](x_0)$$

$$\|\mathcal{K}g - \lambda g\| \leq \varepsilon$$

$$\Rightarrow g(x_n) = \lambda^n g(x_0) + \mathcal{O}(n\varepsilon)$$

**Use  $\varepsilon$  to filter evals!**

**Figure: Mean binary accuracy over test years 2012-2020.**

(IceNet: Andersson et al, "Seasonal Arctic sea ice forecasting with probabilistic deep learning." *Nature Communications*, 2021.)

**What about forecast bounds and dictionary learning?**

# Principle Angle Decomposition (PAD)

**Algorithm 3.1** Principal angles and observables between  $\mathcal{V}$  and  $\mathcal{KV}$ .

**Input:** Matrices  $\mathbf{G}$ ,  $\mathbf{A}$  and  $\mathbf{L}$  in (2.3) and (2.9) for dictionary  $\{\psi_j\}_{j=1}^N$ , subspace selection matrix  $\mathbf{B} \in \mathbb{C}^{N \times n}$ , cut-off tolerance  $\epsilon_c \geq 0$ . For RKHS, use instead the matrices  $\mathbf{G}$ ,  $\mathbf{A}$  and  $\mathbf{R}$  (instead of  $\mathbf{L}$ ) from (2.6) and (2.10).

- 1: Compress the matrices to form  $\mathbf{G}_{\mathcal{V}} = \mathbf{B}^* \mathbf{G} \mathbf{B}$ ,  $\mathbf{A}_{\mathcal{V}} = \mathbf{B}^* \mathbf{A} \mathbf{B}$ , and  $\mathbf{L}_{\mathcal{V}} = \mathbf{B}^* \mathbf{L} \mathbf{B}$ .
- 2: Compute the matrix

$$\mathbf{J}_{\mathcal{V}} = \begin{pmatrix} \mathbf{G}_{\mathcal{V}} & \mathbf{A}_{\mathcal{V}} \\ \mathbf{A}_{\mathcal{V}}^* & \mathbf{L}_{\mathcal{V}} \end{pmatrix} \in \mathbb{C}^{2n \times 2n},$$

and its eigendecomposition  $\mathbf{J}_{\mathcal{V}} = \mathbf{U} \mathbf{D} \mathbf{U}^*$ .

- 3: Set  $\mathcal{I} = \{j : \mathbf{D}_{jj} > \epsilon_c, j = 1, \dots, 2n\}$ , and compute

$$\mathbf{C}_1 = \sqrt{\mathbf{D}(\mathcal{I}, \mathcal{I})} (\mathbf{U}(1:n, \mathcal{I}))^*, \quad \mathbf{C}_2 = \sqrt{\mathbf{D}(\mathcal{I}, \mathcal{I})} (\mathbf{U}(n+1:2n, \mathcal{I}))^*.$$

- 4: Compute the principal angles and observables

$$\{[\theta_j]_{j=1}^q, \mathbf{U}_1, \mathbf{U}_2\} = \text{subspacea}(\mathbf{C}_1, \mathbf{C}_2), \quad q \leq n,$$

and convert back to the original dictionary via (3.1) and (3.2).



**Output:** Principal angles  $\{\theta_j\}_{j=1}^q$  and principal observables  $\{(u_j, v_j)\}_{j=1}^q$ .

**Algorithm 3.2** Principal Angle Decomposition (PAD).

**Input:** Matrices  $\mathbf{G}$ ,  $\mathbf{A}$ ,  $\mathbf{K}$  and  $\mathbf{L}$  in (2.3), (2.4), and (2.9) for dictionary  $\{\psi_j\}_{j=1}^N$ , cut-off tolerance  $\epsilon_c \geq 0$ , number of post-compression modes  $r > 0$ .

- 1: Compute principal angles  $\{\theta_j\}_{j=1}^q$  and observables  $\mathbf{U}_1$  via Algorithm 3.1 ( $\mathbf{B} = \mathbf{I}_N$ ).
- 2: Define  $\mathbf{U}'_1 = \mathbf{U}_1(1:q, :) + \mathbf{K} \mathbf{U}_1(q+1:2q, :)$ , and truncate to  $\mathbf{U} = \mathbf{U}'_1(:, 1:r)$ .
- 3: Construct the matrices  $\mathbf{A}_{\text{pad}} = \mathbf{U}^* \mathbf{A} \mathbf{U} = \mathbf{K}_{\text{pad}}$  and  $\mathbf{L}_{\text{pad}} = \mathbf{U}^* \mathbf{L} \mathbf{U}$ .

**Output:** The compressed matrices  $\mathbf{A}_{\text{pad}}$ ,  $\mathbf{K}_{\text{pad}}$  and  $\mathbf{L}_{\text{pad}}$ .



# Principle Angle Decomposition (PAD)

**Algorithm 3.1** Principal angles and observables between  $\mathcal{V}$  and  $\mathcal{KV}$ .

**Input:** Matrices  $\mathbf{G}$ ,  $\mathbf{A}$  and  $\mathbf{L}$  in (2.3) and (2.9) for dictionary  $\{\psi_j\}_{j=1}^N$ , subspace selection matrix  $\mathbf{B} \in \mathbb{C}^{N \times n}$ , cut-off tolerance  $\epsilon_c \geq 0$ . For RKHS, use instead the matrices  $\mathbf{G}$ ,  $\mathbf{A}$  and  $\mathbf{R}$  (instead of  $\mathbf{L}$ ) from (2.6) and (2.10).

- 1: Compress the matrices to form  $\mathbf{G}_{\mathcal{V}} = \mathbf{B}^* \mathbf{G} \mathbf{B}$ ,  $\mathbf{A}_{\mathcal{V}} = \mathbf{B}^* \mathbf{A} \mathbf{B}$ , and  $\mathbf{L}_{\mathcal{V}} = \mathbf{B}^* \mathbf{L} \mathbf{B}$ .
- 2: Compute the matrix

$$\mathbf{J}_{\mathcal{V}} = \begin{pmatrix} \mathbf{G}_{\mathcal{V}} & \mathbf{A}_{\mathcal{V}} \\ \mathbf{A}_{\mathcal{V}}^* & \mathbf{L}_{\mathcal{V}} \end{pmatrix} \in \mathbb{C}^{2n \times 2n},$$

and its eigendecomposition  $\mathbf{J}_{\mathcal{V}} = \mathbf{U} \mathbf{D} \mathbf{U}^*$ .

- 3: Set  $\mathcal{I} = \{j : \mathbf{D}_{jj} > \epsilon_c, j = 1, \dots, 2n\}$ , and compute

$$\mathbf{C}_1 = \sqrt{\mathbf{D}(\mathcal{I}, \mathcal{I})} (\mathbf{U}(1:n, \mathcal{I}))^*, \quad \mathbf{C}_2 = \sqrt{\mathbf{D}(\mathcal{I}, \mathcal{I})} (\mathbf{U}(n+1:2n, \mathcal{I}))^*.$$

- 4: Compute the principal angles and observables

$$\{[\theta_j]_{j=1}^q, \mathbf{U}_1, \mathbf{U}_2\} = \text{subspacea}(\mathbf{C}_1, \mathbf{C}_2), \quad q \leq n,$$

and convert back to the original dictionary via (3.1) and (3.2).




**Output:** Principal angles  $\{\theta_j\}_{j=1}^q$  and principal observables  $\{(u_j, v_j)\}_{j=1}^q$ .

**Algorithm 3.2** Principal Angle Decomposition (PAD).

**Input:** Matrices  $\mathbf{G}$ ,  $\mathbf{A}$ ,  $\mathbf{K}$  and  $\mathbf{L}$  in (2.3), (2.4), and (2.9) for dictionary  $\{\psi_j\}_{j=1}^N$ , cut-off tolerance  $\epsilon_c \geq 0$ , number of post-compression modes  $r > 0$ .

- 1: Compute principal angles  $\{\theta_j\}_{j=1}^q$  and observables  $\mathbf{U}_1$  via Algorithm 3.1 ( $\mathbf{B} = \mathbf{I}_N$ ).
- 2: Define  $\mathbf{U}'_1 = \mathbf{U}_1(1:q, :) + \mathbf{K} \mathbf{U}_1(q+1:2q, :)$ , and truncate to  $\mathbf{U} = \mathbf{U}'_1(:, 1:r)$ .
- 3: Construct the matrices  $\mathbf{A}_{\text{pad}} = \mathbf{U}^* \mathbf{A} \mathbf{U} = \mathbf{K}_{\text{pad}}$  and  $\mathbf{L}_{\text{pad}} = \mathbf{U}^* \mathbf{L} \mathbf{U}$ .

**Output:** The compressed matrices  $\mathbf{A}_{\text{pad}}$ ,  $\mathbf{K}_{\text{pad}}$  and  $\mathbf{L}_{\text{pad}}$ .




# Dictionary learning and forecast multistep bounds

- Pluto and its 5 satellites
- Comparable mass of Pluto and Charon → **complex dynamics**
- Obtain error bounds for both  $L^2$  and RKHS cases


apply to dictionary refinement

obtain pointwise error bounds, expected error surrogates



# Dictionary learning and forecast bounds

- Apply algorithms to RKHS setting using Matérn kernel




# Dictionary learning and forecast bounds


- Apply algorithms to RKHS setting using Matérn kernel
- Expected errors using **Gaussian processes** avoid overestimation

- E.g., replace  $\|Ax\| \leq \|A\|\|x\|$  with  $\|Ax\| \approx \mathbb{E}_{\mathcal{C}}[A]\|x\|$  where

$$\mathbb{E}_{\mathcal{C}}[A] = \mathbb{E} \left[ \frac{\|Ax\|}{\|x\|} \right], \quad x \sim \mathcal{GP}(0, \mathcal{C})$$



# Dictionary learning and forecast bounds





# Shameless plug 1: CUP book out August 2026...

MATTHEW J. COLBROOK

## Infinite-Dimensional Spectral Computations

Foundations, Algorithms, and Modern Applications



### Contents

|   |       |
|---|-------|
| <i>Foreword by Wilhelm Schlag</i>   | ix    |
| <i>Preface</i>  | xiii  |
| <i>Notation</i>   | xxiii |
| <i>Example Classifications for Spectral Sets</i>                                | xxvii |
| <b>1 Spectral Problems in Infinite Dimensions</b>                               | 1     |
| <b>2 The Solvability Complexity Index: A Toolkit for Classifying Problems</b>   | 41    |
| <b>3 Computing Spectra with Error Control</b>                                   | 77    |
| <b>4 Spectral Measures of Self-Adjoint Operators</b>                            | 151   |
| <b>5 Spectral Measures of Unitary Operators</b>                                 | 219   |
| <b>6 Spectral Types of Self-Adjoint and Unitary Operators</b>                   | 261   |
| <b>7 Quantifying the Size of Spectra</b>  | 293   |
| <b>8 Essential Spectra</b>  | 345   |
| <b>9 Spectral Radii, Abscissas, and Gaps</b>                                    | 381   |
| <b>10 Nonlinear Spectral Problems</b>   | 416   |
| <b>11 Data-Driven Koopman Spectral Problems for Nonlinear Dynamical Systems</b> | 472   |
| <i>Appendix A Some Brief Preliminaries</i>                                      | 573   |
| <i>Appendix B A Bluffer's Guide to the SCI Hierarchy</i>                        | 590   |
| <i>Bibliography</i>   | 592   |
| <i>Index</i>  | 657   |

**100s of:**

- Classifications
- Algorithms
- Examples (full code)
- Exercises (full solutions)

**If something of interest – speak to me!**

*This talk*

# Shameless plug 2: Issac Newton Institute Programme <sup>50</sup>

## Operator Methods for Dynamical Systems


OMD

3 August 2026 to 28 August 2026

### Programme theme

Dynamical systems lie at the heart of our understanding of complex phenomena, whether in modelling weather and ocean currents, molecular dynamics, population growth or stock market fluctuations. However, the nonlinear nature of systems poses a challenge to describing their behaviour. Operator methods offer a powerful way to tackle this challenge. By representing a finite-dimensional nonlinear system's evolution as a linear operator acting on an infinite-dimensional space of functions, tools from linear algebra and spectral analysis can be used to gain insights into the system's long-term behaviour. This operator-based perspective has roots in 20th-century mathematics. It has proven fruitful in classical settings like statistical mechanics, where it helped connect chaotic systems with well-understood linear techniques.

In recent years, interest in operator methods for dynamical systems has surged, driven by data-driven techniques to approximate and analyse these operators from real-world or simulated data. A wide range of frameworks and algorithms have emerged, creating an exciting opportunity and need to develop a unifying foundation for these approaches. This programme aims to bring together experts and young researchers from various communities who use operator-theoretic perspectives (for example, the Koopman and transfer operator frameworks) to study dynamics. By uniting participants from pure theory to practical applications, the programme will spark new collaborations and jointly tackle key open questions, building a more cohesive research community. In particular, a significant focus will be on exploring the spectral properties of these operators—essentially, understanding their eigenvalues, modes, and related features—as these provide crucial insights into a system's long-term behaviour and how such behaviour can be effectively analysed and computed.



### Organisers

- Matthew Colbrook *University of Cambridge*
- Gary Froyland *University of New South Wales*
- Nathan Kutz *University of Washington*
- Julia Slipantschuk *University of Warwick; Universität Bayreuth*
- Caroline Wornell *University of Sydney*

### Participants

- Wael Bahsoun *Loughborough University*
- Steve Brunton *University of Washington*
- Christopher Budd *University of Bath; Institute of Mathematics and its Applications*

Visit to find out more about operator theory and machine learning in dynamical systems!

(and to enjoy Cambridge in the summer!)

# Pointers

1. Data-driven spectral problems for Koopman operators are hugely popular.  
**BUT: Standard truncation methods can fail – NEED TO GO INF-DIM!**
2. **General methods with convergence for spectral properties**  
 (spectra, pseudospectra, spectral measures, etc.) of K. operators!  
*E.g., Verification of approximate eigenfunctions leads to practical gains.*
3. **SCI hierarchy** classifies computational problems:  
**Lower bounds** through method of adversarial dynamics.  
**Upper bounds**  $\implies$  new “inf.-dim.” algorithms. Rigorous, optimal, practical.  
 **$\longrightarrow$  We now have a near complete spectral picture for K. on  $L^2(\mathcal{X}, \omega)$  and RKHS!**
4. **Matrix  $L$**  leads to forecast bounds, ways to train dictionary, and PAD.

# Lots of SCI upper bounds lurking in Koopman literature!

**SCI:** Fewest number of limits needed to solve a computational problem.

| Algorithm                           | Comments/Assumptions  | Spectral Problem's Corresponding SCI Upper Bound |                       |  |   |
|-------------------------------------|---|--|-----------------------|--|---|
|                                     |   | <i>KMD</i>                                       | <i>Spectrum</i>       | <i>Spectral Measure (if m.p.)</i>  | <i>Spectral Type (if m.p.)</i>                                |
| Extended DMD [47]                   | general $L^2$ spaces  | $\text{SCI} \leq 2^*$                            | N/C                   | N/C  | n/a   |
| Residual DMD [44]                   | general $L^2$ spaces  | $\text{SCI} \leq 2^*$                            | $\text{SCI} \leq 3^*$ | $\text{SCI} \leq 2^*$  | varies, see [84]<br>e.g., a.c. density: $\text{SCI} \leq 2^*$ |
| Measure-preserving EDMD [45]        | m.p. systems  | $\text{SCI} \leq 1$                              | N/C                   | $\text{SCI} \leq 2^*$ (general)<br>$\text{SCI} \leq 1$ (delay-embedding) | n/a   |
| Hankel DMD [85]                     | m.p. ergodic systems  | $\text{SCI} \leq 2^*$                            | N/C                   | N/C  | n/a   |
| Periodic approximations [86]        | m.p. + $\omega$ a.c.  | $\text{SCI} \leq 2$                              | N/C                   | $\text{SCI} \leq 2$ (see [87])   | a.c. density: $\text{SCI} \leq 3$                             |
| Christoffel–Darboux kernel [40]     | m.p. ergodic systems  | $\text{SCI} \leq 3$                              | n/a                   | $\text{SCI} \leq 2$  | e.g., a.c. density: $\text{SCI} \leq 2$                       |
| Generator EDMD [88]                 | cts.-time, samples $\nabla F$<br>(otherwise additional limit) | $\text{SCI} \leq 2$                              | N/C                   | $\text{SCI} \leq 2$ (see [89])   | n/a   |
| Compactification [42]               | cts.-time, m.p. ergodic systems                               | $\text{SCI} \leq 4$                              | N/C                   | $\text{SCI} \leq 4$  | n/a   |
| Resolvent compactification [43]     | cts.-time, m.p. ergodic systems                               | $\text{SCI} \leq 5$                              | N/C                   | $\text{SCI} \leq 5$  | n/a   |
| Diffusion maps [90] (see also [10]) | cts.-time, m.p. ergodic systems                               | $\text{SCI} \leq 3$                              | n/a                   | n/a  | n/a   |

**Are these sharp?**

**Previous techniques prove upper bounds on SCI.**

“N/C”: method need not converge. “n/a”: algorithm not applicable to problem.

Also in Ulam’s method for Markov processes, SRB measure computation, control,...

# References

- [1] Colbrook, Matthew J., and Alex Townsend. "Rigorous data-driven computation of spectral properties of Koopman operators for dynamical systems." *Communications on Pure and Applied Mathematics* 77.1 (2024): 221-283.
- [2] Colbrook, Matthew J., Loma J. Ayton, and Máté Szóke. "Residual dynamic mode decomposition: robust and verified Koopmanism." *Journal of Fluid Mechanics* 955 (2023): A21.
- [3] Colbrook, M. J., Li, Q., Raut, R. V., & Townsend, A. "Beyond expectations: residual dynamic mode decomposition and variance for stochastic dynamical systems." *Nonlinear Dynamics* 112.3 (2024): 2037-2061.
- [4] Colbrook, Matthew J. "The Multiverse of Dynamic Mode Decomposition Algorithms." *Handbook of Numerical Analysis*, vol. 25, pp. 127-230. Elsevier, 2024..
- [5] Colbrook, Matthew J. "The mpEDMD algorithm for data-driven computations of measure-preserving dynamical systems." *SIAM Journal on Numerical Analysis* 61.3 (2023): 1585-1608.
- [6] Colbrook, Matthew J., Catherine Drysdale, and Andrew Horning. "Rigged Dynamic Mode Decomposition: Data-Driven Generalized Eigenfunction Decompositions for Koopman Operators." *SIAM Journal on Applied Dynamical Systems* 24, no. 2 (2025): 1150-1190.
- [7] Boullé, Nicolas, and Matthew J. Colbrook. "Multiplicative Dynamic Mode Decomposition." *SIAM Journal on Applied Dynamical Systems* 24, no. 2 (2025): 1945-1968.
- [8] Boullé, Nicolas and Matthew J. Colbrook, "On the Convergence of Hermitian Dynamic Mode Decomposition" *Physica D: Nonlinear Phenomena*, 472 (2025).
- [9] Colbrook, Matthew J., Andrew Horning, and Tianyiwa Xie. "Computing Generalized Eigenfunctions in Rigged Hilbert Spaces." *arXiv preprint arXiv:2410.08343* (2024).
- [10] Zagli, Niccolò, et al. "Bridging the Gap between Koopmanism and Response Theory: Using Natural Variability to Predict Forced Response." *arXiv preprint arXiv:2410.01622* (2024).
- [11] Colbrook, Matthew J. "Another look at Residual Dynamic Mode Decomposition in the regime of fewer Snapshots than Dictionary Size." *Physica D: Nonlinear Phenomena* 469 (2024).
- [12] Colbrook, Matthew. "The foundations of infinite-dimensional spectral computations." *Diss. University of Cambridge*, 2020.
- [13] Ben-Artzi, J., Colbrook, M. J., Hansen, A. C., Nevanlinna, O., & Seidel, M. (2020). "Computing Spectra—On the Solvability Complexity Index Hierarchy and Towers of Algorithms." *arXiv preprint arXiv:1508.03280*.
- [14] Colbrook, Matthew J., Vegard Antun, and Anders C. Hansen. "The difficulty of computing stable and accurate neural networks: On the barriers of deep learning and Smale's 18th problem." *Proceedings of the National Academy of Sciences* 119.12 (2022): e2107151119.
- [15] Colbrook, Matthew, Andrew Horning, and Alex Townsend. "Computing spectral measures of self-adjoint operators." *SIAM review* 63.3 (2021): 489-524.
- [16] Colbrook, Matthew J., Bogdan Roman, and Anders C. Hansen. "How to compute spectra with error control." *Physical Review Letters* 122.25 (2019): 250201.
- [17] Colbrook, Matthew J., and Anders C. Hansen. "The foundations of spectral computations via the solvability complexity index hierarchy." *Journal of the European Mathematical Society* (2022).
- [18] Colbrook, Matthew J. "Computing spectral measures and spectral types." *Communications in Mathematical Physics* 384 (2021): 433-501.
- [19] Colbrook, Matthew J., and Anders C. Hansen. "On the infinite-dimensional QR algorithm." *Numerische Mathematik* 143 (2019): 17-83.
- [20] Colbrook, Matthew J. "On the computation of geometric features of spectra of linear operators on Hilbert spaces." *Foundations of Computational Mathematics* (2022): 1-82.
- [21] Brunton, Steven L., and Matthew J. Colbrook. "Resilient Data-driven Dynamical Systems with Koopman: An Infinite-dimensional Numerical Analysis Perspective."
- [22] Colbrook, Matthew J., Igor Mezić, and Alexei Stepanenko. "Limits and Powers of Koopman Learning." *arXiv preprint arxiv:2407.06312* (2024).
- [23] Herwig, April, Matthew J. Colbrook, Oliver Junge, Péter Koltai, and Julia Slipantschuk. "Avoiding spectral pollution for transfer operators using residuals." *arXiv preprint arXiv:2507.16915* (2025).
- [24] Boullé, Nicolas, Matthew J. Colbrook, and Gustav Conradie. "Convergent Methods for Koopman Operators on Reproducing Kernel Hilbert Spaces." *arXiv preprint arXiv:2506.15782* (2025).
- [25] Drysdale, Catherine, Matthew Colbrook, and Michael TM Woodley. "Computation and Verification of Spectra for Non-Hermitian Systems." *Physical review letters* 135.17 (2025): 170202.
- [26] Colbrook, Matthew J., and Catherine Drysdale. "Universal Methods for Nonlinear Spectral Problems." *arXiv preprint arXiv:2504.17012* (2025).
- [27] Colbrook, Matthew J., Zlatko Drmač, and Andrew Horning. "An Introductory Guide to Koopman Learning." *arXiv preprint arXiv:2510.22002* (2025).