Fast computation of orthogonal systems with a skew-symmetric differentiation matrix

Arieh Iserles

Department of Applied Mathematics and Theoretical Physics Centre for Mathematical Sciences University of Cambridge Wilberforce Rd, Cambridge CB4 1LE United Kingdom

> Marcus Webb* Department of Computer Science KU Leuven Celestijnenlaan 200A, 3001 Leuven Belgium

> > January 24, 2019

Abstract

Orthogonal systems in $L_2(\mathbb{R})$, once implemented in spectral methods, enjoy a number of important advantages if their differentiation matrix is skew-symmetric and highly structured. Such systems, where the differentiation matrix is skewsymmetric, tridiagonal and irreducible, have been recently fully characterised. In this paper we go a step further, imposing the extra requirement of fast computation: specifically, that the first N coefficients can be computed to high accuracy in $\mathcal{O}(N \log_2 N)$ operations. We consider two settings, one approximating a function f directly in $(-\infty,\infty)$ and the other approximating [f(x) + f(-x)]/2 and [f(x) - f(-x)]/2 separately in $[0, \infty)$. In each setting we prove that there is a single family, parametrised by $\alpha, \beta > -1$, of orthogonal systems with a skewsymmetric, tridiagonal, irreducible differentiation matrix and whose coefficients can be computed as Jacobi polynomial coefficients of a modified function. The four special cases where $\alpha, \beta = \pm 1/2$ are of particular interest, since coefficients can be computed using fast sine and cosine transforms. Banded, Toeplitz-plus-Hankel multiplication operators are also possible for representing variable coefficients in a spectral method. In Fourier space these orthogonal systems are related to an apparently new generalisation of the Carlitz polynomials.

 $^{^{*}\}mathrm{This}$ author is also affiliated with Research Foundation – Flanders (FWO) as a postdoctoral research fellow.

1 Introduction

This paper continues a project which we have commenced in (Iserles & Webb 2019b), to investigate complete systems in $L_2(\mathbb{R})$ with a skew-symmetric differentiation matrix. Let us begin by explaining briefly the underlying concepts, motivating our work and revisiting in a very abbreviated manner the contents of (Iserles & Webb 2019b).

The concept of a *differentiation matrix* originates in the theory of finite differences — essentially, this is the matrix taking a vector of function values to those of an approximation of the first derivative. An elementary example,

$$\frac{1}{2\Delta x} \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ -1 & 0 & 1 & \ddots & \vdots \\ 0 & \ddots & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & \ddots & 1 \\ 0 & \cdots & 0 & -1 & 0 \end{bmatrix},$$
(1.1)

is obtained from a second-order central difference approximation on a uniform grid with spacing $\Delta x > 0$ — note that it is skew-symmetric. Skew-symmetry mimics the self-adjointness of the first derivative operator in the standard L₂ Hilbert space with either zero Dirichlet or periodic boundary conditions, and it confers a wide range of advantages on a numerical method. We refer to (Hairer & Iserles 2016, Iserles 2014, Iserles 2016) for a wealth of specific examples: in essence, once a differentiation matrix is skew symmetric, it is often easy to prove stability for linear PDEs, as well as conservation of energy whenever it is mandated by the underlying equation.

The matrix in (1.1) is skew symmetric, yet it is a sobering thought that this second-order approximation of the derivative is as good as it gets: no skew-symmetric finite-difference differentiation matrix on a uniform grid may exceed order 2 (Iserles 2014). It is possible (but not easy) to obtain higher-order differentiation matrices of this kind carefully choosing specific non-uniform grids (Hairer & Iserles 2016, Hairer & Iserles 2017), but this is far from easy for high orders and, at any rate, finite differences are not the approach of choice in this paper.

An example that motivates much of our work is the *linear Schrödinger equation* in a semiclassical regime,

$$i\varepsilon \frac{\partial u}{\partial t} = \varepsilon^2 \frac{\partial^2 u}{\partial x^2} - V(x, t)u, \qquad a \le x \le b, \quad t \ge 0,$$
(1.2)

(for simplicity, we focus here on the univariate case), typically given with periodic boundary conditions. The parameter $\varepsilon > 0$ is small and V is the potential energy (Jin, Markowich & Sparber 2011). Typically, (1.2) is solved by a spectral method (or spectral collocation) in space, followed by a combination of splittings (e.g. Strang splitting or Zassenhaus splitting) and, in the case of time-dependent potential, the Magnus expansion or a version thereof in time (Bader, Iserles, Kropielnicka & Singh 2016, Blanes, Casas & Thalhammer 2017, Iserles, Kropielnicka & Singh 2018).

The definition of a differentiation matrix extends naturally into the realm of spectral methods. In any spectral method we expand the unknown in a (truncated) orthonormal basis $\{\varphi_m\}_{m\in\mathbb{Z}_+}$ of the underlying L₂ Hilbert space, where the φ_m s are suitably regular. The (infinite-dimensional) linear map \mathcal{D} taking $\{\varphi_m\}_{m\in\mathbb{Z}_+}$ into $\{\varphi'_m\}_{m\in\mathbb{Z}_+}$ is called a *differentiation matrix*. In other words, $\varphi'_m = \sum_{k=0}^{\infty} \mathcal{D}_{m,k}\varphi_k$, $m \in \mathbb{Z}_+$. The virtues of skew-symmetry remain undimmed in this setting and they immediately imply stability and energy conservation for (1.2).

Because of periodic boundary conditions, it is natural to use a Fourier basis for (1.2), leading to a skew-Hermitian (and diagonal) differentiation matrix.¹ To all intents and purposes, a skew-Hermitian matrix shares the benefits of a skew-symmetric one. Moreover, a Fourier basis has another critical virtue: we can approximate the first N expansion coefficients using FFT in just $\mathcal{O}(N \log_2 N)$ operations. Therefore, there is little incentive to seek alternative bases.

However, the traditional setting – finite interval, periodic boundary conditions – of (1.2) and other dispersive equations of quantum mechanics is being increasingly challenged, since it is inappropriate for long-term integration (or even short-term integration in the presence of quantum scattering) because sooner or later, wave packets reach a boundary and periodicity leads to non-physical solutions. Modern applications of quantum mechanics, not least quantum control (Shapiro & Brumer 2003), are an important example of this behaviour. Arguably, the most natural setting for (1.2) is that of a Cauchy problem on the entire real line, with solutions confined to $L_2(\mathbb{R})$. Zero Dirichlet boundary conditions on an interval correspond to a *closed* quantum system which can be interpreted as a quantum system on the real line with u(x, 0) = 0for $x \notin (a, b)$ or where the potential V is sufficiently large at the endpoints that no tunnelling is possible (Patriarca 1994).

In (Iserles & Webb 2019*b*) we have developed a comprehensive theory of orthonormal bases of $L_2(\mathbb{R})$ with a skew-symmetric, tridiagonal, irreducible differentiation matrix

$$\mathcal{D} = \begin{bmatrix} 0 & b_0 & 0 & & \\ -b_0 & 0 & b_1 & 0 & \\ 0 & -b_1 & 0 & b_2 & 0 \\ & \ddots & \ddots & \ddots & \ddots \end{bmatrix},$$
(1.3)

where without loss of generality $b_m > 0$, $m \in \mathbb{Z}$. There exists a one-to-one relationship between absolutely continuous real Borel measures $d\mu$ with all their moments bounded, symmetric with respect to the origin and supported by the entire real line on the one hand, orthonormal bases $\Phi = \{\varphi_m\}_{m \in \mathbb{Z}_+}$, complete in $L_2(\mathbb{R})$ and with a differentiation matrix of the form (1.3) on the other.

Specifically, let $d\mu(\xi) = w(\xi) d\xi$ be such Borel measure and $\{p_m\}_{m \in \mathbb{Z}_+}$ the underlying monic orthogonal polynomials. Because of symmetry, they obey a three-term recurrence relation of the form

$$p_{m+1}(x) = x p_m(x) - \lambda_m p_{m-1}(x), \qquad m \in \mathbb{Z}_+,$$
 (1.4)

where $\lambda_0 = 0$ and $\lambda_m > 0$, $m \in \mathbb{N}$. Set $b_m = \sqrt{\lambda_{m+1}}$, $m \in \mathbb{Z}_+$, and

$$\varphi_m(x) = \frac{(-i)^n}{\sqrt{2\pi}} \frac{1}{\|p_m\|} \int_{-\infty}^{\infty} g(\xi) p_m(\xi) e^{-ix\xi} d\xi, \qquad m \in \mathbb{Z}_+,$$
(1.5)

 $^{^1\}mathrm{We}$ typically index a Fourier basis by integers, rather than non-negative integers, but this causes no difficulty in our framework.

where $||p_m||^2 = \int_{-\infty}^{\infty} p_m^2(x) d\mu(x)$ and g is a complex-valued function with even real part and odd imaginary part such that $|g(\xi)|^2 = w(\xi)$. It has been proved in (Iserles & Webb 2019*b*) that $\{\varphi_m\}_{m \in \mathbb{Z}_+}$ is indeed orthonormal and complete in $L_2(\mathbb{R})$ (essentially due to the Plancharel theorem) and has the differentiation matrix (1.3). On the other hand, given such a basis, using the Favard Theorem we can always associate it with a symmetric Borel measure supported by the real line: such measure is unique if the corresponding Hamburger moment problem is determinate (Chihara 1978, p. 73).

Realistic implementation of the aforementioned bases, however, requires that two computations can be performed explicitly, namely explicit formation of the Orthogonal Polynomial System (OPS) $\{p_m\}_{m\in\mathbb{Z}_+}$ given $d\mu$ and the integrals (1.5), since otherwise we will not have the basis Φ in an explicit form. Here we are rapidly running against the limitations of current theory of orthogonal polynomials. An obvious and familiar example of Hermite functions

$$\varphi_m(x) = \frac{(-1)^m}{(2^m m!)^{1/2} \pi^{1/4}} e^{-x^2/2} H_m(x), \quad b_m = \left(\frac{m+1}{2}\right)^{1/2}, \qquad m \in \mathbb{Z}_+,$$

where the H_m are Hermite polynomials, aside, very few symmetric measures supported by the entire real line are known for which the underlying OPS (or even the recurrence coefficients λ_m) can be written explicitly: Iserles & Webb (2019*b*) list just the generalised Hermite polynomials and Carlitz polynomials. Indeed, the resolution of the *Freud problem*, precise asymptotic determination of the λ_m s for $w(\xi) = e^{-\xi^n + q(\xi)}$, where $n \geq 2$ is an even integer and q an even polynomial of degree $\leq n - 2$, using the Riemann-Hilbert transform (Deift, Kriecherbauer, McLaughlin, Venakides & Zhou 1999, Fokas, Its & Kitaev 1992), is considered a recent triumph of the theory of orthogonal polynomials – and in our setting we need explicit values, not just asymptotics! Thus, our basic challenge in this paper is to search for systems Φ by alternative means.

We seek systems that confer a tangible advantage when compared to the default of Hermite functions, and in this paper we find systems that allow for faster computation of expansion coefficients via Fast Cosine Transforms (FCT) or Fast Sine Transforms (FST). Given $f \in L_2(\mathbb{R}) \cap C^{\infty}(\mathbb{R})$, we wish to compute

$$\hat{f}_m = \int_{-\infty}^{\infty} f(x)\varphi_m(x) \,\mathrm{d}x, \qquad m = 0, \dots, N-1,$$

to sufficiently high accuracy in the least possible number of operations. We can identify three options.

Firstly, it is possible to use the *fast multipole method* to compute expansion coefficients in $\mathcal{O}(N \log N + N \log \varepsilon^{-1})$ operations to accuracy ε , provided that a table of nodes and weights of a suitable Gauss–Christoffel quadrature is available (Dutt, Gu & Rokhlin 1996). In the Hermite case, Gauss–Hermite quadrature is required and we note that its nodes and weights can be computed in $\mathcal{O}(N)$ operations (Townsend, Trogdon & Olver 2016). For other bases of the form (1.5) quadrature rules may take up to $\mathcal{O}(N^2)$ operations to compute.

Secondly, by the Fourier-transform on $L_2(\mathbb{R})$, the evaluation of the \hat{f}_m s reduces to an expansion in the polynomial basis $\{p_m\}_{m\in\mathbb{Z}_+}$. This, however, confers an advantage only if we can compute the polynomial coefficients in a fast manner. Of all fast and stable methodologies known to the present authors, this restricts us to $d\mu$ with bounded support, which has been ruled out earlier. The reason why we require $\sup \mu = \mathbb{R}$ is the completeness of the expansion: if, without loss of generality, $\sup \mu = (-1, 1)$ then Φ is complete not in $L_2(\mathbb{R})$ but in the *Paley–Wiener space* $\mathcal{PW}_{[-1,1]}(\mathbb{R})$ (Iserles & Webb 2019b). This is of no clear interest in the design of spectral methods for PDEs but might be relevant, for example, to signal processing.

In this paper we follow a third option. We seek a basis of φ_m s so that the expansion coefficients \hat{f}_m are equal to standard orthogonal polynomial coefficients of a modified function. If the standard polynomials are, for example, the Chebyshev polynomials (which a priori is not necessarily even possible) then the first N coefficients can be approximated in $\mathcal{O}(N \log_2 N)$ operations using the FCT. In Section 2, this line of reasoning leads us to consider the model,

$$\varphi_m(x) = \Theta(x)q_m(H(x)), \qquad m \in \mathbb{Z}_+, \tag{1.6}$$

where $\Theta \in L_2(\mathbb{R})$, $\{q_m\}_{m \in \mathbb{Z}_+}$ is an orthonormal OPS with respect to a measure W(t) dt supported in (-1, 1), and H is a differentiable, strictly monotonically increasing function mapping $(-\infty, \infty)$ onto (-1, 1).

We prove in Section 2 that enforcing orthonormality on such a model means that the coefficients are equal to the coefficients of $f(h(t))/\Theta(h(t))$ in the $\{q_m\}_{m\in\mathbb{Z}_+}$ basis, where h is the inverse function of H. When we further require that our system has the differentiation matrix in (1.3), all options, up to an affine change of variables, are whittled down to

$$\varphi_m^{(\alpha,\beta)}(x) = \frac{(-1)^m}{\sqrt{g_m^{(\alpha,\beta)}}} (1-X)^{\frac{\alpha+1}{2}} (1+X)^{\frac{\beta+1}{2}} \mathcal{P}_m^{(\alpha,\beta)}(X), \qquad X = \tanh x, \qquad (1.7)$$

where $\alpha, \beta > -1$, $\mathbf{P}_m^{(\alpha,\beta)}$ is the *m*th Jacobi polynomial and

$$g_m^{(\alpha,\beta)} = \int_{-1}^1 (1-t)^{\alpha} (1+t)^{\beta} [\mathbf{P}_m^{(\alpha,\beta)}(t)]^2 \, \mathrm{d}t = \frac{2^{1+\alpha+\beta} \Gamma(1+\alpha+m) \Gamma(1+\beta+m)}{m! (1+\alpha+\beta+2m) \Gamma(1+\alpha+\beta+m)}$$
(1.8)

(Rainville 1960, p. 260).

The first N expansion coefficients of a function with respect to Φ as in equation (1.7) can always be approximated in $\mathcal{O}(N(\log N)^2)$ operations by fast polynomial transform techniques (Townsend, Webb & Olver 2018). The cases $\alpha, \beta = \pm 1$ correspond to the Chebyshev polynomials of four different kinds (see (Olver, Lozier, Boisvert & Clark 2010, 18.3)), all of whose coefficients can be approximated in $\mathcal{O}(N \log_2 N)$ operations via various flavours of FCT and FST. Furthermore, a given expansion $f(x) = \sum_{k=0}^{N-1} c_k \varphi_k^{(\alpha,\beta)}(x)$ can be evaluated at a single point $x \in \mathbb{R}$ using Clenshaw's algorithm (Clenshaw 1955).

In Section 3 we turn our gaze to a more complicated model,

$$\varphi_{2m}(x) = \Theta_{\mathcal{E}}(x)r_m(H(x)), \qquad (1.9)$$
$$\varphi_{2m+1}(x) = \Theta_{\mathcal{O}}(x)s_m(H(x)), \qquad m \in \mathbb{Z}_+.$$

Here $\Theta_{\rm E}$ and $\Theta_{\rm O}$ are given functions, the first even and the second odd, $\{r_m\}_{m\in\mathbb{Z}_+}$ and $\{s_m\}_{m\in\mathbb{Z}_+}$ are orthonormal OPSs with respect to the measures $w_{\rm E}dt$ and $w_{\rm O}dt$, both supported by (-1, 1), and H maps strictly monotonically $(0, \infty)$ to (-1, 1). Neither $w_{\rm E}$ nor $w_{\rm O}$ need be symmetric with respect to the origin. Advancing along similar lines to our analysis of (1.6), computation of coefficients corresponding to the system (1.7) can be reduced to the expansion in the two OPS, $\{r_m\}_{m\in\mathbb{Z}_+}$ and $\{s_m\}_{m\in\mathbb{Z}_+}$.

The exploration of (1.9) is a lengthier process. Step after step we winnow the possibilities for H, $\Theta_{\rm E}$, $\Theta_{\rm O}$, $w_{\rm E}$ and $w_{\rm O}$ – at the end we are left with just a one-parameter family of such systems. We prove that (1.9) is consistent with orthonormality and with a skew-symmetric, tridiagonal, irreducible differentiation matrix if and only if $w_{\rm E}(t) = (1-t)^{\alpha}(1+1)^{-\frac{1}{2}}$ and $w_{\rm O}(t) = (1-t)^{\alpha}(1+t)^{\frac{1}{2}}$ for some $\alpha > -1$ – in other words, the r_m s and s_m s are (normalised) Jacobi polynomials $P_m^{(\alpha,-\frac{1}{2})}$ and $P_m^{(\alpha,\frac{1}{2})}$ respectively. However, looking deeper, we demonstrate that these systems are *identical* to the full-range system (1.7) with $\alpha = \beta$, implemented in a half-range mode.

Why do we need to discuss both the 'full-length' systems of Section 2 and the 'two half-length' systems of Section 3, in particular as the latter are nothing but an alternative implementation of the former? As things stand, they exhibit the same range of benefits: they are orthonormal and complete in $L_2(\mathbb{R})$, have skew-symmetric, tridiagonal differentiation matrices, while their expansion coefficients can be approximated in a fast and stable manner. We touch upon the advantages of either approach in Section 5. However, these are early days in the investigation of orthogonal systems with skew-symmetric differentiation matrices and their applications to spectral methods. There is a great merit, we believe, in exploring their theory, potential and limitations in the broadest possible sense.

In Section 4 we give the functions g and measures $d\mu(\xi)$ which appear in the formula (1.5) for these orthonormal bases, for all $\alpha, \beta > -1$. They turn out to be the weights for a generalised version of the Carlitz polynomials (now with two parameters) discussed in (Iserles & Webb 2019*b*), which have the explicit form,

$$d\mu_{\alpha,\beta}(\xi) = C_{\alpha,\beta}^2 \left| \Gamma\left(\frac{\alpha+1}{2} + \frac{i\xi}{2}\right) \Gamma\left(\frac{\beta+1}{2} - \frac{i\xi}{2}\right) \right|^2 d\xi, \qquad \alpha,\beta > -1.$$

The case $\beta = -\alpha$ corresponds to the one-parameter family of Carlitz polynomials. The original interest in Carlitz polynomials stems from the fact that their moments can be written in terms of Bernoulli polynomials. We are not aware of work on this two-parameter generalisation, nor can we anticipate its relevance to the theory of orthogonal polynomials on the real line.

In Section 5 we list some conclusions of our work, and flesh out the basis for applications of these functions to computing the Fourier transform and to a fast Olver–Townsend-type spectral method on the real line (Olver & Townsend 2013).

This is the place to mention (Iserles & Webb 2019*a*), a companion paper to the current one, where we explore complex-valued orthonormal systems with skew-Hermitian tridiagonal differentiation matrices. *Inter alia* we demonstrate there the existence of another system with coefficients that can be computed in $\mathcal{O}(N \log_2 N)$ operations, this time using the *Fast Fourier Transform*.

2 The full-range setting

In this section we characterise all orthonormal systems $\Phi = \{\varphi_m\}_{m \in \mathbb{Z}_+}$ with a skewsymmetric, irreducible, tridiagonal differentiation matrix, with the additional constraint that the expansion coefficients of a function $f \in L_2(\mathbb{R})$ are equal to expansion coefficients for a weighted and mapped version of f, in an orthonormal polynomial basis on (-1, 1). Explicitly, for the differentiation matrix, we require

$$\varphi'_m = -b_{m-1}\varphi_{m-1} + b_m\varphi_{m+1}, \qquad m \in \mathbb{Z}_+, \tag{2.1}$$

where $b_{-1} = 0$ and, without loss of generality, $b_m > 0$ for $m \in \mathbb{Z}_+$. For the expansion coefficients, we require

$$\hat{f}_m = \int_{-1}^1 \theta(t) f(h(t)) q_m(t) W(t) \, \mathrm{d}t, \qquad (2.2)$$

where $\{q_m\}_{m\in\mathbb{Z}}$ are orthonormal polynomials with respect to the absolutely continuous measure W(t) dt on (-1, 1), and $h, \theta : (-1, 1) \to \mathbb{R}$.

We will make the following assumptions about θ , h and W. Later we deduce much more indeed about these functions as necessary consequences of our model.

- h maps onto the whole of \mathbb{R} , is differentiable with h' being a measurable, positive function.
- This implies the existence of an inverse function $H : \mathbb{R} \to (-1, 1)$ which is differentiable with h'(t)H'(h(t)) = 1, therefore H' is also positive and measurable.
- θ is such that $t \mapsto \theta(t)\sqrt{W(t)/h'(t)} \in L_{\infty}(-1,1)$. A careful application of the weighted Cauchy-Schwarz inequality with weight h' shows that this implies that \hat{f}_m is finite for all $m \in \mathbb{Z}_+$.

Changing variables to x = h(t) yields

$$\hat{f}_m = \int_{-\infty}^{\infty} f(x)\Theta(x)q_m(H(x))\,\mathrm{d}x,\tag{2.3}$$

where $\Theta(x) = \theta(H(x))H'(x)W(H(x))$. For this to hold for all $f \in L_2(\mathbb{R})$, φ_m must be of the form,

$$\varphi_m(x) = \Theta(x)q_m(H(x)), \qquad m \in \mathbb{Z}_+.$$
(2.4)

The rest of the section is devoted to proving the following surprisingly simple result.

Theorem 1 All orthonormal systems of the form in equation (2.4) with a skewsymmetric, tridiagonal, irreducible differentiation matrix are, up to an affine change of variables, of the form

$$\varphi_m(x) = \frac{(-1)^m}{\sqrt{g_m^{(\alpha,\beta)}}} (1 - \tanh x)^{(\alpha+1)/2} (1 + \tanh x)^{(\beta+1)/2} \mathbf{P}_m^{(\alpha,\beta)}(\tanh x), \qquad m \in \mathbb{Z}_+$$
(2.5)

for $\alpha, \beta > -1$, with $g_m^{(\alpha,\beta)}$ as in (1.8). The coefficients b_m are given by (2.8).

We will discuss properties of these orthonormal systems in Section 4. In particular, we will show that they are in fact complete orthonormal bases for $L_2(\mathbb{R})$ by deriving the measure $d\mu(\xi)$ and function $g(\xi)$ in the Fourier transform expression which these systems must satisfy as in equation (1.5) and (Iserles & Webb 2019*b*).

2.1 Necessary conditions

Let us work out necessary consequences of Φ being an orthonormal system in $L_2(\mathbb{R})$. For all $n, m \in \mathbb{Z}_+$,

$$\int_{-\infty}^{\infty} \varphi_n(x)\varphi_m(x) \, \mathrm{d}x = \int_{-\infty}^{\infty} \Theta(x)^2 q_m(H(x))q_n(H(x)) \, \mathrm{d}x$$
$$= \int_{-1}^{1} q_m(t)q_n(t)\Theta(h(t))^2 h'(t) \, \mathrm{d}t.$$

It follows at once that

$$W(t) = \Theta(h(t))^2 h'(t) \quad \text{and} \quad \Theta(x) = \sqrt{H'(x)W(H(x))}.$$
(2.6)

We will return to this later.

It is considerably more complicated to ensure the existence of a skew-symmetric, tridiagonal differentiation matrix. First note that by setting m = 0 in equation (2.4), we see that Θ is infinitely differentiable and in $L_2(\mathbb{R})$, because it is proportional to φ_0 (which is an infinitely differentiable function in $L_2(\mathbb{R})$, being the Fourier transform of a superalgebraically decaying $L_2(\mathbb{R})$ function by (1.5)).

Inserting equation (2.4) into equation (2.1), we obtain for all $m \in \mathbb{Z}_+$,

$$\Theta'q_m(H) + \Theta q'_m(H)H' = -b_{m-1}\Theta q_{m-1}(H) + b_m\Theta q_{m+1}(H).$$

Setting m = 0 implies that $\Theta'(x) = b_0 q_1(H(x))\Theta(x)/q_0(H(x))$, which can be substituted back into the equation for general $m \in \mathbb{Z}_+$ to find

$$\Theta b_0 q_1(H) q_m(H) / q_0(H) + \Theta q'_m(H) H' = -b_{m-1} \Theta q_{m-1}(H) + b_m \Theta q_{m+1}(H).$$

Dividing through by $H'(x)\Theta(x)$ and changing variables to t = H(x) gives us

$$q'_{m}(t) = h'(t) \left[-b_{m-1}q_{m-1}(t) + b_{m}q_{m+1}(t) - \frac{b_{0}}{q_{0}}q_{1}(t)q_{m}(t) \right].$$
 (2.7)

Here we used the fact discussed earlier that H'(h(t))h'(t) = 1.

Now, the left-hand-side of equation (2.7) is a polynomial of degree exactly m-1, while the right-hand-side is h'(t) times a polynomial of degree at most m+1. It follows from the case m = 1 that $h'(t) = 1/(at^2 + bt + c)$ for some real numbers a, b and c. Since $h: (-1, 1) \to \mathbb{R}$ and h' > 0, necessarily $h'(\pm 1) = \infty$ and it follows that, up to an affine change of variables, $h(t) = \operatorname{arctanh} t$ and $H(x) = \operatorname{tanh} x$.

Back to equation (2.6). Substituting in the necessary form of h we have,

$$W(t) = \frac{\Theta(\operatorname{arctanh} t)^2}{1 - t^2}.$$

Using the fact that $\Theta'(x) = b_0 q_1(H(x))\Theta(x)/q_0(H(x))$, we conclude after simple algebra that

$$W'(t) = \frac{2b_0q_1(t)/q_0 + 2t}{1 - t^2}W(t).$$

Writing $2b_0q_1(t)/q_0 + 2t = (1+t)\alpha + (1-t)\beta$ for real numbers α and β , the solution to this ODE with W(0) = 1 without loss of generality, is

$$W(t) = (1-t)^{\alpha}(1+t)^{\beta}.$$

It is none other than than a *Jacobi weight*. We deduce that, if at all possible, the polynomials in the system (2.4) must be the (orthonormal) Jacobi polynomials,

$$q_m(t) = \frac{(-1)^{k_m}}{\sqrt{g_m^{(\alpha,\beta)}}} \mathcal{P}_m^{(\alpha,\beta)}(t)$$

where $g_m^{(\alpha,\beta)}$ has been defined in (1.8) and k_m is an integer — for the time being, to allow for simpler algebra, we assume that $k_m = 0$ but will change this later.

Substituting W(t) into equation (2.6) shows that

$$\Theta(x) = (1 - \tanh x)^{\frac{\alpha+1}{2}} (1 + \tanh x)^{\frac{\beta+1}{2}}.$$

2.2 Sufficient conditions

All that remains is to check whether these systems actually satisfy the requirements set out at the start of the section. Firstly, the functions are in $L_2(\mathbb{R})$ if and only if $\alpha, \beta > -1$ (which also ensures that W(t) dt is a finite measure). A quick change of variables will show that these functions are indeed orthonormal for all $\alpha, \beta > -1$. For the final requirement on the differentiation matrix, we have,

$$\begin{split} \varphi_m'(t) &= -\frac{\alpha+1}{2} \frac{1}{\sqrt{g_m^{(\alpha,\beta)}}} (1-t^2)(1-t)^{\frac{\alpha-1}{2}} (1+t)^{\frac{\beta+1}{2}} \mathbf{P}_m^{(\alpha,\beta)}(t) \\ &+ \frac{\beta+1}{2} \frac{1}{\sqrt{g_m^{(\alpha,\beta)}}} (1-t)^{\frac{\alpha+1}{2}} (1+t)^{\frac{\beta-1}{2}} \mathbf{P}_m^{(\alpha,\beta)}(t) \\ &+ \frac{1}{\sqrt{g_m^{(\alpha,\beta)}}} (1-t^2)(1-t)^{\frac{\alpha+1}{2}} (1+t)^{\frac{\beta+1}{2}} \frac{\mathrm{d}\mathbf{P}_m^{(\alpha,\beta)}(t)}{\mathrm{d}t} \\ &= \frac{(1-t)^{\frac{\alpha+1}{2}} (1+t)^{\frac{\beta+1}{2}}}{\sqrt{g_m^{(\alpha,\beta)}}} \left[\left(\frac{\beta-\alpha}{2} - \frac{\alpha+\beta+2}{2} t \right) \mathbf{P}_m^{(\alpha,\beta)}(t) + (1-t^2) \frac{\mathrm{d}\mathbf{P}_m^{(\alpha,\beta)}}{\mathrm{d}t} \right] \end{split}$$

According to (Olver et al. 2010, 18.9.17–18)

$$(1-t^2)\frac{\mathrm{d}\mathbf{P}_m^{(\alpha,\beta)}(t)}{\mathrm{d}t}(t) = m\left(\frac{\alpha-\beta}{\alpha+\beta+2m}-t\right)\mathbf{P}_m^{(\alpha,\beta)}(t) + \frac{2(\alpha+m)(\beta+m)}{\alpha+\beta+2m}\mathbf{P}_{m-1}^{(\alpha,\beta)}(t)$$

$$= (\alpha + \beta + m + 1) \left(\frac{\alpha - \beta}{\alpha + \beta + 2m + 2} + t \right) \mathbf{P}_{m-1}^{(\alpha,\beta)}(t)$$
$$= \frac{2(m+1)(\alpha + \beta + m + 1)}{\alpha + \beta + 2m + 2} \mathbf{P}_{m+1}^{(\alpha,\beta)}(t),$$

while the three-term recurrence relation for Jacobi polynomials is

$$t\mathbf{P}_m^{(\alpha,\beta)}(t) = A_m \mathbf{P}_{m-1}^{(\alpha,\beta)} + B_m \mathbf{P}_m^{(\alpha,\beta)}(t) + C_m \mathbf{P}_{m+1}^{(\alpha,\beta)}(t),$$

where

$$A_{m} = \frac{2(\alpha + m)(\beta + m)}{(\alpha + \beta + 2m)(\alpha + \beta + 2m + 1)},$$

$$B_{m} = -\frac{\alpha^{2} - \beta^{2}}{(\alpha + \beta + 2m)(\alpha + \beta + 2m + 2)},$$

$$C_{m} = \frac{2(m + 1)(\alpha + \beta + m + 1)}{(\alpha + \beta + 2m + 1)(\alpha + \beta + 2m + 2)}$$

(Rainville 1960, p. 263). In other words,

$$\varphi'_m(x) = \eta_m \varphi_{m-1}(x) + \theta_m \varphi_m(x) + b_m \varphi_{m+1}(x),$$

where

$$\eta_m = \sqrt{\frac{g_{m-1}^{(\alpha,\beta)}}{g_m^{(\alpha,\beta)}}} \left[-\frac{\alpha+\beta+2}{2} + \alpha+\beta+m+1 \right] A_m,$$

$$\theta_m = \frac{\beta-\alpha}{2} - \frac{\alpha+\beta+2}{2} B_m + m \left(\frac{\alpha-\beta}{\alpha+\beta+2m} - B_m\right),$$

$$b_m = \sqrt{\frac{g_{m+1}^{(\alpha,\beta)}}{g_m^{(\alpha,\beta)}}} \left(-\frac{\alpha+\beta+2}{2} - m \right) C_m.$$

We require that $\eta_m = -b_{m-1}$ and $\theta_m = 0$. However, before we look further into the above quantities, we note that necessarily $b_m < 0$. Fortunately, this is an artefact of our choice of $k_m = 0$: once we replace this with $k_m = m$, we obtain $b_m > 0$, $m \in \mathbb{Z}_+$, as required. This requires long, yet simple algebra, best performed using a symbolic algebra package. It follows from (1.8) that

$$\sqrt{\frac{g_{m-1}^{(\alpha,\beta)}}{g_m^{(\alpha,\beta)}}} = \sqrt{\frac{m(\alpha+\beta+m)(\alpha+\beta+2m+1)}{(\alpha+m)(\beta+m)(\alpha+\beta+2m-1)}},$$
$$\sqrt{\frac{g_{m+1}^{(\alpha,\beta)}}{g_m^{(\alpha,\beta)}}} = \sqrt{\frac{\alpha+m+1)(\beta+m+1)(\alpha+\beta+2m+1)}{(m+1)(\alpha+\beta+m+1)(\alpha+\beta+2m+3)}},$$

hence all the conditions are satisfied for

$$b_m = \sqrt{\frac{(m+1)(\alpha+m+1)(\beta+m+1)(\alpha+\beta+m+1)}{(\alpha+\beta+2m+1)(\alpha+\beta+2m+3)}}, \qquad m \in \mathbb{Z}_+.$$
 (2.8)

3 A half-range setting

In this section we take the model from the previous section a step further. Rather than requiring that our coefficients are equal to coefficients of a modified function in a single orthogonal polynomial basis, we consider two polynomial bases which are linked to the odd and even basis elements.

Specifically, suppose we have an orthonormal system Φ in $L_2(\mathbb{R})$ such that φ_m is an even function if m is even and an odd function if m is odd; let us call this an *even-odd system*. Then we can re-express the expansion coefficients in the form

$$\hat{f}_{2m} = 2 \int_0^\infty f_{\mathcal{E}}(x) \varphi_{2m}(x) \, \mathrm{d}x, \quad \hat{f}_{2m+1} = 2 \int_0^\infty f_{\mathcal{O}}(x) \varphi_{2m+1}(x) \, \mathrm{d}x, \qquad m \in \mathbb{Z}_+,$$

where

$$f_{\rm E}(x) = \frac{1}{2}[f(x) + f(-x)], \quad f_{\rm O}(x) = \frac{1}{2}[f(x) - f(-x)], \qquad x \in (0,\infty).$$

In other words, we can 'translate' the setting from the real line to $(0, \infty)$ and this forms the theme of this section. We suppose that the odd and even coefficients can each be expressed in a similar fashion to the previous section, but with separate expressions for each case, as follows. Let $\{r_m\}_{m\in\mathbb{Z}_+}$ and $\{s_m\}_{m\in\mathbb{Z}_+}$ be orthonormal polynomial systems with respect to the absolutely continuous measures $w_{\rm E}(t)dt$ and $w_{\rm O}(t)dt$, respectively, on (-1, 1). We wish to find all orthonormal, even-odd systems Φ with a tridiagonal, skew-symmetric differentiation matrix, i.e.

$$\varphi'_m = -b_{m-1}\varphi_{m-1} + b_m\varphi_{m+1}, \qquad m \in \mathbb{Z}_+, \tag{3.1}$$

and such that the coefficients are equal to

$$\hat{f}_{2m} = \int_{-1}^{1} \theta_{\rm E}(t) f_{\rm E}(h(t)) r_m(t) w_{\rm E}(t) \,\mathrm{d}t, \qquad (3.2)$$
$$\hat{f}_{2m+1} = \int_{-1}^{1} \theta_{\rm O}(t) f_{\rm O}(h(t)) s_m(t) w_{\rm O}(t) \,\mathrm{d}t, \qquad m \in \mathbb{Z}_+,$$

where $h: (-1,1) \to (0,\infty)$ and $\theta_{\rm E}, \theta_{\rm O}: (-1,1) \to \mathbb{R}$.

We will make the following assumptions about h, $\theta_{\rm E}$, $\theta_{\rm O}$, $w_{\rm E}$ and $w_{\rm O}$. Just as in the previous section, we will *deduce* more from these basic assumptions and our model.

- h maps onto the whole of $(0, \infty)$, is differentiable with h' a measurable function, and h'(t) > 0 for all $t \in (-1, 1)$.
- This implies the existence of an inverse function $H : (0, \infty) \to (-1, 1)$ which is differentiable with h'(t)H'(h(t)) = 1, which implies that H' is also positive and measurable.
- $\theta_{\rm E}$ is such that $t \mapsto \theta_{\rm E}(t) \sqrt{w_{\rm E}(t)/h'(t)} \in L_{\infty}(\mathbb{R})$, and $\theta_{\rm O}$ satisfies the analogous property. The motivation is exactly as in the previous section.

Changing variables to x = h(t) yields

$$\hat{f}_{2m} = \int_0^\infty \Theta_{\mathcal{E}}(x) f_{\mathcal{E}}(x) r_m(H(x)) \, \mathrm{d}x, \qquad (3.3)$$
$$\hat{f}_{2m+1} = \int_0^\infty \Theta_{\mathcal{O}}(x) f_{\mathcal{O}}(x) s_m(H(x)) \, \mathrm{d}x, \qquad m \in \mathbb{Z}_+,$$

where $\Theta_{\rm E}(x) = \theta_{\rm E}(H(x))H'(x)w_{\rm E}(H(x))$ and $\Theta_{\rm O}(x) = \theta_{\rm O}(H(x))H'(x)w_{\rm O}(H(x))$. For this to hold for all $f \in L_2(\mathbb{R})$, we must necessarily have the 'half-range model',

$$\varphi_{2m}(x) = \Theta_{\mathcal{E}}(x)r_m(H(x))$$

$$\varphi_{2m+1}(x) = \Theta_{\mathcal{O}}(x)s_m(H(x)), \qquad m \in \mathbb{Z}_+.$$
(3.4)

We extend H to the whole of \mathbb{R} by setting H(-x) = H(x). Since φ_0 is an even function, we must have that Θ_E is even function and likewise Θ_O is an odd function. Note that φ_m is an infinitely differentiable function for all $m \in \mathbb{Z}_+$ because it is the Fourier transform of a superalgebraically decaying function by equation (1.5). Therefore, $\varphi_1(0) = 0$ in order to have oddness. It follows that $\Theta_O(0) = 0$. Allow us to place one more assumption into the mix: $\Theta_E(0) \neq 0$. Otherwise all basis functions vanish at the origin, rendering it clearly unsuitable for approximation of functions which are in general *nonzero* at the origin.

The rest of the section is devoted to proving the following.

Theorem 2 All the systems (3.4) which are orthonormal in $L_2(\mathbb{R})$ and possess a tridiagonal, skew-symmetric differentiation matrix are, up to a linear change of variables,

$$\varphi_{2m}(x) = \frac{2^{(2\alpha+1)/4}}{\sqrt{g_m^{(\alpha,-\frac{1}{2})}}} \frac{1}{\cosh^{1+\alpha} x} \mathbf{P}_m^{(\alpha,-\frac{1}{2})} \left(1 - \frac{2}{\cosh^2 x}\right), \qquad (3.5)$$
$$\varphi_{2m+1}(x) = -\frac{2^{(2\alpha+3)/4}}{\sqrt{g_m^{(\alpha,\frac{1}{2})}}} \frac{\sinh x}{\cosh^{2+\alpha} x} \mathbf{P}_m^{(\alpha,\frac{1}{2})} \left(1 - \frac{2}{\cosh^2 x}\right), \qquad m \in \mathbb{Z}_+,$$

for any $\alpha > -1$.

We show that these bases are equal to the bases in Theorem 1 with $\beta = \alpha$, something which is far from obvious at first sight. All discussions of mathematical properties of these functions such as completeness in $L_2(\mathbb{R})$ can therefore be derived from properties of the functions in Theorem 1.

3.1 Necessary conditions

The first condition to explore is orthonormality. Since the φ_m s have the parity of m on the real line, we have

$$\int_{-\infty}^{\infty} \varphi_{2m}(x)\varphi_{2n+1}(x) \,\mathrm{d}x = 0, \qquad m, n \in \mathbb{Z}_+$$

and need to check orthogonality only within each set. Changing variables, it follows from (3.4) that

$$\int_{-\infty}^{\infty} \varphi_{2m}(x)\varphi_{2n}(x) \,\mathrm{d}x = 2 \int_{0}^{\infty} \varphi_{2m}(x)\varphi_{2n}(x) \,\mathrm{d}x = 2 \int_{-1}^{1} \Theta_{\mathrm{E}}^{2}(h(t))h'(t)r_{m}(t)r_{n}(t) \,\mathrm{d}t.$$

Since $\{r_m\}_{m\in\mathbb{Z}_+}$ is an orthonormal set with respect to $w_{\rm E}(t)dt$, we must have

$$w_{\rm E}(t) = 2\Theta_{\rm E}^2(h(t))h'(t), \qquad t \in (-1,1),$$
(3.6)

and, by the same token,

$$w_{\rm O}(t) = 2\Theta_{\rm O}^2(h(t))h'(t), \qquad t \in (-1,1).$$
 (3.7)

Note that, by the monotonicity of h, both weight functions are nonnegative, as required. These expressions can be inverted: changing back to x,

$$\Theta_{\rm E}(x) = \sqrt{\frac{1}{2}H'(x)w_{\rm E}(H(x))}, \quad \Theta_{\rm O}(x) = \pm \sqrt{\frac{1}{2}H'(x)w_{\rm O}(H(x))}, \qquad x \in (0,\infty).$$
(3.8)

Observe that we need to be very careful in our choice of sign. As things stand, we allow for both options.

Our next, considerably more challenging task is to ensure the existence of a differentiation matrix of the correct form. Recall that both $\Theta_{\rm E}$ and $\Theta_{\rm O}$ are smooth functions on \mathbb{R} . Substituting (3.4) into (3.1), we have

$$\Theta'_{\mathrm{E}}r_{m}(H) + \Theta_{\mathrm{E}}H'r'_{m}(H) = \Theta_{\mathrm{O}}[b_{2m-1}s_{m-1}(H) + b_{2m}s_{m}(H)], \qquad (3.9)$$

$$\Theta'_{\mathrm{O}}s_{m}(H) + \Theta_{\mathrm{O}}H's'_{m}(H) = \Theta_{\mathrm{E}}[-b_{2m-1}r_{m}(H) + b_{2m+1}r_{m+1}(H)], \qquad m \in \mathbb{Z}_{+}.$$

Setting m = 0 yields

$$\Theta'_{\mathrm{E}}(x) = \Theta_{\mathrm{O}}(x)b_0s_0/r_0, \qquad (3.10)$$
$$\Theta'_{\mathrm{O}}(x) = \Theta_{\mathrm{E}}(x)b_1r_1(H(x))/s_0.$$

Substituting these back into (3.9) and changing variables back to t, we have,

$$r'_{m}(t) = h'(t) \sqrt{\frac{w_{\rm O}(t)}{w_{\rm E}(t)}} A_{m}(t)$$
(3.11)

$$s'_{m}(t) = h'(t)\sqrt{\frac{w_{\mathrm{E}}(t)}{w_{\mathrm{O}}(t)}}B_{m}(t), \qquad m \in \mathbb{N},$$
(3.12)

where

$$A_m(t) = -b_{2m-1}s_{m-1}(t) + b_{2m}s_m(t) - \frac{b_0s_0}{r_0}r_m(t),$$

$$B_m(t) = -b_{2m}r_m(t) + b_{2m+1}r_{m+1}(t) + \frac{b_0r_0 - b_1r_1(t)}{s_0}s_m(t), \qquad m \in \mathbb{N}.$$

The way forward rests upon the observation that the left-hand sides of both (3.11) and (3.12) are (m-1)-degree polynomials, and this places important constrains upon their right-hand sides. This is similar to the analysis of Section 2 yet considerably more complicated.

Setting m = 1, taking products of equations (3.11) and (3.12) and performing some simple algebra, we obtain,

$$h'(t) = \sqrt{\frac{r'_1 s'_1}{A_1(t)B_1(t)}}$$
 and $\frac{w_{\rm O}(t)}{w_{\rm E}(t)} = \frac{r'_1 B_1(t)}{s'_1 A_1(t)}.$

Since r_1 , s_1 , and A_1 are polynomials of degree 1 and B_1 is a polynomial of degree 2, there exist constants $a, b, c, \gamma_1, \gamma_2$ (which are real except that b and c may be complex conjugates) such that

$$h'(t) = \frac{\gamma_1}{\sqrt{(1-at)(1-bt)(1-ct)}} \quad \text{and} \quad \frac{w_{\rm O}(t)}{w_{\rm E}(t)} = \gamma_2 \frac{(1-bt)(1-ct)}{1-at}.$$
 (3.13)

Substituting the ratio of $w_{\rm O}(t)$ and $w_{\rm E}(t)$ for $t \in (-1, 1)$ into equations (3.6) and (3.7), we find

$$\Theta_{\rm O}(h(t)) = \sqrt{\gamma_2 \frac{(1-bt)(1-ct)}{1-at}} \Theta_{\rm E}(h(t)).$$
(3.14)

Since h(-1) = 0, $\Theta_{\rm O}(0) = 0$ and $\Theta_{\rm E}(0) \neq 0$ we deduce (without loss of generality) that c = -1. Since $\lim_{t\to 1} h(t) = +\infty$, we must have, integrating h'(t),

$$\lim_{t \to 1} \int_{-1}^{t} \frac{1}{\sqrt{(1-as)(1-bs)(1+s)}} \, \mathrm{d}s = +\infty.$$

This is only possible if a = b = 1. Therefore,

$$h'(t) = \frac{\gamma_1}{(1-t)\sqrt{1+t}}, \qquad w_{\rm O}(t) = \gamma_2(1+t)w_{\rm E}(t).$$

The formula for h'(t) is readily integrated using the substitution $\tanh u = \sqrt{(1+s)/2}$, giving

$$h(t) = \sqrt{2\gamma_1} \operatorname{arctanh} \sqrt{\frac{1+t}{2}}.$$

Inverting, and ignoring a linear change of variables in x, we have whittled our way down to a single option,

$$H(x) = 1 - \frac{2}{\cosh^2 x}.$$

Now let us find $\Theta_{\rm E}(x)$. We know that $\Theta'_{\rm E}(x) = \Theta_{\rm O}(x)b_0s_0r_0$ by equation (3.10) and $\Theta_{\rm O}(x) = \sqrt{\gamma_2(1+H(x))}\Theta_{\rm E}(x)$ by equation (3.14). Hence,

$$\frac{\mathrm{d}}{\mathrm{d}t}\Theta_{\mathrm{E}}(h(t)) = \Theta_{\mathrm{E}}'(h(t))h'(t) = \frac{\Theta_{E}(h(t))}{1-t}\sqrt{\gamma_{2}}\gamma_{1}b_{0}s_{0}/r_{0}.$$

In consequence, $\Theta_{\rm E}(h(t)) \propto (1-t)^{\frac{\alpha+1}{2}}$ for some $\alpha > -1$, and, converting to the x variable, we have

$$\Theta_{\rm E}(x) \propto \frac{1}{\cosh^{\alpha+1}(x)}, \qquad \Theta_{\rm O}(x) \propto \frac{\sinh(x)}{\cosh^{2+\alpha}(x)}.$$

Equations (3.6) and (3.7) give us the weights,

$$w_{\rm E}(t) = (1-t)^{\alpha}(1+t)^{-\frac{1}{2}}, \qquad w_{\rm O}(t) = (1-t)^{\alpha}(1+t)^{\frac{1}{2}},$$

where without loss of generality $w_{\rm E}(0) = w_{\rm O}(0) = 1$.

3.2 Sufficient conditions

As things stand, we have identified one – and just one – one-parameter family of weights $\{w_{\rm E}, w_{\rm O}\}$ for which we might be able to obtain an orthonormal system (3.4) with a tridiagonal skew-symmetric differentiation matrix:

$$\begin{split} w_{\rm E}(t) &= (1-t)^{\alpha} (1+t)^{-\frac{1}{2}}, \quad w_{\rm O}(t) = (1-t)^{\alpha} (1+t)^{\frac{1}{2}}, \qquad t \in (-1,1), \quad \alpha > -1, \\ r_m(t) &= \frac{1}{\sqrt{g_m^{(\alpha,-\frac{1}{2})}}} {\rm P}_m^{(\alpha,-\frac{1}{2})}(t), \quad s_m(t) = \frac{1}{\sqrt{g_m^{(\alpha,\frac{1}{2})}}} {\rm P}_m^{(\alpha,\frac{1}{2})}(t), \qquad m \in \mathbb{Z}_+, \\ h(t) &= \tanh^{-1} \sqrt{\frac{1+t}{2}}, \qquad H(x) = 1 - \frac{2}{\cosh^2 x}, \\ \Theta_{\rm E}(x) &= 2^{\frac{1}{4} + \frac{\alpha}{2}} \frac{1}{\cosh^{1+\alpha} x}, \qquad \Theta_{\rm O}(x) = -2^{\frac{3}{4} + \frac{\alpha}{2}} \frac{\sinh x}{\cosh^{2+\alpha} x}, \end{split}$$

where we have used (3.8) (with a minus sign) to determine $\Theta_{\rm E}$ and $\Theta_{\rm O}$. The question is, do the systems here actually satisfy our requirements for all $\alpha > -1$? The following subsection answers this question in the affirmative by relating these functions to the full-range systems from Section 2.

3.3 The connection to full-range systems

Further investigation of these half-range systems (which we omit here for the sake of brevity and lack of utility) leads one to the conclusion the half-range systems of Theorem 2 are a special case of full-range systems of Theorem 1 with $\beta = \alpha$. The formulæ look completely different, but as we will now show, they are identical.

For half-range functions

$$\varphi_{2m}(x) = \frac{2^{\frac{\alpha+1}{4}}}{\sqrt{g_m^{(\alpha,-\frac{1}{2})}}} \frac{1}{\cosh^{\alpha+1}x} \mathcal{P}_m^{(\alpha,-\frac{1}{2})} \left(1 - \frac{2}{\cosh^2 x}\right),$$

while for full range,

$$\varphi_{2m}(x) = \frac{1}{\sqrt{g_{2m}^{(\alpha,\alpha)}}} (1 - \tanh^2 x)^{\frac{\alpha+1}{2}} \mathcal{P}_{2m}^{(\alpha,\alpha)}(\tanh x)$$

$$=\frac{2^{\frac{\alpha+1}{2}}}{\sqrt{g_{2m}^{(\alpha,\alpha)}}}\frac{1}{\cosh^{\alpha+1}x}\mathbf{P}_{2m}^{(\alpha,\alpha)}\left(\sqrt{1-\frac{1}{\cosh^2 x}}\right).$$

We now evoke Proposition 5 (cf. Appendix A):

$$\mathbf{P}_{2m}^{(\alpha,\alpha)}\left(\sqrt{1-\frac{1}{\cosh^2 x}}\right) = \frac{(\alpha+m+1)!m!}{(2m)!}\mathbf{P}_m^{(\alpha,-\frac{1}{2})}\left(1-\frac{2}{\cosh^2 x}\right)$$

and, substituting the values of $g_m^{(\alpha,-\frac{1}{2})}$ and $g_{2m}^{(\alpha,\alpha)}$ from (1.8), we deduce that the two expressions are indeed identical.

Likewise, in a half-range formalism,

$$\varphi_{2m+1}(x) = -\frac{2^{\frac{\alpha}{2} + \frac{3}{4}}}{\sqrt{g_m^{(\alpha, \frac{1}{2})}}} \frac{\sin h}{\cosh^{2+\alpha} x} \mathcal{P}_m^{(\alpha, \frac{1}{2})} \left(1 - \frac{2}{\cosh^2 x}\right).$$

while the full-range expression is

$$\varphi_{2m+1}(x) = \frac{(-1)^m}{\sqrt{g_{2m+1}^{(\alpha,\alpha)}}} (1 - \tanh^2 x)^{\frac{\alpha+1}{2}} \mathcal{P}_{2m+1}^{(\alpha,\alpha)}(\tanh x)$$
$$= \frac{2^{\frac{\alpha+1}{2}}}{\sqrt{g_{2m+1}^{(\alpha,\alpha)}}} \frac{1}{\cosh^{\alpha+1} x} \mathcal{P}_{2m+1}^{(\alpha,\alpha)}\left(\sqrt{1 - \frac{1}{\cosh^2 x}}\right).$$

It follows from Proposition 6 and from

$$1 - \frac{1}{\cosh^2 x} = \tanh^2 x$$

that the two expressions are identical.

4 The tanh-Jacobi functions

In Section 2 we identified the following orthonormal bases $\Phi^{(\alpha,\beta)} = \{\varphi_m^{(\alpha,\beta)}\}_{m\in\mathbb{Z}_+}$ for $\alpha,\beta>-1$,

$$\varphi_m^{(\alpha,\beta)}(x) = \frac{(-1)^m}{\sqrt{g_m^{(\alpha,\beta)}}} (1-\tanh x)^{\frac{\alpha+1}{2}} (1+\tanh x)^{\frac{\beta+1}{2}} \mathcal{P}_m^{(\alpha,\beta)}(\tanh x), \qquad m \in \mathbb{Z}_+.$$

These orthonormal bases have a tridiagonal, irreducible, skew-symmetric differentiation matrix as in equation (1.3), with

$$b_m^{(\alpha,\beta)} = \sqrt{\frac{(m+1)(\alpha+m+1)(\beta+m+1)(\alpha+\beta+m+1)}{(\alpha+\beta+2m+1)(\alpha+\beta+2m+3)}}, \qquad m \in \mathbb{Z}_+.$$

We call these function the Tanh-Jacobi functions, for obvious reasons.



Figure 4.1: The first five basis functions for the Hermite functions (top-left), Tanh-Chebyshev-T functions (top-right), Tanh-Legendre functions (bottom-left) and the Tanh-Chebyshev-U functions (bottom-right). The darkest coloured line represents m = 0, progressing up to the lighted coloured, representing m = 4.

In Section 3 we showed that in the special case where $\beta = \alpha$, there is an alternative, equivalent expression which separates the functions into odd and even parts. For all $\alpha > -1$, let

$$\varphi_{2m}^{\alpha}(x) = \frac{2^{(2\alpha+1)/4}}{\sqrt{g_m^{(\alpha,-\frac{1}{2})}}} \frac{1}{\cosh^{1+\alpha} x} \mathbf{P}_m^{(\alpha,-\frac{1}{2})} \left(1 - \frac{2}{\cosh^2 x}\right),$$
$$\varphi_{2m+1}^{\alpha}(x) = -\frac{2^{(2\alpha+3)/4}}{\sqrt{g_m^{(\alpha,\frac{1}{2})}}} \frac{\sinh x}{\cosh^{2+\alpha} x} \mathbf{P}_m^{(\alpha,\frac{1}{2})} \left(1 - \frac{2}{\cosh^2 x}\right), \qquad m \in \mathbb{Z}_+.$$

Then $\varphi_m^{\alpha} = \varphi_m^{(\alpha,\alpha)}$ for all $m \in \mathbb{Z}_+$, at least mathematically speaking. For computation of coefficients, this basis is different, as is discussed below.

4.1 Expansion coefficients

The expansion coefficients of a function $f \in L_2(\mathbb{R})$ in the orthonormal basis $\Phi^{(\alpha,\beta)}$ are equal to

$$\hat{f}_m = \frac{(-1)^m}{\sqrt{g_m^{(\alpha,\beta)}}} \int_{-1}^1 \frac{f(\operatorname{arctanh} t)}{(1-t)^{\frac{\alpha+1}{2}}(1+t)^{\frac{\beta+1}{2}}} \mathcal{P}_m^{(\alpha,\beta)}(t)(1-t)^{\alpha}(1+t)^{\beta} \, \mathrm{d}t, \tag{4.1}$$

which are the Jacobi coefficients of the modified function

$$F(t) = \frac{f(\operatorname{arctanh} t)}{(1-t)^{\frac{\alpha+1}{2}}(1+t)^{\frac{\beta+1}{2}}}$$

with a diagonal scaling. The regularity of F determines the convergence of the coefficients, since Jacobi polynomials bases have spectral convergence properties (this is an elementary consequence of integration by parts and derivative identity (Olver et al. 2010, 18.19.6)).

For the half-range model, while the coefficients are equal to those given above with $\beta = \alpha$, they take the following form. Let

$$f_{\rm E}(x) = \frac{1}{2}[f(x) + f(-x)], \qquad f_{\rm O}(t) = \frac{1}{2}[f(x) - f(-x)], \qquad x \in (0,\infty),$$

be the even and odd portions of f, respectively. Then, using the transformation $x = h(t) = \operatorname{arctanh} \sqrt{\frac{1+t}{2}}$, we have

$$\hat{f}_{2m} = \int_{-\infty}^{\infty} f(x)\varphi_{2m}(x) \,\mathrm{d}x = 2 \int_{0}^{\infty} f_{\mathrm{E}}(x)\varphi_{2m}(x) \,\mathrm{d}x = 2 \int_{-1}^{1} f_{\mathrm{E}}(h(t))h'(t)\varphi_{2m}(h(t)) \,\mathrm{d}t$$
$$= \frac{2^{-\frac{1}{4}}}{\sqrt{g_{m}^{(\alpha,-\frac{1}{2})}}} \int_{-1}^{1} f_{\mathrm{E}}\left(\arctan\sqrt{\frac{1+t}{2}}\right) (1-t)^{-\frac{1}{2}(1-\alpha)}(1+t)^{-\frac{1}{2}} \mathrm{P}_{m}^{(\alpha,-\frac{1}{2})}(t) \,\mathrm{d}t$$
$$= \frac{2^{-\frac{1}{4}}}{\sqrt{g_{m}^{(\alpha,-\frac{1}{2})}}} \int_{-1}^{1} \frac{f_{\mathrm{E}}\left(\arctan\sqrt{\frac{1+t}{2}}\right)}{(1-t)^{(1+\alpha)/2}} \mathrm{P}_{m}^{(\alpha,-\frac{1}{2})}(t) w_{\mathrm{E}}(t) \,\mathrm{d}t.$$
(4.2)

Likewise,

$$\hat{f}_{2m+1} = \int_{-\infty}^{\infty} \varphi_{2m}(x) \, \mathrm{d}x = 2 \int_{0}^{\infty} f_{\mathrm{O}}(x) \varphi_{2m+1}(x) \, \mathrm{d}x$$
$$= 2 \int_{-1}^{1} f_{\mathrm{O}}(h(t)) h'(t) \varphi_{2m+1}(h(t)) \, \mathrm{d}t$$
$$= -\frac{2^{\frac{1}{4}}}{\sqrt{g_{m}^{(\alpha,\frac{1}{2})}}} \int_{-1}^{1} \frac{f_{\mathrm{O}}\left(\arctan\sqrt{\frac{1+t}{2}}\right)}{(1-t)^{(1+\alpha)/2}(1+t)^{\frac{1}{2}}} \mathbf{P}_{m}^{(\alpha,\frac{1}{2})}(t) w_{\mathrm{O}}(t) \, \mathrm{d}t.$$
(4.3)

Convergence of the coefficients is determined by the functions

$$F_{\rm E}(t) = \frac{f_{\rm E}\left(\operatorname{arctanh}\sqrt{\frac{1+t}{2}}\right)}{(1-t)^{(1+\alpha)/2}}, \qquad F_{\rm O}(t) = \frac{f_{\rm O}\left(\operatorname{arctanh}\sqrt{\frac{1+t}{2}}\right)}{(1-t)^{(1+\alpha)/2}(1+t)^{\frac{1}{2}}}.$$

Exactly how properties of the function f itself determine the rate of convergence is a topic for another paper. It is clear from the outset that it is some combination of regularity and decay at infinity for the function f which will determine the regularity of F — if for example f only decays algebraically at infinity, then F will be unbounded in (-1, 1)!

With regards to computation, the first N Jacobi coefficients of a given function can be approximated in $\mathcal{O}(N(\log N)^2)$ operations using fast polynomial transform techniques described in (Townsend et al. 2018). An efficient and straightforward Julia implementation exists in the software package, APPROXFUN (Olver 2018).

The cases where $\alpha, \beta = \pm \frac{1}{2}$ correspond to Chebyshev polynomials of various kinds. There are four kinds of Chebyshev polynomials:

- 1. Chebyshev polynomials of the first kind T_m , orthogonal in (-1,1) with the weight function $(1-t^2)^{-\frac{1}{2}}$;
- 2. Chebyshev polynomials of the second kind U_m , orthogonal in (-1, 1) with the weight function $(1 t^2)^{\frac{1}{2}}$;
- 3. Chebyshev polynomials of the third kind V_m , orthogonal in (-1,1) with the weight function $(1-t)^{\frac{1}{2}}(1+t)^{-\frac{1}{2}}$; and
- 4. Chebyshev polynomials of he fourth kind W_m , orthogonal in (-1,1) with the wights function $(1-t)^{-\frac{1}{2}}(1+t)^{\frac{1}{2}}$

(Olver et al. 2010, 18.3). They are all scaled Jacobi polynomials with $\alpha, \beta \in \{-\frac{1}{2}, +\frac{1}{2}\}$ and are expressible in terms of trigonometric functions,

$$T_m(\cos\theta) = \cos m\theta, \qquad U_m(\cos\theta) = \frac{\sin(m+1)\theta}{\sin\theta}, V_m(\cos\theta) = \frac{\sin(m+\frac{1}{2})\theta}{\sin\frac{1}{2}\theta}, \qquad W_m(\cos\theta) = \frac{\cos(m+\frac{1}{2})\theta}{\cos\frac{1}{2}\theta}$$

(Olver et al. 2010, 18.5.1–4). Therefore, letting $t = \cos \theta$, the integral in equation (4.1) can be converted into trigonometric integrals over $(0, \pi)$ which correspond to either Cosine or Sine Transform. Either can be discretised and computed in $\mathcal{O}(N \log_2 N)$ operations using Fast Cosine Transform (FCT) or Fast Sine Transform (FST).

Because of this algorithmic advantage, and odd-even symmetry, the bases $\Phi^{(\alpha,\beta)}$ with $(\alpha,\beta) = (-\frac{1}{2},-\frac{1}{2})$ and $(\alpha,\beta) = (\frac{1}{2},\frac{1}{2})$, are contenders for practical application. The Tanh-Chebyshev-T functions are given by

$$\varphi_m^{\left(-\frac{1}{2},-\frac{1}{2}\right)}(x) = (-1)^m \sqrt{2/\pi} \operatorname{sech}^{\frac{1}{2}} x \tilde{\mathrm{T}}_m(\tanh x), \qquad m \in \mathbb{Z}_+,$$
$$b_m^{\left(-\frac{1}{2},-\frac{1}{2}\right)} = \frac{1}{2} \left(m + \frac{1}{2}\right),$$

where $\tilde{T}_m = T_m$ if m > 0 and $\tilde{T}_0 = 1/\sqrt{2}$. The Tanh-Chebyshev-U functions are given by

$$\varphi_m^{\left(\frac{1}{2},\frac{1}{2}\right)}(x) = (-1)^m \sqrt{\frac{2}{\pi}} \operatorname{sech}^{\frac{3}{2}} x \operatorname{U}_m(\tanh x), \qquad m \in \mathbb{Z}_+,$$
$$b_m^{\left(\frac{1}{2},\frac{1}{2}\right)} = \frac{1}{2} \left(m + \frac{3}{2}\right).$$

For the half-range model, equations (4.2) and (4.3) can also be converted into Sine and Cosine transforms. Specifically, with $\alpha = -\frac{1}{2}$ we obtain the combination of Chebyshev polynomials of the first and the fourth kind, T_m and W_m , which can be computed with FCT-I and FCT-II respectively – cf. (Duhamel & Vetterli 1990) for different flavours of FCT and FST. Likewise, $\alpha = \frac{1}{2}$ results in a combination of U_m and V_m , computable with FST-I and FST-II respectively.

4.2 The Fourier transform representation and completeness

We will now prove that these bases are complete in $L_2(\mathbb{R})$. By (Iserles & Webb 2019b, Thms 6, 8) (see also equation (1.5)) there exists a complex-valued function $g_{\alpha,\beta}(\xi)$ with even real part and odd imaginary part such that

$$\varphi_m^{(\alpha,\beta)}(x) = \frac{(-\mathrm{i})^m}{\sqrt{2\pi}} \int_{-\infty}^{\infty} g_{\alpha,\beta}(\xi) p_m(\xi) \mathrm{e}^{\mathrm{i}x\xi} \,\mathrm{d}\xi, \tag{4.4}$$

where $P = \{p_m\}_{m \in \mathbb{Z}_+}$ are the orthonormal polynomials with respect to $d\mu(\xi) = w(\xi) d\xi = |g_{\alpha,\beta}(\xi)|^2 d\xi$. By (Iserles & Webb 2019*b*, Thm. 9), the functions $\Phi^{(\alpha,\beta)}$ are complete in $L_2(\mathbb{R})$ if $w(\xi) > 0$ for all $\xi \in \mathbb{R}$ and polynomials are dense in the space $L_2(\mathbb{R}, d\mu(\xi))$.

Now, applying the Fourier transform to both sides of equation (4.4) and setting m = 0, we have

$$g_{\alpha,\beta}(\xi) \propto \int_{-\infty}^{\infty} \varphi_0^{(\alpha,\beta)}(x) \mathrm{e}^{-\mathrm{i}x\xi} \mathrm{d}x.$$

We can normalise this function after we find an expression. Let us use the change of variables $\tau = e^x$ to manipulate this integral into a more reasonable form.

$$\begin{split} & \int_{-\infty}^{\infty} \varphi_0^{(\alpha,\beta)}(x) \mathrm{e}^{-\mathrm{i}x\xi} \, \mathrm{d}x \\ &= \int_{-\infty}^{\infty} (1 - \tanh x)^{\frac{\alpha+1}{2}} (1 + \tanh x)^{\frac{\beta+1}{2}} \mathrm{e}^{-\mathrm{i}x\xi} \, \mathrm{d}x \\ &= \int_{-\infty}^{\infty} \left(\frac{2\mathrm{e}^{-x}}{\mathrm{e}^x + \mathrm{e}^{-x}}\right)^{\frac{\alpha+1}{2}} \left(\frac{2\mathrm{e}^x}{\mathrm{e}^x + \mathrm{e}^{-x}}\right)^{\frac{\beta+1}{2}} \mathrm{e}^{-\mathrm{i}x\xi} \, \mathrm{d}x \\ &= 2^{\alpha+\beta+1} \int_{0}^{\infty} \frac{\tau^{-\frac{\alpha+1}{2}} \tau^{\frac{\beta+1}{2}} \tau^{-\mathrm{i}\xi}}{(\tau + \tau^{-1})^{\frac{\alpha+\beta}{2}+1}} \frac{\mathrm{d}\tau}{\tau} \\ &= 2^{\alpha+\beta+1} \int_{0}^{\infty} \frac{\tau^{\beta-\mathrm{i}\xi}}{(1 + \tau^2)^{\frac{\alpha+\beta}{2}+1}} \frac{\mathrm{d}\tau}{\tau}. \end{split}$$

The change of variables $\sigma = \tau^2$ transforms this into

$$\int_{-\infty}^{\infty} \varphi_0^{(\alpha,\beta)}(x) e^{-ix\xi} dx = 2^{\alpha+\beta} \int_0^{\infty} \frac{\sigma^{\frac{\beta-1}{2} - \frac{i\xi}{2}}}{(1+\sigma)^{\frac{\alpha+\beta}{2} + 1}} d\sigma$$

This is one of the standard integral formulae for Euler's Beta function (Olver et al. 2010, 5.12.3), meaning that

$$g_{\alpha,\beta}(\xi) \propto \mathbf{B}\left(\frac{\alpha+1}{2} + \frac{\mathrm{i}\xi}{2}, \frac{\beta+1}{2} - \frac{\mathrm{i}\xi}{2}\right).$$

Using the identity $B(a, b) = \Gamma(a)\Gamma(b)/\Gamma(a+b)$ for all $a, b \in \mathbb{C}$, we deduce the existence of a real constant $C_{\alpha,\beta}$ such that

$$g_{\alpha,\beta}(\xi) = C_{\alpha,\beta}\Gamma\left(\frac{\alpha+1}{2} + \frac{\mathrm{i}\xi}{2}\right)\Gamma\left(\frac{\beta+1}{2} - \frac{\mathrm{i}\xi}{2}\right).$$

Note that $g_{\alpha,\beta}$ is complex-valued in general, but has an even real part and odd imaginary part, which implies that $\varphi_m^{\alpha,\beta}(x)$ is real-valued for all $x \in \mathbb{R}$. Barnes's Beta Integral (Olver et al. 2010, 5.13.3) can be used to compute the constant $C_{\alpha,\beta}$ which makes the measure $d\mu_{\alpha,\beta}(\xi) = |g_{\alpha,\beta}(\xi)|^2 d\xi$ have mass equal to one.

We have thus proved the following theorem.

Theorem 3 The Fourier-space measure associated with $\{\varphi_m^{(\alpha,\beta)}\}_{m=0}^{\infty}$ is

$$d\mu_{\alpha,\beta}(\xi) = C_{\alpha,\beta}^2 \left| \Gamma\left(\frac{\alpha+1}{2} + \frac{i\xi}{2}\right) \Gamma\left(\frac{\beta+1}{2} - \frac{i\xi}{2}\right) \right|^2 d\xi, \qquad \alpha,\beta > -1.$$

There is little hope of simplifying the expression for $d\mu_{\alpha,\beta}$ from Theorem 3 for general $\alpha, \beta > -1$ except in some special cases.

Let us first consider the case $\beta = -\alpha$, where necessarily $\alpha \in (-1, 1)$. In this case, the reflection formula, $\Gamma(z)\Gamma(1-z) = \pi \operatorname{cosec} \pi z$ (Olver et al. 2010, 5.5.3), implies

$$g_{\alpha,-\alpha}(\xi) = \pi C_{\alpha,-\alpha} \frac{1}{\cos\left(\frac{\pi}{2}(\alpha + \mathrm{i}\xi)\right)}$$

and using the basic multiple angle formula, the associated measure is equal to

$$d\mu_{\alpha,-\alpha}(\xi) = 2\pi^2 C_{\alpha,-\alpha}^2 \frac{1}{\cosh(\pi\xi) + \cos(\pi\alpha)} d\xi.$$

The polynomials associated to this measure are known as the *Carlitz polynomials* (after mapping them from a line in the complex plane to the real line) (Carlitz 1959). They were mentioned in (Iserles & Webb 2019*b*), but now we have the full picture of their relationship to orthogonal systems with a tridiagonal, skew-symmetric differentiation matrix.

The case $\alpha = \beta$ yields

$$g_{\alpha,\alpha}(\xi) = C_{\alpha,\alpha} \left| \Gamma\left(\frac{\alpha+1}{2} + \frac{\mathrm{i}\xi}{2}\right) \right|^2,\tag{4.5}$$

since $\Gamma(\overline{z}) = \overline{\Gamma(z)}$.

Lemma 4 In the case $\alpha = \beta \ge 0$ being integers we have

$$g_{2n,2n}(\xi) = \pi C_{n,n} \frac{\prod_{j=0}^{n-1} [(j+\frac{1}{2})^2 + \frac{1}{4}\xi^2]}{\cosh(\frac{\pi\xi}{2})}, \qquad (4.6)$$
$$g_{2n+1,2n+1}(\xi) = \frac{\pi}{2} C_{2n+1,2n+1} \frac{\xi \prod_{j=1}^{n} (j^2 + \frac{1}{4}\xi^2)}{\sinh(\frac{\pi\xi}{2})}, \qquad n \in \mathbb{Z}_+.$$

Proof In the case n = 0 (4.6) is confirmed by direct computation. Otherwise we use the standard recurrence formula $\Gamma(z+1) = z\Gamma(z)$. (4.5) implies that

$$\frac{g_{n,n}(\xi)}{C_{n,n}} = \frac{(n-1)^2 + \xi^2}{4} \frac{g_{n-2,n-2}(\xi)}{C_{n-2,n-2}}, \qquad n \ge 2,$$

and (4.6) follows by simple induction.

Regarding completeness of $\Phi^{(\alpha,\beta)}$ in $L_2(\mathbb{R})$, as mentioned above, $g_{\alpha,\beta}$ being nonzero for all $\xi \in \mathbb{R}$ (because the Γ function has no roots in the complex plane), all that remains is the question of density of polynomials in the space $L_2(\mathbb{R}, d\mu)$. It would be sufficient that $w(\xi)$ have exponential decay as $\xi \to \pm \infty$ (Akhiezer 1965, pp. 45,86,86).

To show that these measures have exponential decay, we use the asymptotic formula (Olver et al. 2010, 5.11.9),

$$|\Gamma(x+iy)| \sim \sqrt{2\pi} |y|^{x-\frac{1}{2}} e^{-\pi|y|/2}$$
 as $|y| \to \infty$.

This implies,

$$\left|\Gamma\left(\frac{\alpha+1}{2}+\frac{i\xi}{2}\right)\Gamma\left(\frac{\beta+1}{2}-\frac{i\xi}{2}\right)\right|^2 \sim 4\pi^2 \left|\frac{\xi}{2}\right|^{\alpha+\beta} e^{-\pi\xi} \text{ as } |\xi| \to \infty,$$

as required.

An intriguing fact is that (4.5) can be alternatively derived from a little-known formula due to Ramanujan (2000), namely

$$\int_{-\infty}^{\infty} |\Gamma(a+\mathrm{i}\xi)|^2 \mathrm{e}^{\mathrm{i}x\xi} \,\mathrm{d}\xi = \frac{\sqrt{\pi}\,\Gamma(a)\Gamma(a+\frac{1}{2})}{\cosh^{2a}\left(\frac{x}{2}\right)}, \qquad a > 0.$$

Taking $a = (\alpha + 1)/2$, a trivial change of variable takes $g_{\alpha,\alpha}$ to the correct φ_0 and the proof follows by inverting the Fourier transform. Going in the reverse direction, we can obtain a the generalisation of Ramanujan's formula, to the Fourier transform of $\Gamma(a + i\xi)\Gamma(b - i\xi)$ for a, b > 0.

Except for Carlitz polynomials and their immediate generalisations, orthogonal polynomials associated with measures $d\mu_{\alpha,\beta}$ do not appear to have been studied in the literature. They might be an interesting object for further study, being examples of measures supported by the entire real line, yet distinct from the more familiar Freud-type measures.

5 Conclusions

We set ourselves a goal in this paper: to identify and characterise orthonormal systems, complete in $L_2(\mathbb{R})$ and with a skew-symmetric, tridiagonal, irreducible differentiation matrix whose expansion coefficients can be computed rapidly. In particular we are interested in the computation of the first N expansion coefficients in $\mathcal{O}(N \log_2 N)$ operations, utilising familiar transforms, e.g. FFT, FCT or FST.

In Section 2 we introduced the Tanh-Jacobi functions, of which the four cases where $\alpha, \beta = \pm \frac{1}{2}$ have expansion coefficients which can be computed using FCTs or FSTs. Subsequently, in Section 3 we described two kinds of half-range expansions (i.e., treating the even and the odd part of a function separately) which can be computed rapidly with either FCT or FST: the first based on a combination of Chebyshev polynomials of the first and the fourth kind, the second on such polynomials of the second and third kind.

Mathematically, the two approaches are identical when $\alpha = \beta$, although their computation is somewhat different. Which is preferable? As things stand, there is no clear answer (and things are complicated by the availability of yet another approach of this kind, using skew-Hermitian differentiation matrices, which will be described in (Iserles & Webb 2019*a*)). The full-range approximation has the virtue of simplicity, hence of easier implementation. A possible advantage of a half-range approximation is more subtle. Once \mathcal{D} approximates the first derivative, \mathcal{D}^2 approximates the second one and, provided \mathcal{D} is skew symmetric, \mathcal{D}^2 is negative semi-definite – this is in line with the Laplace operator being negative semi-definite and is vital for stability.

As a square of a skew-symmetric, tridiagonal matrix, \mathcal{D}^2 neatly separates even and odd functions. Specifically, let $\mathfrak{E} \oplus \mathfrak{O} = L_2(\mathbb{R}) \cap C^2(\mathbb{R})$ be a representation of square-integrable, twice differentiable functions on the line as a direct sum of even functions \mathfrak{E} and odd functions \mathfrak{O} . A derivative takes \mathfrak{E} to \mathfrak{O} and vice versa, hence a second derivative is invariant in both \mathfrak{E} and \mathfrak{O} . There is thus a virtue, at least once both first and second derivatives are present, to work separately in \mathfrak{E} and \mathfrak{O} , as is the case with half-range approximations.

Is simplicity preferable to even-odd separation? Are there additional considerations at play? By this stage it is impossible to provide a definitive answer. The purpose of the paper is to present a range of new results that improve our knowledge of approximation on the real line in the context of spectral methods.

The outlook for spectral methods on the real line using Tanh-Chebyshev-T functions appears promising. Consider the basic first order differential operator,

$$\mathcal{L}u(x) = u'(x) + a(x)u(x),$$

where a is a bounded function such that a rapidly convergent expansion of the form $a(x) = \sum_{m=0}^{\infty} a_m \tilde{T}_m(\tanh x)$ is possible². In coefficient space for the Tanh-Chebyshev-T functions, the operator \mathcal{L} becomes the infinite dimensional matrix $\mathcal{D} + \mathcal{A}$, where \mathcal{D} is the skew-symmetric tridiagonal differentiation matrix (1.3) with $b_m = \frac{1}{2}(m + \frac{1}{2})$,

²With the absence of the sech $\frac{1}{2}x$ weight, these functions need not be square-integrable on the real line, which is perfectly consistent with which functions a(x) allow \mathcal{L} to be a bounded operator on the Sobolev space $\mathrm{H}^{1}(\mathbb{R})$.

and

$$\mathcal{A} = \begin{bmatrix} a_0 & a_1 & a_2 & a_3 & \cdots \\ a_1 & a_0 & a_1 & a_2 & \ddots \\ a_2 & a_1 & a_0 & a_1 & \ddots \\ a_3 & a_2 & a_1 & a_0 & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \end{bmatrix} + \begin{bmatrix} a_1 & a_2 & a_3 & a_4 & \cdots \\ a_2 & a_3 & a_4 & a_5 & \cdot \\ a_3 & a_4 & a_5 & a_6 & \cdot \\ a_4 & a_5 & a_6 & a_7 & \cdot \\ \vdots & \ddots & \ddots & \ddots & \ddots \end{bmatrix}$$

The matrix is Toeplitz-plus-Hankel and, if a is sufficiently regular then the matrix is effectively banded, because if $a_m = 0$ for m > M for some integer M then the resulting operator has bandwidth M. This implies that an Olver–Townsend type approach for infinite-dimensional QR solution is possible in principle. An rth order differential operator with variable coefficients whose expansions have a maximum of M terms yields a matrix with bandwidth of at most r+M. Furthermore, the resulting matrices respect certain symmetries that the operator \mathcal{L} may have, something which is not true of the original Olver–Townsend ultraspherical spectral method. For example, if \mathcal{L} is self-adjoint on $L_2(\mathbb{R})$ then L is a self-adjoint operator on ℓ_2 .

One final, credible, but as of yet unexplored, application of this work is for the computation of the Fourier transform of certain functions on the real line. Let $f \in L_2(\mathbb{R})$ have a rapidly convergent expansion in one of the Tanh-Jacobi bases, $f(x) = \sum_{m=0}^{\infty} c_m \varphi_m^{(\alpha,\beta)}(x)$. An approximation $f_N(x) = \sum_{m=0}^{N} c_m^N \varphi_m^{(\alpha,\beta)}(x)$ computed using either fast polynomial transforms or the FCT/FST as discussed in Section 4. By the identity in equation (1.5) and (Iserles & Webb 2019*b*), the Fourier transform of f_N is an expansion in the generalised Carlitz polynomials weighted by $g_{\alpha,\beta}$ (see equation (4.2)),

$$\mathcal{F}[f](\xi) = g_{\alpha,\beta}(\xi) \sum_{m=0}^{N} (-i)^m c_m^N p_m(\xi).$$
(5.1)

By Theorem 6 of (Iserles & Webb 2019b), these generalised Carlitz polynomials, which are orthonormal with respect to $d\mu_{\alpha,\beta}(\xi)$ in Theorem 3, satisfy

$$p_{m+1}(\xi) = \frac{\xi}{b_m} p_m(\xi) - \frac{b_{m-1}}{b_m} p_{m-1}(\xi),$$
(5.2)

where b_m is given in equation (2.8). Clenshaw's algorithm can be used to evaluate this expansion at a single point $\xi \in \mathbb{R}$, using purely the coefficients $\{b_m\}_{m \in \mathbb{Z}_+}$ (Clenshaw 1955). This is similar to the work of Weber in which one computes a series expansion with orthogonal rational functions whose Fourier transforms are Laguerre functions, utilising the Fast Fourier Transform (Weber 1980).

Much work is needed to be done, particularly on understanding the approximation theory of the Tanh-Jacobi bases, but we hope these functions find application and interest well beyond that discussed in this paper.

References

Akhiezer, N. I. (1965), The classical moment problem: and some related questions in analysis, Vol. 5, Oliver & Boyd.

- Bader, P., Iserles, A., Kropielnicka, K. & Singh, P. (2016), 'Efficient methods for linear Schrödinger equation in the semiclassical regime with time-dependent potential', *Proc. Royal Soc. A.* 472(2193), 20150733, 18.
- Blanes, S., Casas, F. & Thalhammer, M. (2017), 'High-order commutator-free quasi-Magnus exponential integrators for non-autonomous linear evolution equations', *Comput. Phys. Commun.* 220, 243–262.
- Carlitz, L. (1959), 'Bernoulli and Euler numbers and orthogonal polynomials', Duke Math. J. 26, 1–15.
- Chihara, T. S. (1978), An Introduction to Orthogonal Polynomials, Gordon and Breach Science Publishers, New York–London–Paris. Mathematics and its Applications, Vol. 13.
- Clenshaw, C. (1955), 'A note on the summation of Chebyshev series', *Mathematics of Computation* **9**(51), 118–120.
- Deift, P., Kriecherbauer, T., McLaughlin, K. T.-R., Venakides, S. & Zhou, X. (1999), 'Strong asymptotics of orthogonal polynomials with respect to exponential weights', Comm. Pure Appl. Math. 52(12), 1491–1552.
- Duhamel, P. & Vetterli, M. (1990), 'Fast Fourier transforms: a tutorial review and a state of the art', Signal Process. 19(4), 259–299.
- Dutt, A., Gu, M. & Rokhlin, V. (1996), 'Fast algorithms for polynomial interpolation, integration, and differentiation', SIAM J. Numer. Anal. 33(5), 1689–1711.
- Fokas, A. S., Its, A. R. & Kitaev, A. V. (1992), 'The isomonodromy approach to matrix models in 2D quantum gravity', Comm. Math. Phys. 147(2), 395–430.
- Hairer, E. & Iserles, A. (2016), 'Numerical stability in the presence of variable coefficients', Found. Comput. Math. 16(3), 751–777.
- Hairer, E. & Iserles, A. (2017), 'Banded, stable, skew-symmetric differentiation matrices of high order', *IMA J. Numer. Anal.* 37(3), 1087–1103.
- Iserles, A. (2014), 'On skew-symmetric differentiation matrices', IMA J. Numer. Anal. 34(2), 435–451.
- Iserles, A. (2016), 'The joy and pain of skew symmetry', Found. Comput. Math. 16(6), 1607–1630.
- Iserles, A. & Webb, M. (2019a), A family of orthogonal rational functions and other orthogonal systems with a skew-Hermitian differentiation matrix, Technical report, DAMTP, University of Cambridge. To appear.
- Iserles, A. & Webb, M. (2019b), 'Orthogonal systems with a skew-symmetric differentiation matrix', Found. Comp. Maths. To appear.

- Iserles, A., Kropielnicka, K. & Singh, P. (2018), 'Magnus-Lanczos methods with simplified commutators for the Schrödinger equation with a time-dependent potential', *SIAM J. Numer. Anal.* 56(3), 1547–1569.
- Jin, S., Markowich, P. & Sparber, C. (2011), 'Mathematical and computational methods for semiclassical Schrödinger equations', Acta Numer. 20, 121–209.
- Olver, F. W. J., Lozier, D. W., Boisvert, R. F. & Clark, C. W., eds (2010), NIST Handbook of Mathematical Functions, U.S. Department of Commerce, National Institute of Standards and Technology, Washington, DC; Cambridge University Press, Cambridge. With 1 CD-ROM (Windows, Macintosh and UNIX).
- Olver, S. (2018), 'Approxfun. jl v0.10.3'.
- Olver, S. & Townsend, A. (2013), 'A fast and well-conditioned spectral method', SIAM Review 55(3), 462–489.
- Patriarca, M. (1994), 'Boundary conditions for the Schrödinger equation in the numerical simulation of quantum systems', *Physical Review E* 50(2), 1616.
- Rainville, E. D. (1960), Special Functions, The Macmillan Co., New York.
- Ramanujan, S. (2000), Some definite integrals [Messenger Math. 44 (1915), 10–18], in 'Collected papers of Srinivasa Ramanujan', AMS Chelsea Publ., Providence, RI, pp. 53–58.
- Shapiro, M. & Brumer, P. (2003), Principles of the Quantum Control of Molecular Processes, Wiley-Interscience, Hoboken, N.J.
- Townsend, A., Trogdon, T. & Olver, S. (2016), 'Fast computation of Gauss quadrature nodes and weights on the whole real line', *IMA J. Numer. Anal.* **36**(1), 337–358.
- Townsend, A., Webb, M. & Olver, S. (2018), 'Fast polynomial transforms based on Toeplitz and Hankel matrices', Math. Comp. 87(312), 1913–1934.
- Weber, H. (1980), 'Numerical computation of the Fourier transform using Laguerre functions and the fast Fourier transform', *Numerische Mathematik* **36**(2), 197–209.

A Two identities for Jacobi polynomials

Proposition 5 For every $\alpha > -1$, $m \in \mathbb{Z}_+$ and $X \in \mathbb{C}$

$$P_{2m}^{(\alpha,\alpha)}(\sqrt{1-X}) = \frac{(\alpha+m+1)_m m!}{(2m)!} P_m^{(\alpha,-\frac{1}{2})}(1-2X), \qquad m \in \mathbb{Z}_+.$$
(A.1)

Proof Exploiting the connection between $P_m^{(\alpha,\alpha)}$ and *Gegenbauer polynomials* C_m^{ν} (Rainville 1960, p. 276), we have

$$\mathbf{P}_{2m}^{(\alpha,\alpha)}(\sqrt{1-X}) = \frac{(1+\alpha)_{2m}}{(1+2\alpha)_{2m}} \mathbf{C}_{2m}^{\alpha+\frac{1}{2}}(\sqrt{1-X})$$

$$= \frac{(1+\alpha_{2m})}{(1+2\alpha)_{2m}} \sum_{\ell=0}^{m} \frac{(-1)^{\ell} (\alpha+\frac{1}{2})_{2m-\ell}}{\ell!(2m-2\ell)!} (2\sqrt{1-X})^{2(m-\ell)}$$

$$= (-1)^{m} \frac{(1+\alpha)_{2m}}{(1+2\alpha)_{2m}} \sum_{\ell=0}^{m} \frac{(-1)^{\ell} (\alpha+\frac{1}{2})_{m+\ell}}{(2\ell)!(m-\ell)!} 4^{\ell} (1-X)^{\ell}$$

$$= (-1)^{m} \frac{(1+\alpha)_{2m} (\alpha+\frac{1}{2})_{m}}{(1+2\alpha)_{2m} m!} {}_{2}F_{1} \begin{bmatrix} -m, \alpha_{m} + \frac{1}{2}; \\ \frac{1}{2}; \end{bmatrix},$$

while, using formula (3) from (Rainville 1960, p. 254),

$$\mathbf{P}_{m}^{(\alpha,-\frac{1}{2})}(1-2X) = (-1)^{m} \frac{(2m)!}{4^{m} m!^{2}} {}_{2}F_{1} \begin{bmatrix} -m, \alpha+m+\frac{1}{2}; \\ \frac{1}{2}; \end{bmatrix} \cdot \mathbf{1} - X \end{bmatrix}.$$

The proposition follows since

$$\frac{(1+\alpha)_{2m}(\alpha+\frac{1}{2})_m}{(1+2\alpha)_{2m}} = \frac{[(\alpha+1)(\alpha+2)\cdots(\alpha+2m)][(\alpha+\frac{1}{2})(\alpha+\frac{3}{2})\cdots(\alpha+m-\frac{1}{2})]}{(2\alpha+1)(2\alpha+2)\cdots(2\alpha+2m)} = \frac{[(\alpha+1)(\alpha+2)\cdots(\alpha+2m)][(2\alpha+1)(2\alpha+3)\cdots(2\alpha+2m-1)]}{4^m[(2\alpha+1)(2\alpha+3)\cdots(2\alpha+2m-1)][(\alpha+1)(\alpha+2)\cdots(\alpha+m)]} = \frac{(\alpha+m+1)_m}{4^m}.$$

	-	-	
			1

Proposition 6 For every $\alpha > -1$, $m \in \mathbb{Z}_+$ and $X \in \mathbb{C}$

$$P_{2m+1}^{(\alpha,\alpha)}(\sqrt{1-X}) = \frac{(\alpha+m+1)_{m+1}m!}{(2m+1)!}\sqrt{1-X}P_m^{(\alpha,\frac{1}{2})}(1-2X).$$
(A.2)

Proof Identical to the proof of the previous proposition.

Although this plays no role in this paper, it is interesting to recast (A.1) and (A.2) with the substitution $t = \sqrt{1-X}$, whereby, since $P_m^{(\alpha,\beta)}(-t) = (-1)^m P_m^{(\beta,\alpha)}(t)$ (Rainville 1960, p. 256), they become

$$P_{2m}^{(\alpha,\alpha)}(t) = (-1)^m \frac{m!(\alpha+m+1)_m}{(2m)!} P_m^{(-\frac{1}{2},\alpha)}(1-t^2),$$

$$P_{2m+1}^{(\alpha,\alpha)}(t) = (-1)^m \frac{m!(\alpha+m+1)_{m+1}}{(2m+1)!} t P_m^{(\frac{1}{2},\alpha)}(1-t^2).$$

We are not aware of this representation of ultraspherical polynomials in terms of Jacobi polynomials of (roughly) half the degree, which might be of an independent interest.