

# TWO-PHASE SEMILINEAR FREE BOUNDARY PROBLEM WITH A DEGENERATE PHASE

NORAYR MATEVOSYAN AND ARSHAK PETROSYAN

ABSTRACT. We study minimizers of the energy functional

$$\int_D [|\nabla u|^2 + \lambda(u^+)^p] dx$$

for  $p \in (0, 1)$  without any sign restriction on the function  $u$ . The distinguished feature of the problem is the lack of nondegeneracy in the negative phase. The main result states that in dimension two the free boundaries  $\Gamma^+ = \partial\{u > 0\} \cap D$  and  $\Gamma^- = \partial\{u < 0\} \cap D$  are  $C^{1,\alpha}$ -regular, provided  $1 - \epsilon_0 < p < 1$ . The proof is obtained by a careful iteration of the Harnack inequality to obtain a nontrivial growth estimate in the negative phase, compensating for the a priori unknown nondegeneracy.

## 1. INTRODUCTION AND MAIN RESULTS

1.1. **The problem.** In this paper we study a two-phase free boundary problem, obtained by minimizing the functional

$$(1.1) \quad J(u) := \int_D (|\nabla u|^2 + 2F(u)) dx$$

in an open subset  $D \subset \mathbb{R}^n$ , where  $F$  is a Hölder continuous function

$$(1.2) \quad F(u) := \lambda(u^+)^p, \quad \lambda > 0, \quad 0 < p < 1.$$

(Hereafter we denote  $u^\pm = \max\{\pm u, 0\}$ .) By a *minimizer*, we understand a function  $u \in W^{1,2}(D)$  such that

$$J(u) \leq J(v), \quad \text{for any } v \in u + W_0^{1,2}(D).$$

The existence of minimizers with a given Sobolev trace boundary data  $u_0 \in W^{1,2}(D)$  follows easily by the direct methods of the calculus of variations, however there is generally no uniqueness as  $J$  is not convex. The minimizers satisfy

$$\begin{aligned} \Delta u &= p\lambda u^{p-1} & \text{in } \Omega^+(u) &:= \{u > 0\} \\ \Delta u &= 0 & \text{in } \Omega^-(u) &:= \{u < 0\} \end{aligned}$$

and our objective is to study the interfaces or *free boundaries*

$$\Gamma^\pm(u) := \partial\Omega^\pm(u) \cap D.$$

---

2000 *Mathematics Subject Classification.* Primary 35R35.

*Key words and phrases.* Regularity of the free boundary, two-phase free boundary problem, lack of nondegeneracy, homogeneous global solutions.

N. Matevosyan is partially supported by the WWTF (Wiener Wissenschafts, Forschungs und Technologiefonds) project No. CI06 003 and by award No. KUK-I1-007-43, made by King Abdullah University of Science and Technology (KAUST).

A. Petrosyan is supported in part by NSF grant DMS-0701015.

The problem should be viewed in the general framework of two-phase free boundary problems, such as a chemical reaction problem with instantaneous reaction along the free boundary (interface), see e.g. [CH70, Wei01]. See also [BN80] for the one phase case.

The main distinguished feature of the problem considered in this paper is the *lack of nondegeneracy in the negative phase* in the sense that there is no a priori bound on how slowly  $u^-$  can grow near  $\Gamma^-$ . This creates a substantial difficulty in using the most essential tool - the rescaling of solutions (blowup) at an appropriate rate (see Section 3). If  $u^-$  would grow slower than that rate, the blowups would be identically zero, and the information carried by the negative phase would be lost in the blowup limit.

On the other hand, it is relatively easy to show that  $u^+(x)$  grows as  $\text{dist}(x, \Gamma^+)^{\beta}$  away from  $\Gamma^+$ ,  $\beta = 2/(2-p)$  for  $x$  near  $\Gamma^+$  on compact subsets of  $\Omega^+$ . This is a direct consequence of the positivity of  $\lambda$ .

The nondegenerate version of the problem has been recently studied by Lindgren and Petrosyan in [LP08], where the authors consider the minimizers of (1.1) with

$$F(u) = \lambda_+(u^+)^p + \lambda_-(u^-)^p, \quad \lambda_{\pm} > 0, \quad 0 < p < 1.$$

Because of the positivity of both  $\lambda_+$  and  $\lambda_-$ , both phases  $u^{\pm}$  grow as  $\text{dist}(x, \Gamma^{\pm})^{\beta}$  and one can apply Aleksandrov's reflection-comparison type argument (which heavily relies on nondegeneracy) to show that the free boundaries  $\Gamma^{\pm}$  are  $C^1$  regular in dimension  $n = 2$ . However, even in this nondegenerate case the problem is completely open in higher dimensions, which is related to the lack of the classification of homogeneous global solutions rather than the nondegeneracy (see Section 3).

The corresponding one-phase problem (i.e., nonnegative minimizers of  $u$ ) has been studied in a series of papers by Phillips [Phi83a, Phi83b] and Alt and Phillips [AP86]. In the latter paper it has been proved that there exists a singular set  $\Sigma \subset \Gamma^+$  of  $(n-1)$ -Hausdorff measure zero such that  $\Gamma^+ \setminus \Sigma$  is  $C^{\infty}$  (actually real analytic). Moreover, they have shown that in dimension  $n = 2$  the singular set  $\Sigma = \emptyset$ , i.e. the free boundary is *fully regular*.

**1.2. Main result.** The main result in this paper states that despite the lack of nondegeneracy in the negative phase, the free boundary is still fully regular for  $p$  near 1 in dimension  $n = 2$ .

**Theorem I.** *Let  $u$  be a minimizer of (1.1) in dimension  $n = 2$ . Then there exists an absolute constant  $\epsilon_0 > 0$  such that  $\Gamma^+(u)$  and  $\Gamma^-(u)$  are locally  $C^{1,\alpha}$  curves, provided  $1 - \epsilon_0 < p < 1$ . Moreover,  $\Gamma^-(u) \subset \Gamma^+(u)$ .*

The last part of the theorem basically says that the fattening of  $\{u = 0\}$  cannot occur between the phases  $\Omega^{\pm}$ , but only as the “dead core” in  $\Omega^+$ .

The paper is organized as follows.

- Section 2 contains preliminary results, such as the optimal regularity and the general structure of the free boundary.
- Section 3 gives the notion of rescalings and the blowups suitable for this problem. It also contains the proof of the nondegeneracy in the positive components and the classification of blowups in dimension 2. We should mention that the methods in this section are rather standard but of fundamental importance for the proof of the main theorem.




FIGURE 1.  $\Gamma^+ = \partial\{u > 0\} \cap D$ ,  $\Gamma^- = \partial\{u < 0\} \cap D$

- The heart of the paper is Section 4, more specifically Proposition 4.2, which deals with the lack of the nondegeneracy in the negative phase. The main idea consists in a careful iteration of the Harnack inequality in the negative phase, which provides with a nontrivial growth rate there, compensating for the apriori unknown nondegeneracy.

**Acknowledgment.** We would like to thank the anonymous referee for valuable comments that have contributed to the improvement of the paper.

## 2. PRELIMINARIES

**2.1. Optimal regularity and the Euler-Lagrange equation.** Perhaps the first question associated with the variational problem (1.1) is the optimal regularity of the minimizers. While studying the one-phase free boundary problem, Phillips [Phi83a] has established that the nonnegative minimizers of  $J$  satisfy

$$u \in C_{\text{loc}}^{1,\beta-1}(D), \quad \beta = \frac{2}{2-p}.$$

Later, Giaquinta and Giusti [GG84] have extended the above result to general minimizers of  $J$  with no restrictions on their sign. This is the best regularity possible, as the one-dimensional example  $u(x_1) = C_0(x_1^+)^{\beta}$  shows for a suitably chosen  $C_0 = C_0(\lambda, p) > 0$ .

The Euler-Lagrange equation associated with the variational problem (1.1) is

$$(2.1) \quad \Delta u = p\lambda(u^+)^{p-1}\chi_{\{u>0\}} \quad \text{in } D.$$

However, since even the local integrability of  $(u^+)^{p-1}\chi_{\{u>0\}}$  in  $D$  is apriori unknown, we have to specify in which sense we understand (2.1). But first notice that since  $\Omega^{\pm}(u) = \{\pm u > 0\}$  are open, the equation (2.1) is clearly satisfied in the classical sense in  $\Omega^{\pm}(u)$ , and moreover,  $u$  is real analytic there. In general, we understand (2.1) in the sense of *domain variation*:

$$\frac{d}{d\epsilon} \Big|_{\epsilon=0} J(u(x + \epsilon\phi(x))) = \int_D [2\nabla u \cdot D\phi \nabla u - (\text{div } \phi)(|\nabla u|^2 + 2F(u))] dx = 0,$$

for any  $\phi \in C_0^{\infty}(D, \mathbb{R}^n)$ .

**2.2. Structure of the free boundary.** Here we briefly discuss the structure of the free boundary (see also Fig. 1).

1° First let us note that  $\Gamma^- \subset \Gamma^+$ . Indeed, if  $x_0 \in \Gamma^- \setminus \Gamma^+$ , then  $u \leq 0$  in  $B_\delta(x_0)$  for some small  $\delta > 0$ . This implies that the term  $F(u)$  in the integrand of (1.1) vanishes identically in  $B_\delta(x_0)$ , and consequently that the minimizer  $u$  is harmonic there. But then by the maximum principle,  $u \equiv 0$  in  $B_\delta(x_0)$ , which contradicts to the fact that  $x_0 \in \partial\{u < 0\}$ . Hence,  $\Gamma^- \subset \Gamma^+$ . This essentially means that no thickening of the level set  $\{u = 0\}$  can occur between  $\Gamma^-$  and  $\Gamma^+$ , nor that there could be “dead cores” of  $\{u = 0\}$  in  $\{u \leq 0\}$ .

2° However, it is possible that  $\Gamma^+ \setminus \Gamma^-$  is nonempty, e.g. when  $u$  is nonnegative. In fact, for any  $x_0 \in \Gamma^+ \setminus \Gamma^-$  there exists a ball  $B_\delta(x_0) \subset D$  such that  $u \geq 0$  in  $B_\delta(x_0)$ . We call such  $x_0$  a *one-phase* free boundary point. The analysis of the regularity of the free boundary near one-phase points is reduced to the case already studied by Alt and Phillips [AP86]. In particular, in dimension  $n = 2$ ,  $\delta$  can be chosen so small that  $B_\delta(x_0) \cap \Gamma^+$  will be a real-analytic surface.

3° We say that  $x_0$  is a *two-phase* free boundary point, if

$$x_0 \in \Gamma^+ \cap \Gamma^- (= \Gamma^-).$$

We distinguish two types of two-phase points. The first kind is so-called *branching points*, where the condition

$$|\nabla u(x_0)| = 0$$

is satisfied. This terminology is reminiscent of the fact that in similar two-phase free-boundary problems this condition holds automatically at  $x_0 \in \Gamma^+ \cap \Gamma^- \cap \overline{\{u = 0\}^\circ}$ , i.e., at points  $x_0$  where the free boundary branches out to  $\Gamma^\pm$ . By 1° above, the branching in this narrow sense can never occur in our case; however, apriori we may not exclude the existence of two-phase points with vanishing gradient. *In fact, the proof of Theorem I consists in showing that such points don't exist when  $n = 2$  and  $1 - \epsilon_0 < p < 1$ .*

The second kind of two-phase points are the *non-branching points* is where

$$|\nabla u(x_0)| > 0.$$

Since  $u \in C_{\text{loc}}^{1,\beta-1}(D)$ , the implicit function theorem implies that for such points there exists a small  $\delta$  such that  $B_\delta(x_0) \cap \Gamma^+ = B_\delta(x_0) \cap \Gamma^-$  is a graph of a  $C^{1,\alpha}$  function.

### 3. RESCALINGS AND BLOWUPS

**3.1. Rescalings.** A fundamental tool that we will use throughout the article is the notion of “blowup” at a free boundary point.

For a minimizer  $u$  of (1.1), a point  $x_0 \in \Gamma^+ \cup \Gamma^- (= \Gamma^+)$  and  $r > 0$ , let us define the *rescaling*

$$u_{x_0,r}(x) := \frac{u(x_0 + rx)}{r^\beta}, \quad \text{for } \beta = \frac{2}{2-p} \quad \text{and } x \in D_{x_0,r} = \frac{1}{r}(D - x_0).$$

We will use the notation  $u_r$  for  $u_{x_0,r}$  if  $x_0 = 0$ . If  $x_0 \in (\Gamma^+ \cup \Gamma^-) \cap K$  for  $K \Subset D$  and is such that  $|\nabla u(x_0)| = 0$ , we have the uniform estimates

$$|u_{x_0,r}(x)| \leq C_K |x|^\beta, \quad \text{for } |x| \leq \frac{\delta}{r},$$

where  $\delta = \frac{1}{2} \text{dist}(K, \Gamma^+)$ . This follows from the optimal  $C_{\text{loc}}^{1, \beta-1}$ -regularity of  $u$ . Hence, for a fixed  $x_0$ , we may extract a sequence  $r_j \rightarrow 0$  such that

$$u_{x_0, r_j} \rightarrow u_0 \quad \text{in } C_{\text{loc}}^1(\mathbb{R}^n),$$

where  $u_0 \in C^{1, \beta-1}(\mathbb{R}^n)$ . We will call  $u_0$  a *blowup* of  $u$  at  $x_0$ . It is a simple exercise to show that  $u_0$  is a global minimizer of functional  $J$ , i.e. it minimizes  $J$  on every subdomain  $U \Subset \mathbb{R}^n$  among the functions in  $W^{1,2}(U)$  with the same trace on  $\partial U$  as  $u$ . Note that the blowup is not defined at free boundary points  $x_0$  where  $|\nabla u(x_0)| > 0$ , i.e. at non-branching points. Moreover, at points where blowups exist, it is not clear apriori if the blowup is unique. Namely, taking a different subsequence  $r'_j \rightarrow 0$  may result in convergence of  $u_{x_0, r'_j}$  to a different blowup  $u'_0$ . This may happen, e.g. when the free boundary “spirals” near  $x_0$ , see Fig. 2.




FIGURE 2. Possible nonuniqueness of blowups: spiraling free boundary

One of the most important properties of blowups is their homogeneity.

**Proposition 3.1** (Homogeneity of blowups). *Let  $u$  be a minimizer of (1.1) and  $x_0 \in (\Gamma^+ \cup \Gamma^-) \cap \{|\nabla u| = 0\}$ . Then any blowup  $u_0$  of  $u$  at  $x_0$  is a homogeneous function of degree  $\beta$  with respect to the origin, i.e.,*

$$u_0(rx) = r^\beta u_0(x), \quad x \in \mathbb{R}^n, \quad r > 0.$$

*Proof.* See [LP08, Proposition 3.2] for a quick proof based on the monotonicity formula of Weiss [Wei98].  $\square$

**3.2. Nondegeneracy in positive phase.** The following nondegeneracy properties of the minimizers in the positive phase will guarantee the nontriviality of blowups. In fact, it is the lack of a similar property in the negative phase that is the major complication in our problem.

**Lemma 3.2** (Nondegeneracy in positive phase). *Let  $u$  be a minimizer of (1.1) and let  $x_0 \in \Omega^+ \cup \Gamma^+$ . Then for any  $r > 0$  with  $B_r(x_0) \Subset D$  we have*

$$\sup_{\Omega^+ \cap \partial B_r(x_0)} u \geq c_0 r^\beta,$$

with  $c_0 = c_0(\lambda, n, p) > 0$ .

In fact, we will need the following more refined version of the nondegeneracy lemma.

**Lemma 3.3** (Nondegeneracy in positive components). *Let  $u$  be a minimizer of (1.1) and let  $x_0 \in D$ . For any  $B_r(x_0) \Subset D$ , let  $V$  be a connected component of  $\Omega_r^+ = \Omega^+ \cap B_r(x_0)$  such that  $V \cap B_{r/2}(x_0) \neq \emptyset$ . Then*

$$\sup_{\partial V \cap \partial B_r(x_0)} u \geq c_0 r^\beta$$

with  $c_0 = c_0(\lambda, n, p) > 0$ .

*Proof.* The proof is a slight modification of the one in [LP08], which in turn is similar to the one in [Phi83a], which follows the original idea in [CR76].

Fix a connected component  $V$  of  $\Omega^+ \cap B_r(x_0)$ . Then we can write

$$\partial V = E \cup F,$$

where

$$\begin{aligned} E &:= \partial V \cap B_r(x_0) \subset \Gamma^+ \\ F &:= \partial V \cap \partial B_r(x_0) \subset \partial B_r(x_0). \end{aligned}$$

Next, pick  $y_0 \in V \cap B_{r/2}(x_0)$  and consider

$$w(x) = |u(x)|^{2/\beta} - c|x - y_0|^2, \quad x \in D$$

for some constant  $c > 0$  to be specified later. Then by a direct computation, we have

$$\Delta w = (2/\beta)p\lambda + (2/\beta)(2/\beta - 1)|\nabla u|^2|u|^{-p} - 2nc \quad \text{in } V.$$

Hence, by choosing  $c = p\lambda/\beta n$ , we make  $\Delta w \geq 0$  in  $V$ . Since  $w(y_0) > 0$  and  $w$  is subharmonic, there must exist  $z_0 \in \partial V$  such that  $w(z_0) > 0$ . On the other hand, since  $w \leq 0$  on  $E \subset \Gamma^+$ , necessarily  $z_0 \in F$ , which gives that

$$\sup_F w > 0.$$

By construction we have that  $|x - y_0| \geq r/2$  for any  $x \in F \subset B_r(x_0)$  and therefore we obtain that

$$\sup_F u^{2/\beta} > cr^2/4,$$

which completes the proof of the lemma with  $c_0 = (c/4)^{\beta/2}$ .  $\square$

**3.3. Classification of homogeneous global minimizers.** Since the blowups of minimizers  $u$  are homogeneous of degree  $\beta$ , it would be desirable to obtain the classification of such global minimizers. This poses a challenging open problem even in the one-phase case in higher dimensions. In dimension  $n = 2$ , the problem is much simpler and reduces to identifying the solutions of an ODE with period  $2\pi$ .

**Proposition 3.4** (Classification of blowups). *Let  $u_0$  be a homogeneous global minimizer of  $J$  in dimension  $n = 2$ . Then after a suitable rotation of coordinate axes*

$$u_0(x) = C_0(x_1^+)^{\beta},$$

where  $C_0 = C_0(p) > 0$ .

We emphasize again that the above classification is not known in higher dimensions. The proof below is very specific for two dimensions and is based on the analysis of positivity and negativity sets of  $u_0$ , which are unions of cones. This is done in the next two lemmas. For convenience, we use polar coordinates in the statement of these lemmas.

**Lemma 3.5** (Positive solutions in angles). *Let  $u(r, \theta) = r^\beta f(\theta)$  be a positive solution of  $\Delta u = p\lambda u^{p-1}$  in the cone  $\mathcal{C}_\gamma = \{(r, \theta) : r > 0, \theta \in (0, \gamma)\}$ , vanishing continuously on  $\partial\mathcal{C}_\gamma$ :  $u(r, 0) = u(r, \gamma) = 0$ . Suppose also that  $u \in C^1(\overline{\mathcal{C}_\gamma})$ . Then*

$$\frac{\pi}{\beta} \leq \gamma \leq \pi.$$

*Furthermore, if  $f'(0) = 0$  or  $f'(\gamma) = 0$  then  $\gamma = \pi$ . Conversely, if  $\gamma = \pi$  then necessarily  $f'(0) = f'(\pi) = 0$ . Moreover, in this case  $f(\theta) = C_0(\sin \theta)^\beta$  for some  $C_0 = C_0(\lambda, p)$ .*

*Proof.* See the proof of [LP08, Lemma 4.2].  $\square$

**Lemma 3.6** (Negative solutions in angles). *Let  $u(r, \theta) = r^\beta f(\theta)$  be a negative harmonic function in the cone  $\mathcal{C}_\gamma$ , continuously vanishing on  $\partial\mathcal{C}_\gamma$ . Then*

$$\gamma = \frac{\pi}{\beta}$$

*and  $f(\theta) = -C \sin(\beta\theta)$  for some  $C > 0$ . In particular,  $|f'(0)| = |f'(\gamma)| > 0$ .*

*Proof.* Since  $u$  is harmonic when negative, the proof follows from elementary arguments in complex analysis.  $\square$

*Proof of Proposition 3.4.* Consider two cases:

1°  $0$  is a positive one-phase point, i.e.,  $0 \in \Gamma^+(u_0) \setminus \Gamma^-(u_0)$ . In this case  $u_0 \geq 0$ . Consider then the positivity set  $\Omega^+(u_0)$ . From the homogeneity, the connected components of  $\Omega^+$  are cones. Lemma 3.5 implies that the cones have openings between  $\pi/\beta$  and  $\pi$ . In fact, since  $|\nabla u_0| = 0$  on  $\Gamma^+$  for nonnegative solutions, the openings of the components of  $\Omega^+$  are exactly  $\pi$ . Hence, there are either two, or just one components of  $\Omega^+$  of opening  $\pi$ , which after a rotation, correspond to

$$u_0(x) = C_0|x_1|^\beta$$

and

$$u_0(x) = C_0(x_1^+)^beta,$$

respectively. The former case is actually impossible, since for nonnegative minimizers the zero set  $\{u_0 = 0\}$  must have nonzero Lebesgue density at free boundary points, see [Phi83b].

2°  $0$  is a two-phase point, i.e.,  $0 \in \Gamma^+(u_0) \cap \Gamma^-(u_0)$ . In this case both  $\Omega^+$  and  $\Omega^-$  are nonempty. By Lemmas 3.5–3.6 each component of  $\Omega^\pm$  is a cone of opening between  $\pi/\beta$  and  $\pi$ . Since  $\beta < 2$  there could be no more than 3 different components in  $\Omega^\pm$ .

If there are three components, then we have two possibilities: either there are two components of the same sign sharing a common side, or the set  $\{u = 0\}$  has a nonempty interior. In both cases,  $|\nabla u| = 0$  on one side of at least two of the components. By Lemmas 3.5–3.6, these components must be positive and have opening  $\pi$ . This doesn't leave space for the third component.

Hence, there are precisely two components, one necessarily in  $\Omega^+$ , the other in  $\Omega^-$ , since we assume that  $0$  is a two-phase free boundary point. Since the negative component must have opening  $\pi/\beta < \pi$  and the positive one at most  $\pi$ , the zero set  $\{u = 0\}$  will have nonempty interior and therefore  $|\nabla u| = 0$  on one side of both components. But then again by Lemmas 3.5–3.6, both components must be

positive, which is a contradiction. Thus, 0 cannot be a two-phase free boundary point.  $\square$

#### 4. PROOF OF THE MAIN THEOREM

The main ingredient in the proof of Theorem I is the following fact:

**Proposition 4.1.** *Let  $u$  be a minimizer of (1.1). Then there exist no free boundary branching points, i.e.,  $\Gamma^+ \cap \Gamma^- \cap \{|\nabla u| = 0\} = \emptyset$ , provided  $p$  is close enough to 1.*

In fact, we are going to reformulate this proposition as follows.

**Proposition 4.2.** *Let  $u$  be a minimizer of (1.1) in  $B_1$  and suppose that*

$$0 \in \Gamma^+, \quad |\nabla u(0)| = 0.$$

*If  $p$  is close enough to 1, then there exists a small  $r = r_u > 0$  such that  $u \geq 0$  in  $B_r$ .*

The proof of this proposition is subdivided into four steps.

1 $^\circ$  We start by showing that the free boundary becomes flatter as we approach to the origin.

**Claim 1.** *For any  $\sigma > 0$ , there exists an  $r_\sigma > 0$  such that for all  $0 < r < r_\sigma$  there exists a direction  $e_r \in \mathbb{R}^2$ ,  $|e_r| = 1$  such that*

$$\begin{aligned} u > 0 & \text{ in } \{x \cdot e_r > \sigma r\} \cap B_r =: D_{\sigma,r}^+, \\ u \leq 0 & \text{ in } \{x \cdot e_r < -\sigma r\} \cap B_r =: D_{\sigma,r}^-. \end{aligned}$$




FIGURE 3. Claim 1

*Proof.* The proof follows from the fact that any blowup is a halfspace solution and that  $u_+$  is nondegenerate. Indeed, suppose that the claim fails. Then there exist a sequence  $r_j \rightarrow 0+$  such that for the rescalings

$$u_{r_j}(x) = u_{0,r_j}(x) = \frac{u(r_j x)}{r_j^\beta}$$

at least one of the conditions

$$\begin{aligned} u_{r_j} &> 0 \text{ in } \{x \cdot e > \sigma\} \cap B_1, \\ u_{r_j} &\leq 0 \text{ in } \{x \cdot e < -\sigma\} \cap B_1 \end{aligned}$$

fails for any choice of the direction  $e$ . Passing to a subsequence, if necessary, we may assume that  $u_{r_j} \rightarrow u_0$  in  $C_{\text{loc}}^{1,\alpha}(\mathbb{R}^2)$ . By Propositions 3.1 and 3.4 we have that  $u_0(x) = C_0(x_1^+)^{\beta}$ , after a suitable rotation of coordinate axes. Then we claim that

$$\begin{aligned} u_{r_j} &> 0 \text{ in } \{x_1 > \sigma\} \cap B_1, \\ u_{r_j} &\leq 0 \text{ in } \{x_1 < -\sigma\} \cap B_1 \end{aligned}$$

for sufficiently large  $j$ . Indeed, the first inequality follows from the uniform convergence of  $u_{r_j} \rightarrow u_0$ . The second one is a direct corollary of the nondegeneracy (see Lemma 3.2): if  $\{u_{r_j} > 0\} \cap \{x_1 < -\sigma\} \cap B_1 \neq \emptyset$ , then  $\sup_{\{x_1 < -\sigma/2\} \cap B_2} u_{r_j} \geq c_0(\sigma/2)^{\beta}$  and consequently  $\sup_{\{x_1 < -\sigma/2\} \cap B_2} u_0 \geq c_0(\sigma/2)^{\beta}$ , which is a contradiction. Thus, we also arrive at a contradiction with the assumption made on  $u_{r_j}$ . This proves the claim.  $\square$

2° Since  $u$  is harmonic in  $\{u \leq 0\}^{\circ}$ , by the maximum principle, in  $D_{\sigma,r}^- = \{x \cdot e_r < -\sigma r\}$  there are two possibilities:

$$\begin{aligned} i^{\circ} \quad &u \equiv 0 \text{ in } D_{\sigma,r}^- \\ ii^{\circ} \quad &u < 0 \text{ in } D_{\sigma,r}^- \end{aligned}$$

and in principle different alternatives may hold for different  $r$ . However, our next claim says that the same alternative holds for all small  $r$ .

**Claim 2.** *If  $u < 0$  in  $D_{\sigma,r}^-$  for some  $r = r_0 < r_{\sigma}$ , then also  $u < 0$  in  $D_{\sigma,r}^-$  for all  $0 < r \leq r_0$  provided  $\sigma$  is small enough.*

*Proof.* Suppose  $r_0 \geq r \geq r_0(1 - \sigma)$ . From Claim 1 we know that  $D_{\sigma,r}^-$  cannot intersect  $D_{\sigma,r_0}^+$ . But then  $D_{\sigma,r}^-$  must intersect  $D_{\sigma,r_0}^-$ , since otherwise  $D_{\sigma,r}^-$  should be contained in a strip  $|x \cdot e_{r_0}| < \sigma r_0$ , which is too narrow if  $\sigma$  is small. This yields that  $u < 0$  in  $D_{\sigma,r}^-$ . By iteration, the claim follows.  $\square$




FIGURE 4. Claim 2

3° Now let us show, that

**Claim 3.** *If the exponent  $p \in (0, 1)$  in (1.1) is sufficiently close to 1, the alternative  $u < 0$  in  $D_{\sigma,r}^-$  for  $r < r_{\sigma}$  is not possible for small  $\sigma$ .*

*Proof.* Let  $\sigma = \epsilon/N$ , with  $\epsilon > 0$  small and  $N$  large to be specified later. For some  $r_0 < r_{\epsilon/N}$  consider the sequence of radii

$$r_n = r_0(1 - \epsilon)^n$$

as well as a sequence of points

$$P_n = -\frac{1}{2}r_n e_{r_n},$$

where  $e_r$ ,  $r = r_n$ , are as in Claim 1.

The proof consists of the repetitive application of the Harnack inequality, to obtain an estimate for  $|u(P_n)|$  from below, which will contradict to the growth estimate  $|u(x)| \leq C|x|^\beta$ . More detailed outline is as follows:

*Step 1:* Estimate the distance between  $P_n$  and  $P_{n+1}$ .

*Step 2:* Apply the Harnack inequality with precise constants to  $u$  in  $D_{\frac{\epsilon}{N}, r}^-$  at points  $P_n$  and  $P_{n+1}$  to obtain

$$(4.1) \quad -u(P_{n+1}) > -\lambda u(P_n),$$

which implies  $-u(P_n) > \alpha \lambda^n$ , where  $\alpha = -u(P_0)$

*Step 3:* On the other hand, we have that

$$|u(P_n)| \leq C r_n^\beta = C r_0^\beta (1 - \epsilon)^{\beta n}.$$

Thus, if we show that (4.1) holds with  $\lambda > (1 - \epsilon)^\beta$  then we will arrive at a contradiction.

We now start implementing this strategy.

**Step 1:** *Distance between  $P_n$  and  $P_{n+1}$ .*

Actually, we will estimate the distance between  $P_0$  and  $P_1$ , since the general case can be easily obtained from this by scaling.




FIGURE 5. Claim 3: Step 1

We start by estimating the rotation of the unit vector  $e_r$ . Namely, we want to find a control of

$$\theta = \text{angle}(e_{r_0}, e_{r_1}).$$

Since  $D_{\frac{\epsilon}{N}, r_1}^+$  cannot intersect  $D_{\frac{\epsilon}{N}, r_0}^-$ , using elementary geometry we obtain that

$$\begin{aligned}\sin \theta &\leq \frac{\epsilon/N}{1-\epsilon}, \\ \cos \theta &\geq \sqrt{1 - \frac{\epsilon^2/N^2}{(1-\epsilon)^2}}.\end{aligned}$$

Then, recalling that

$$P_0 = -\frac{1}{2}r_0e_{r_0}, \quad P_1 = -\frac{1}{2}r_0(1-\epsilon)e_{r_1},$$

we obtain

$$\begin{aligned}\text{dist}(P_0, P_1)^2 &= \frac{r_0^2}{4} (1 + (1-\epsilon)^2 - 2(1-\epsilon)\cos \theta) \\ &\leq \frac{r_0^2}{4} \left( 1 + (1-\epsilon)^2 - 2\sqrt{(1-\epsilon)^2 - \frac{\epsilon^2}{N^2}} \right) \\ &= \frac{r_0^2}{4} \left[ \left( 1 + \frac{1}{N^2} \right) \epsilon^2 + O(\epsilon^3) \right].\end{aligned}$$

This implies that

$$\text{dist}(P_0, P_1) < \frac{r_0\epsilon}{2}(1+\delta),$$

where  $\delta = \delta_{\epsilon, N} > 0$  can be made arbitrarily small if  $\epsilon > 0$  is sufficiently small and  $N$  is large.

**Step 2:** *The Harnack constant in dimension 2.*

To simplify the notations in this step we will assume  $r_0 = 1$  and  $e_{r_0} = (0, -1)$ . This will not affect the generality, since the estimates that we are going to use will be scale and rotation invariant. We start by observation that

$$B_{\frac{1}{2}-\frac{\epsilon}{N}}(P_0) \subset D_{\frac{\epsilon}{N}, 1}^-$$

and

$$P_1 \in B_{\frac{\epsilon}{2}(1+\delta)}(P_0).$$

Now, if  $u < 0$  satisfies

$$\Delta u = 0 \quad \text{in } B_R(P_0),$$

for a certain  $R > 0$  then the Harnack inequality for balls in  $\mathbb{R}^2$  gives

$$\frac{R-r}{R+r} \leq \frac{u(P_0)}{u(P_1)} \leq \frac{R+r}{R-r}.$$

In our case

$$R = \frac{1}{2} - \frac{\epsilon}{N} \quad \text{and} \quad r = \frac{\epsilon}{2}(1+\delta).$$

However, it turns out that this estimate is not enough for our purposes. Since  $u < 0$  actually in a larger set  $D_{\frac{\epsilon}{N}, 1}^-$ , we can improve this estimate as follows. Consider a bijective conformal mapping

$$\phi : D_{\frac{\epsilon}{N}, 1}^- \rightarrow B_R \quad \text{such that } \phi(P_0) = 0.$$

Since  $B_R(P_0) \subset D_{\frac{\epsilon}{N}, 1}^-$ , then using complex notations, we have

$$|\phi'(P_0)| < 1.$$




FIGURE 6. Claim 3: Step 2

Consequently, for small  $r$  we have

$$\phi(B_r(P_0)) \subset B_{(1-\kappa)r},$$

for some  $\kappa > 0$ , which can be regarded as an absolute constant if  $\epsilon/N$  is small. Applying the Harnack inequality to  $u \circ \phi^{-1}$  for balls  $B_{(1-\kappa)r} \subset B_R$ , we obtain that for our  $u$  we have

$$\frac{u(P_0)}{u(P_1)} \leq \frac{R + (1-\kappa)r}{R - (1-\kappa)r}.$$

Thus, we get

$$-u(P_1) \geq -\lambda u(P_0)$$

with

$$\lambda := \frac{R - (1-\kappa)r}{R + (1-\kappa)r} = \frac{1 - \frac{2\epsilon}{N} - (1-\kappa)(1+\delta)\epsilon}{1 - \frac{2\epsilon}{N} + (1-\kappa)(1+\delta)\epsilon}.$$

Similarly, we obtain that

$$-u(P_{n+1}) \geq -\lambda u(P_n)$$

for any  $n$  and as a corollary that

$$|u(P_n)| \geq \lambda^n |u(P_0)|.$$

**Step 3: Estimating  $\lambda$ .**

Using that  $(1+x)/(1-y) = (1+x)(1+y+O(y^2)) = 1+x+y+O(x^2+y^2)$ , we find that

$$\begin{aligned} \lambda &= \frac{1 - \frac{2\epsilon}{N} - (1-\kappa)(1+\delta)\epsilon}{1 - \frac{2\epsilon}{N} + (1-\kappa)(1+\delta)\epsilon} \\ &= 1 - \frac{2\epsilon}{N} - (1-\kappa)(1+\delta)\epsilon + \frac{2\epsilon}{N} - (1-\kappa)(1+\delta)\epsilon + O(\epsilon^2) \\ &= 1 - 2(1-\kappa)(1+\delta)\epsilon + O(\epsilon^2). \end{aligned}$$

Now recall that  $\kappa$  is an absolute constant and  $\delta$  can be chosen as small as we wish. Thus, we can make

$$(1 - \kappa)(1 + \delta) < 1 - \kappa/2,$$

and consequently

$$\lambda \geq 1 - 2(1 - \kappa/2)\epsilon + O(\epsilon^2) \geq (1 - \epsilon)^{2-\kappa/2},$$

provided  $\epsilon$  is sufficiently small. This implies that

$$|u(P_n)| \geq \lambda^n |u(P_0)| \geq c |P_n|^{2-\kappa/2},$$

with  $c > 0$ . On the other hand, we have the growth estimate

$$|u(P_n)| \leq C |P_n|^\beta$$

and to arrive at a contradiction we note that

$$\beta = \frac{2}{2-p} > 2 - \kappa/2$$

if  $p$  is close enough to 1.  $\square$

4° So far we have proved that for  $r \leq r_0$

$$\begin{aligned} u &> 0 \text{ in } D_{\sigma,r}^+ \\ u &\equiv 0 \text{ in } D_{\sigma,r}^- \end{aligned}$$

provided  $p$  is close enough to 1.

In this step, rescaling and rotating if necessary, we will assume that  $r_0 = 1$  and  $e_{r_0} = (1, 0)$ .

**Claim 4.** *Under the assumption above,  $u \geq 0$  in  $B_{\frac{1}{4}}$ .*

*Proof.* Assume the contrary and let  $W$  be a connected component of  $\{u < 0\} \cap B_1$  such that  $W \cap B_{\frac{1}{4}} \neq \emptyset$ . Observe that  $W$  must be contained in the strip  $\{|x_1| < \sigma\}$ .

We then consider two cases.

*i°*  $W \Subset B_{\frac{3}{4}}$ . Here we will have a contradiction to the maximum principle, since  $u$  is harmonic in  $W$  and must vanish on  $\partial W$ .


*ii°*  $W \cap \partial B_{\frac{3}{4}} \neq \emptyset$ . Then there exists a simple piecewise linear curve  $\gamma \subset W$  that joins  $\partial B_{\frac{1}{4}}$  and  $\partial B_{\frac{3}{4}}$ . Without loss of generality we may assume  $\gamma \subset \{x_2 > 0\}$ . The curve  $\gamma$  will divide the region

$$U := (B_{\frac{3}{4}} \setminus \overline{B_{\frac{1}{4}}}) \cap \{|x_1| < \frac{1}{4}, x_2 > 0\}$$

into two connected components  $U_+$  and  $U_-$  so that

$$U \setminus \gamma = U_+ \cup U_-, \quad U_+ \supset U \cap \{x_1 > \sigma\}, \quad U_- \supset U \cap \{x_1 < -\sigma\}.$$

(Note that here we have strongly used the topological properties of  $\mathbb{R}^2$ : in higher dimensions  $U \setminus \gamma$  or even  $U \setminus \overline{W}$  may still be connected.) Now pick a point  $P \in U_- \cap W$  with  $|P| = \frac{1}{2}$  such that there is a connected arc  $I$  of the circle  $\partial B_{\frac{1}{2}}$ , fully contained in  $U_-$ , which joins  $P$  and  $U \cap \{x_1 < -\sigma\}$ . Next, take an  $\epsilon$ -neighborhood  $I_\epsilon$  of the arc  $I$  for a small  $\epsilon > 0$  such that  $I \subset I_\epsilon \subset U_-$ . Then we claim that there exists a point  $Q \in I_\epsilon$  such that  $u(Q) > 0$ . Indeed, otherwise  $u \leq 0$  in  $I_\epsilon$  and thus  $u \leq 0$  in  $\{x_1 < -\sigma\} \cup I_\epsilon \cup W$ . However,  $u \equiv 0$  in  $\{x_1 < -\sigma\}$ ,  $u < 0$  in  $W$  and

FIGURE 7. Claim 4: alternative  $ii^\circ$ 

$u$  is harmonic in the union  $\{x_1 < -\sigma\} \cup I_\epsilon \cup W$ . This contradicts to the strong maximum principle.

Thus, there exists  $Q \in I_\epsilon \subset U_-$  with  $u(Q) > 0$ . Note that by our construction  $\frac{1}{2} - \epsilon \leq |Q| \leq \frac{1}{2} + \epsilon$  and consequently  $B_{\frac{1}{8}}(Q) \subset U$ , if  $\epsilon$  is small enough. Let now  $V$  be the connected component of the set  $\{u > 0\} \cap B_{\frac{1}{8}}(Q)$  that contains  $Q$ . Since  $Q \in V \subset U \setminus \gamma$  and  $Q \in U_-$ , necessarily  $V \subset U_-$  and consequently

$$V \subset \{|x_1| < \sigma\}.$$

Next, we apply the nondegeneracy in positive components to arrive at a contradiction. By Lemma 3.3 we must have

$$\sup_{\partial B_{\frac{1}{8}}(Q) \cap V} u \geq c_0 \left(\frac{1}{8}\right)^\beta.$$

On the other hand, since  $u \equiv 0$  in  $\{x_1 < -\sigma\}$ , and  $u \in C^{1, \beta-1}$ , we have

$$|u| \leq C_0 \sigma^\beta \quad \text{in } \{|x_1| < \sigma\} \cap B_1.$$

which is a contradiction if  $\sigma$  is small.

This proves the claim.  $\square$

*Proof of Proposition 4.2.* Note that Claim 4 is just the rescaled version of Proposition 4.2.  $\square$

We are now ready to prove the main theorem.

*Proof of Theorem I.* By Proposition 4.1, we have that  $|\nabla u|$  does not vanish on  $\Gamma^- = \Gamma^- \cap \Gamma^+$ , provided  $p$  is close enough to 1. Hence, by the implicit function theorem,  $\Gamma^-$  is locally a  $C^{1, \alpha}$  graph. On the other hand,  $u \geq 0$  in a neighborhood of any point on  $\Gamma^+ \setminus \Gamma^-$  and using the result of [AP86] in dimension 2 we conclude that  $\Gamma^+ \setminus \Gamma^-$  is  $C^\infty$  and therefore  $C^{1, \alpha}$ . This completes the proof of the theorem.  $\square$

## REFERENCES

- [AP86] H. W. Alt and D. Phillips, *A free boundary problem for semilinear elliptic equations*, J. Reine Angew. Math. **368** (1986), 63–107. MR850615 (88b:35206)
- [BN80] C.-M. Brauner and B. Nicolaenko, *On nonlinear eigenvalue problems which extend into free boundaries problems*, Bifurcation and nonlinear eigenvalue problems (Proc., Session, Univ. Paris XIII, Villetaneuse, 1978), Lecture Notes in Math., vol. 782, Springer, Berlin, 1980, pp. 61–100. MR572251 (83e:35127)
- [CR76] L. A. Caffarelli and N. M. Rivière, *Smoothness and analyticity of free boundaries in variational inequalities*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **3** (1976), no. 2, 289–310. MR0412940 (54 #1061)
- [CH70] J. R. Cannon and C. Denson Hill, *On the movement of a chemical reaction interface*, Indiana Univ. Math. J. **20** (1970/1971), 429–454. MR0279448 (43 #5170)
- [GG84] Mariano Giaquinta and Enrico Giusti, *Sharp estimates for the derivatives of local minima of variational integrals*, Boll. Un. Mat. Ital. A (6) **3** (1984), no. 2, 239–248 (English, with Italian summary). MR753882 (86g:49009)
- [LP08] Erik Lindgren and Arshak Petrosyan, *Regularity of the free boundary in a two-phase semilinear problem in two dimensions*, Indiana Univ. Math. J. **57** (2008), no. 7, 3397–3417. MR2492237
- [Phi83a] Daniel Phillips, *A minimization problem and the regularity of solutions in the presence of a free boundary*, Indiana Univ. Math. J. **32** (1983), no. 1, 1–17. MR684751 (84e:49012)
- [Phi83b] ———, *Hausdorff measure estimates of a free boundary for a minimum problem*, Comm. Partial Differential Equations **8** (1983), no. 13, 1409–1454. MR714047 (85b:35068)
- [Wei98] Georg S. Weiss, *Partial regularity for weak solutions of an elliptic free boundary problem*, Comm. Partial Differential Equations **23** (1998), no. 3-4, 439–455. MR1620644 (99d:35188)
- [Wei01] G. S. Weiss, *An obstacle-problem-like equation with two phases: pointwise regularity of the solution and an estimate of the Hausdorff dimension of the free boundary*, Interfaces Free Bound. **3** (2001), no. 2, 121–128, DOI 10.4171/IFB/35. MR1825655 (2002c:35275)

DEPARTMENT OF APPLIED MATHEMATICS AND THEORETICAL PHYSICS, UNIVERSITY OF CAMBRIDGE, CAMBRIDGE CB3 0WA, UK

*E-mail address:* `n.matevosyan@damtp.cam.ac.uk`

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN 47907

*E-mail address:* `arshak@math.purdue.edu`