

3P1b **Quantum Field Theory: Example Sheet 2** Michaelmas 2024

Corrections and suggestions should be emailed to ac2553@cam.ac.uk.

1. Consider a real scalar field with Lagrangian

$$\mathcal{L} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m^2 \phi^2 . \quad (1)$$

Show that, after normal ordering, the conserved four-momentum $P^\mu = \int d^3x T^{0\mu}$ takes the operator form

$$P^\mu = \int \frac{d^3p}{(2\pi)^3} p^\mu a_{\vec{p}}^\dagger a_{\vec{p}} , \quad (2)$$

where $p^0 = E_{\vec{p}}$ in this expression. From Eq. (2), verify that if $\phi(x)$ is now in the Heisenberg picture, then

$$[P^\mu, \phi(x)] = -i \partial^\mu \phi(x) .$$

2. For a real scalar field with Lagrangian (1), show that in the Heisenberg picture,

$$\dot{\phi}(x) = i[H, \phi(x)] = \pi(x) \quad \text{and} \quad \dot{\pi}(x) = i[H, \pi(x)] = \nabla^2 \phi(x) - m^2 \phi(x) .$$

Hence show that the operator $\phi(x)$ satisfies the Klein-Gordon equation.

3. Let $\phi(x)$ be a real scalar field in the Heisenberg picture with Lagrangian (1). Show that the relativistically normalised states $|p\rangle = \sqrt{2E_{\vec{p}}} a_{\vec{p}}^\dagger |0\rangle$ satisfy

$$\langle 0 | \phi(x) | p \rangle = e^{-ip \cdot x} .$$

- 4* In Example Sheet 1, you showed that the classical angular momentum of a field is given by

$$Q_i = \frac{1}{2} \epsilon_{ijk} \int d^3x (x^j T^{0k} - x^k T^{0j}) .$$

Write down the explicit form of the angular momentum for a free real scalar field with Lagrangian as in Eq.(1). Show that, after normal ordering, the quantum operator Q_i can be written as

$$Q_i = \frac{i}{2} \epsilon_{ijk} \int \frac{d^3p}{(2\pi)^3} a_{\vec{p}}^\dagger \left(p^j \frac{\partial}{\partial p_k} - p^k \frac{\partial}{\partial p_j} \right) a_{\vec{p}} .$$

Hence confirm that the quanta of the scalar field have spin zero (i.e. a one-particle state $|\vec{p}\rangle$ has zero angular momentum in its rest frame).

5. Show that the time ordered product $T(\phi(x_1)\phi(x_2))$ and the normal ordered product $:\phi(x_1)\phi(x_2):$ are both symmetric under the interchange of x_1 and x_2 . Deduce that the Feynman propagator $\Delta_F(x_1 - x_2)$ has the same symmetry property.
- 6* The Schwinger-Dyson equation states that

$$(\square_x + m^2) \langle \phi_x \phi_1 \dots \phi_n \rangle = \langle \mathcal{L}'_{\text{int}}[\phi_x] \phi_1 \dots \phi_n \rangle - i \sum_{j=1}^n \delta^{(4)}(x - x_j) \langle \phi_1 \dots \phi_{j-1} \phi_{j+1} \dots \phi_n \rangle, \quad (3)$$

where $\phi_j \equiv \phi(x_j)$ and $\phi_x \equiv \phi(x)$. Recall that the brackets stand for a shorthand of time-ordering, i.e.,

$$\langle \phi_1 \dots \phi_n \rangle \equiv \langle \Omega | T(\phi_1 \dots \phi_n) | \Omega \rangle,$$

and for simplicity we are assuming that the interacting part of the Lagrangian (\mathcal{L}_{int}) does not include derivatives of the fields.

In lectures we showed (3) for two fields for a QFT that is local and causal. Derive explicitly (3) for three fields by using the same assumptions.

7. Consider the scalar Yukawa theory given by the Lagrangian

$$\mathcal{L} = \partial_\mu \psi^* \partial^\mu \psi + \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - M^2 \psi^* \psi - \frac{1}{2} m^2 \phi^2 - g \psi^* \psi \phi.$$

Calculate the amplitude for meson decay $\phi \rightarrow \psi \bar{\psi}$ to leading order in g using the LSZ formula and the Schwinger-Dyson equation.

Show that the amplitude is only non-zero for $m > 2M$ and explain the physical interpretation of this condition using conservation laws.

8. *Optional: The interaction picture (Srednicki 9.5). See also Sec 3.1 and 3.7 of DT.*

In this problem, we will derive a formula for $\langle \Omega | T(\phi_1 \dots \phi_n) | \Omega \rangle$ using time-dependent perturbation theory for an interacting theory involving a single massive scalar field.

Suppose we have a Hamiltonian density $\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_1$, where

$$\mathcal{H}_0 = \frac{1}{2} \pi^2 + \frac{1}{2} (\nabla \phi)^2 + \frac{1}{2} m^2 \phi^2, \quad (4)$$

and \mathcal{H}_1 is a function of $\pi(0, \vec{x})$ and $\phi(0, \vec{x})$ and their spatial derivatives.

The ground state of the whole system is $|\Omega\rangle$, and a constant is added to the Hamiltonian such that $H|\Omega\rangle = 0$. The ground state of H_0 is given by $|0\rangle$ and a constant is also added such that $H_0|0\rangle = 0$. The Heisenberg picture field is

$$\phi(t, \vec{x}) = e^{iHt} \phi(0, \vec{x}) e^{-iHt}. \quad (5)$$

We also define the interaction-picture field as

$$\phi_I(t, \vec{x}) = e^{iH_0 t} \phi(0, \vec{x}) e^{-iH_0 t}. \quad (6)$$

- (a) Show that $\phi_I(x)$ obeys the Klein-Gordon equation, and hence is a free field.
 (b) Show that

$$\phi(x) = U^\dagger(t)\phi_I(x)U(t) ,$$

where $U(t) = e^{iH_0t}e^{-iHt}$ is unitary.

- (c) Show that $U(t)$ obeys the differential equation

$$i\frac{d}{dt}U(t) = H_I(t)U(t) ,$$

where $H_I(t) = e^{iH_0t}H_1e^{-iH_0t}$ is the interaction Hamiltonian in the interaction picture.

- (d) \mathcal{H}_1 was specified by a particular function of the fields $\pi(0, \vec{x})$ and $\phi(0, \vec{x})$. Show that $\mathcal{H}_I(t)$ is given by the same function of the interaction-picture fields $\pi_I(t, \vec{x})$ and $\phi_I(t, \vec{x})$.
 (e) Show that, for $t > 0$,

$$U(t) = \text{T exp} \left(-i \int_0^t dt' H_I(t') \right)$$

is a solution to the differential equation (7). What is the comparable expression for $t < 0$?

- (f) Define $U(t_2, t_1) := U(t_2)U^\dagger(t_1)$. Show that, for $t_2 > t_1$,

$$U(t_2, t_1) = \text{T exp} \left(-i \int_{t_1}^{t_2} dt' H_I(t') \right)$$

What is the comparable expression for $t_1 > t_2$?

- (g) For any time ordering, show that

$$\begin{aligned} U(t_3, t_1) &= U(t_3, t_2)U(t_2, t_1) , \\ U^\dagger(t_1, t_2) &= U(t_2, t_1) , \end{aligned}$$

and

$$\begin{aligned} U^\dagger(t, 0) &= U^\dagger(\infty, 0)U(\infty, t) , \\ U(t, 0) &= U(t, -\infty)U(-\infty, 0) . \end{aligned}$$

- (h) Replace H_0 with $(1 - i\epsilon)H_0$, and show that $\langle \Omega | U^\dagger(\infty, 0) = \langle \Omega | 0 \rangle \langle 0 |$ and that $U(-\infty, 0) | \Omega \rangle = | 0 \rangle \langle 0 | \Omega \rangle$.
 (i) Show that

$$\phi(x_n) \dots \phi(x_1) = U^\dagger(t_n, 0)\phi_I(x_n)U(t_n, t_{n-1})\phi_I(x_{n-1}) \dots U(t_2, t_1)\phi_I(x_1)U(t_1, 0) ,$$

and

$$\langle \Omega | \phi(x_n) \dots \phi(x_1) | \Omega \rangle = | \langle \Omega | 0 \rangle |^2 \langle 0 | U(\infty, t_n)\phi_I(x_n)U(t_n, t_{n-1})\phi_I(x_{n-1}) \dots U(t_2, t_1)\phi_I(x_1)U(t_1, -\infty) | 0 \rangle .$$

(j) Show that

$$\langle \Omega | T(\phi(x_n) \dots \phi(x_1)) | \Omega \rangle = |\langle \Omega | 0 \rangle|^2 \langle 0 | T(\phi_I(x_n) \dots \phi_I(x_1)) e^{-i \int d^4x \mathcal{H}_I} | 0 \rangle .$$

(k) Show that

$$|\langle \Omega | 0 \rangle|^2 = \left(\langle 0 | T e^{-i \int d^4x \mathcal{H}_I} | 0 \rangle \right)^{-1} .$$

Thus we have

$$\langle \Omega | T(\phi(x_n) \dots \phi(x_1)) | \Omega \rangle = \frac{\langle 0 | T(\phi_I(x_n) \dots \phi_I(x_1)) e^{-i \int d^4x \mathcal{H}_I} | 0 \rangle}{\langle 0 | T e^{-i \int d^4x \mathcal{H}_I} | 0 \rangle} .$$

The right-hand side of this equation involves only interaction-picture fields, hence one can Taylor expand the exponentials and use free-field theory to compute the resulting correlation functions.