Asymptotic Methods: Example Sheet 2

Send corrections to Henrik Latter at: hl278@cam.ac.uk

Michaelmas 2026

Copyright © 2025 University of Cambridge. Not to be quoted or reproduced without permission.

Two more integrals of Laplace type

1. Obtain the first correction to the Stirling formula in the asymptotic expansion of the Gamma function, i.e.,

$$\Gamma(x+1) \sim \sqrt{2\pi x}\Big(\frac{x}{e}\Big)^x\bigg(1+\frac{1}{12x}+\cdots\bigg), \quad \text{as } x\to\infty.$$

2. Derive the expansion

$$\int_0^{\pi/2} e^{x\sin^2 t}\,dt \sim \frac{e^x}{2}\sum_{m=0}^{\infty}(-1)^m\frac{\Gamma(1/2)}{\Gamma(1/2-m)\,m!}\frac{\Gamma(1/2+m)}{x^{m+1/2}}, \quad \text{for } x\to\infty.$$

   By means of a change of variables $t\mapsto \pi/2-t$ and the identity[1]

$$\Gamma(z)\Gamma(1-z)=\frac{\pi}{\sin\pi z},$$

   or otherwise, show that

$$\int_0^{\pi/2} e^{-x\sin^2 t}\,dt \sim \Big(\frac{\pi}{4x}\Big)^{1/2}\bigg(1+\frac{1}{1!}\frac{1^2}{4x}+\frac{1}{2!}\frac{1^2\cdot 3^2}{(4x)^2}+\cdots+\frac{1}{n!}\frac{1^2\cdot 3^2\cdots(2n-1)^2}{(4x)^n}+\cdots\bigg),$$

   as $x\to\infty$. Use this to find an asymptotic expansion for the Bessel function, defined by

$$I_0(x)=\frac{1}{\pi}\int_0^{\pi} e^{x\cos\theta}\,d\theta,$$

   in the same limit.

Oscillatory integrals and Stationary Phase

3. (a) Assume $a<c<b$ and let $f(t)$ be a function that is smooth in $(a,c)\cup(c,b)$ but has a discontinuity at $t=c$. To be precise, assume that, for all $n\in\{0,1,2,\dots\}$, the limits of the nth order derivative $f^{(n)}(t)$ as $t\to a^+$, c^-, c^+, and b^- exist and are designated $f^{(n)}(a^+)$, $f^{(n)}(c^-)$, $f^{(n)}(c^+)$, and $f^{(n)}(b^-)$ respectively. Find the asymptotic expansion, as $|\omega|\to\infty$, of

$$I(\omega)=\int_a^b f(t)e^{i\omega t}\,dt.$$

[1] See the course on Further Complex Methods to derive this identity.

(b) By taking the appropriate limits in part (a), find the asymptotic expansion of $I(\omega) = \int_{-\infty}^{\infty} f(t)e^{i\omega t}dt$, as $|\omega| \to \infty$, when

$$f(t) = \begin{cases} -e^t & t < 0 \\ e^{-t} & t \geq 0. \end{cases}$$

Compare your result with the exact expression for $I(\omega)$.

4. Review Stokes' problem from the lectures on stationary phase (also see p. 22 in Stuart's notes). Obtain the leading asymptotic behaviour of

$$\int_a^{\infty} f(t) \exp\big(ix(t^3 - t)\big)\, dt,$$

as $x \to \infty$, where f is smooth and $f \to 0$ as $t \to \pm\infty$, in the two cases: $a = -1/\sqrt{3}$, and $a = 1$.

5. Show that, as $x \to +\infty$,

$$\int_0^{\pi} \exp[ix(t - \sin t)]\, dt \sim e^{i\pi/6}\left(\frac{6}{x}\right)^{1/3}\Gamma\left(\frac{4}{3}\right).$$

How would this result differ if the lower limit of the integral were $-\pi$?

6. Taking $x \to \infty$, find the leading term in the asymptotic approximations of

(a) $\displaystyle\int_0^1 \cos(xt^p)\, dt,$ with $p > 1$ real,

(b) $\displaystyle\int_0^{\pi/2}\left(1 - \frac{2\theta}{\pi}\right)^{\gamma}\cos(x\cos\theta)\, d\theta,$ for $\gamma = 0$, $\gamma = -1/2$, and $\gamma = -3/4$.

Method of Steepest Descent

7. For θ real and positive, the function $f(\theta)$ is defined by

$$f(\theta) = \frac{1}{2\pi i}\int_{\mathcal{C}} \exp\left(\theta\left(t + \frac{1}{3}t^3\right)\right) dt,$$

where the path $\mathcal{C}$ lies in the complex plane, beginning at ∞ in the sector $-\pi/2 < \arg(t) < -\pi/6$ and ending at ∞ in the sector $\pi/6 < \arg(t) < \pi/2$. Find the two saddle points and show that the two paths of steepest *descent* through these points are

$$x = +\left(\frac{2+y}{3y}\right)^{1/2}(y-1), \quad y > 0$$

and

$$x = -\left(\frac{y-2}{3y}\right)^{1/2}(y+1), \quad y < 0,$$

where $t = x + iy$. You should justify carefully your choice of signs for the square roots. Show that, as $\theta \to \infty$,

$$f(\theta) = (\pi\theta)^{-1/2}\cos\left(\frac{2}{3}\theta - \frac{\pi}{4}\right) + O(\theta^{-1}).$$

Copyright © 2025 University of Cambridge. Not to be quoted or reproduced without permission.

8. Use the method of steepest descent to obtain the first two non-zero terms in the asymptotic approximation

$$\int_0^\infty \exp\left(ix\left(\frac{1}{3}t^3 + t\right)\right) dt \sim i\left(\frac{1}{x} + \frac{2}{x^3} + \cdots + \frac{a_n}{x^n} + \cdots\right),$$

as $x \to +\infty$. Check your answer via an integration by parts argument.

($*$) Find an expression for a_n, for all n.

9. Let $h(t) = i(t + t^2)$. Sketch the path through the point $t = 0$ for which $\mathrm{Im}(h(t)) = \text{const}$. Also sketch the path through the point $t = 1$ for which $\mathrm{Im}(h(t)) = \text{const}$.

By integrating along these paths, show that

$$\int_0^1 t^{-1/2} \exp\big(i\lambda(t + t^2)\big)\, dt \sim \frac{c_1}{\lambda^{1/2}} + c_2 \frac{e^{2i\lambda}}{\lambda} + \cdots,$$

as $\lambda \to \infty$, where the constants c_1 and c_2 are to be determined.

10. ($*$) Apply the method of steepest descent to the integral

$$I(k) = \int_{\gamma - i\infty}^{\gamma + i\infty} \frac{\exp(k(z - 2z^{1/2}))}{z - c}\, dz,$$

for the case $k \to +\infty$. Here the path of integration is parallel to the imaginary axis, and $\gamma > 1$ is a real constant. The branch cut for $z^{1/2}$ is the negative real axis. Show that the two parametrised curves $\tau \mapsto z_\pm(\tau)$, given by

$$z_\pm(\tau) = 1 - \tau^2 \pm 2i\tau, \quad 0 \le \tau < \infty,$$

are the steepest descent paths emanating from the saddle-point $z = 1$. Verify that they form two halves of a parabola crossing the real axis at the saddle point and find the equation of the parabola in real form.

Investigate the asymptotics of $I(k)$ as $k \to +\infty$ in the following cases:

(a) c is real and < 1;

(b) c is real and $1 < c < \gamma$;

(c) $c = ib$, with b real and $b > 2$.

[You may find it convenient to use τ as a variable of integration.]

Copyright © 2025 University of Cambridge. Not to be quoted or reproduced without permission.