

Example Sheet 4

1. Verify the Jacobi identity for Poisson brackets,

$$\{f, \{g, h\}\} + \{g, \{h, f\}\} + \{h, \{f, g\}\} = 0.$$

2. A particle with mass m , position \mathbf{r} and momentum \mathbf{p} has angular momentum $\mathbf{L} = \mathbf{r} \times \mathbf{p}$. Evaluate $\{x_i, L_j\}$, $\{p_i, L_j\}$, $\{L_i, L_j\}$ and $\{L_i, |\mathbf{L}|^2\}$.

The Laplace–Runge–Lenz vector is defined as

$$\mathbf{A} = \mathbf{p} \times \mathbf{L} - mk \hat{\mathbf{r}},$$

where k is a constant and $\hat{\mathbf{r}} = \mathbf{r}/|\mathbf{r}|$. Show that $\{L_i, A_j\} = \epsilon_{ijk} A_k$. For a system described by the Hamiltonian

$$H = \frac{|\mathbf{p}|^2}{2m} - \frac{k}{|\mathbf{r}|},$$

show, using Poisson brackets, that \mathbf{A} is conserved.

3. A particle of charge q moves in a time-independent background magnetic field \mathbf{B} . Show that $\{m\dot{x}_i, m\dot{x}_j\} = q\epsilon_{ijk} B_k$ and $\{x_i, m\dot{x}_j\} = \delta_{ij}$.

A *magnetic monopole* is a particle that produces a radial magnetic field of the form

$$\mathbf{B} = g \frac{\hat{\mathbf{r}}}{r^2},$$

where g is a constant and $\hat{\mathbf{r}} = \mathbf{r}/|\mathbf{r}|$. Consider a charged particle moving in the background of the magnetic monopole. Define the generalized angular momentum,

$$\mathbf{J} = m \mathbf{r} \times \dot{\mathbf{r}} - qg \hat{\mathbf{r}}.$$

Show that $\{\mathbf{J}, H\} = \mathbf{0}$. Why does this imply that \mathbf{J} is conserved?

4. In the lectures we constructed canonical transformations using generating functions. Consider canonical transformations $\mathbf{q} \mapsto \mathbf{Q}(\mathbf{q}, \mathbf{p})$, $\mathbf{p} \mapsto \mathbf{P}(\mathbf{q}, \mathbf{p})$ from the following perspective. Define the $2n$ -dimensional vector $\mathbf{x} = (q_1, \dots, q_n, p_1, \dots, p_n)^\top$ and the $2n \times 2n$ matrix

$$\Omega = \begin{pmatrix} 0 & \mathbf{I}_n \\ -\mathbf{I}_n & 0 \end{pmatrix},$$

where each entry is itself an $n \times n$ matrix.

- (a) Write Hamilton's equations for $\dot{\mathbf{x}}$ in terms of Ω and the Hamiltonian H .

(b) Hence deduce the following equation for the vector $\mathbf{X} = (Q_1, \dots, Q_n, P_1, \dots, P_n)^\top$:

$$\dot{\mathbf{X}} = (\mathbf{J}\Omega\mathbf{J}^\top) \frac{\partial H}{\partial \mathbf{X}},$$

where $J_{ab} = \partial X_a / \partial x_b$ ($a, b = 1, \dots, 2n$) is the Jacobian matrix of the transformation. This implies that, if the Jacobian of a transformation satisfies

$$\mathbf{J}\Omega\mathbf{J}^\top = \Omega,$$

then Hamilton's equations are invariant under that transformation. The transformations with such a Jacobian (said to be *symplectic*) are canonical.

(c) Use the above conclusion to prove that, if the Poisson bracket structure is preserved, then the transformation is canonical.

5. Show that the following transformations are canonical:

$$(a) \quad P = \frac{1}{2}(p^2 + q^2), \quad Q = \arctan\left(\frac{q}{p}\right),$$

$$(b) \quad P = \frac{1}{q}, \quad Q = pq^2,$$

$$(c) \quad P = 2\sqrt{q}(1 + \sqrt{q} \cos p) \sin p, \quad Q = \log(1 + \sqrt{q} \cos p).$$

6. Show that the following transformation is canonical, for any constant λ :

$$\begin{aligned} q_1 &= Q_1 \cos \lambda + P_2 \sin \lambda, & q_2 &= Q_2 \cos \lambda + P_1 \sin \lambda, \\ p_1 &= -Q_2 \sin \lambda + P_1 \cos \lambda, & p_2 &= -Q_1 \sin \lambda + P_2 \cos \lambda. \end{aligned}$$

Given that the original Hamiltonian is

$$H(\mathbf{q}, \mathbf{p}) = \frac{1}{2} (q_1^2 + q_2^2 + p_1^2 + p_2^2),$$

determine the new Hamiltonian $H(\mathbf{Q}, \mathbf{P})$. Hence solve for the dynamics, subject to the constraints $Q_2 = P_2 = 0$.

7. A group of particles, all of the same mass m , have initial heights z_0 and vertical momenta p_0 lying in the rectangle $-a \leq z_0 \leq a$, $-b \leq p_0 \leq b$ in phase space. The particles fall freely in a uniform gravitational field for a time t . Find the region of phase space in which they lie at time t , and show by direct calculation that its area is still $4ab$.

8. A *Poisson structure* on \mathbb{R}^n is an antisymmetric matrix ω^{ab} whose components depend on the coordinates $\xi^a \in \mathbb{R}^n$, $a = 1, \dots, n$, and such that the Poisson bracket

$$\{f, g\} = \sum_{a=1}^n \sum_{b=1}^n \omega^{ab}(\xi) \frac{\partial f}{\partial \xi^a} \frac{\partial g}{\partial \xi^b}$$

satisfies the Jacobi identity.

(a) Show that

$$\{fg, h\} = f\{g, h\} + \{f, h\}g.$$

(b) Assuming that the matrix ω is invertible, show that the antisymmetric matrix $W = \omega^{-1}$ satisfies

$$\partial_a W_{bc} + \partial_c W_{ab} + \partial_b W_{ca} = 0,$$

where $\partial_a = \frac{\partial}{\partial \xi^a}$. [Hint: Note that $\omega^{ab} = \{\xi^a, \xi^b\}$.]

(c) Set $\xi^a = (x, y, z)$. Show that

$$\{x, y\} = z, \quad \{y, z\} = x, \quad \{z, x\} = y$$

defines a Poisson structure on \mathbb{R}^3 , and find Hamilton's equations corresponding to a Hamiltonian $H = Ax^2 + By^2 + Cz^2$, where A , B and C are non-zero constants.

9. Explain what is meant by an *adiabatic invariant* for a mechanical system with one degree of freedom.

A light string passes through a small hole in the roof of a lift compartment of a very high skyscraper, and a small weight is attached to the lower end. Initially, the lift is at rest and the system behaves like a simple pendulum executing small oscillations. Construct a Hamiltonian for the system and use the theory of adiabatic invariants to discuss what happens to the frequency, linear and angular amplitudes of the motion if:

- (a) the lift begins to move upwards with slowly increasing acceleration, with the string attached at the hole;
- (b) the lift stays at rest, but the string is slowly withdrawn through the roof.

10. Consider a system with Hamiltonian

$$H = \frac{p^2}{2m} + \lambda q^{2n},$$

where λ is a positive constant and n is a positive integer. Show that the action variable I and the energy E are related by

$$E = \lambda^{1/(n+1)} \left(\frac{n\pi I}{J_n} \right)^{2n/(n+1)} \left(\frac{1}{2m} \right)^{n/(n+1)},$$

where $J_n = \int_0^1 (1-x)^{1/2} x^{(1-2n)/2n} dx$.

Consider a particle that moves in a potential $V(q) = \lambda q^4$. Assuming that λ varies slowly with time, show that the particle's total energy E is proportional to $\lambda^{1/3}$. Conversely, in the case that λ is fixed, show that the period of the motion is proportional to $(\lambda E)^{-1/4}$.

11. A pulsar of mass m moves in a planar orbit around a luminous supergiant star with mass $M \gg m$. You may regard the supergiant as being fixed at the origin of a plane-polar coordinate system (r, θ) , and the neutron star as moving in a central potential $V(r) = -GMm/r$. Construct the Hamiltonian for the motion, and show that p_θ and the total energy E are constants of motion.

The neutron star is in a non-circular orbit with $E < 0$. Give an expression for the adiabatic invariant $J(E, p_\theta, M)$ associated with the radial motion. The supergiant is steadily losing mass in a radiatively driven wind. Show that, over a long timescale, we have $E \propto M^2$.

Eventually the supergiant becomes a supernova, throwing off its outer layers on a short timescale, and leaving behind a remnant black hole of mass $M/2$. Explain why the theory of adiabatic invariants cannot be used to calculate the new orbit.

[You may find the following integral helpful:

$$\int_{r_1}^{r_2} \left[\left(1 - \frac{r_1}{r}\right) \left(\frac{r_2}{r} - 1\right) \right]^{1/2} dr = \frac{\pi}{2}(r_1 + r_2) - \pi\sqrt{r_1 r_2},$$

where $0 < r_1 < r_2$.]

12. [optional, based on 2010 Paper 4, Section II, Question 15D]

A system is described by the Hamiltonian $H(q, p, t)$. Define the *Poisson bracket* $\{f, g\}$ of two functions $f(q, p, t)$ and $g(q, p, t)$. Show from Hamilton's equations that

$$\frac{df}{dt} = \{f, H\} + \frac{\partial f}{\partial t}.$$

Consider the Hamiltonian

$$H = \frac{1}{2} (p^2 + \omega^2 q^2),$$

where $\omega = \omega(t)$, and define

$$a = \frac{p - i\omega q}{\sqrt{2\omega}}, \quad a^* = \frac{p + i\omega q}{\sqrt{2\omega}},$$

where $i^2 = -1$. Evaluate $\{a, a\}$ and $\{a, a^*\}$, and show that $\{a, H\} = -i\omega a$ and $\{a^*, H\} = i\omega a^*$. Show further that, when $f(q, p, t)$ is regarded as a function of the independent complex variables (a, a^*) and of t , one has

$$\frac{df}{dt} = i\omega \left(a^* \frac{\partial f}{\partial a^*} - a \frac{\partial f}{\partial a} \right) - \frac{1}{2} \frac{\dot{\omega}}{\omega} \left(a \frac{\partial f}{\partial a^*} + a^* \frac{\partial f}{\partial a} \right) + \frac{\partial f}{\partial t}.$$

Deduce that, in the case $d\omega/dt = 0$, both $(\log a^* - i\omega t)$ and $(\log a + i\omega t)$ are constant during the motion.

Consider now the case in which $\omega(t)$ varies slowly with time. Writing $f = (H/\omega)$, show that the time-average of (df/dt) over one period, $(2\pi/\omega)$, is approximately zero (that is, to order $(\dot{\omega}^2, \ddot{\omega})$). [Hint: You might like to start by writing $a = A(t)e^{-i\omega t} = A(0)e^{-i\omega t} + O(\dot{\omega})$.]