

## Vector Calculus: Example Sheet 3

David Tong, February 2022

1. Consider the line integral

$$I = \oint_C -x^2y \, dx + xy^2 \, dy$$

for  $C$  a closed curve traversed anti-clockwise in the  $(x, y)$ -plane.

(i) Evaluate  $I$  when  $C$  is a circle of radius  $R$  centred at the origin. Use Green's theorem to relate the results for  $R = b$  and  $R = a$  to an area integral over an appropriate region, and calculate the area integral directly.

(ii) Now suppose  $C$  is the boundary of a square centred at the origin with sides of length  $\ell$ . Show that  $I$  is independent of the orientation of the square in the  $(x, y)$ -plane.

2. Verify Stokes' theorem for the hemispherical shell  $S = \{x^2 + y^2 + z^2 = 1, z \geq 0\}$ , and the vector field

$$\mathbf{F}(\mathbf{x}) = (y, -x, z).$$

3. By applying Stokes' theorem to the vector field  $\mathbf{a} \times \mathbf{F}$  for  $\mathbf{a}$  constant, or otherwise, show that for a vector field  $\mathbf{F}(\mathbf{x})$

$$\oint_C d\mathbf{x} \times \mathbf{F} = \int_S (d\mathbf{S} \times \nabla) \times \mathbf{F}$$

where  $C = \partial S$ . Verify this result when  $C$  is the unit square in the  $(x, y)$ -plane with opposite vertices at  $(0, 0, 0)$  and  $(1, 1, 0)$  and  $\mathbf{F}(\mathbf{x}) = \mathbf{x}$ .

4. Let  $S = \{\mathbf{x} : |\mathbf{x}| = 1\}$  be the surface of a unit sphere. For the vector field

$$\mathbf{F}(\mathbf{x}) = \frac{\mathbf{x}}{r^3}$$

where  $r = |\mathbf{x}|$ , compute the integral  $\int_S \mathbf{F} \cdot d\mathbf{S}$ . Deduce that there *does not* exist a vector potential for  $\mathbf{F}$ , i.e. there can be no  $\mathbf{A}$  for which  $\mathbf{F} = \nabla \times \mathbf{A}$ . Compute  $\nabla \cdot \mathbf{F}$  and comment on your result.

5\*. Consider the following vector field

$$\mathbf{A}(\mathbf{x}) = \frac{1}{(x^2 + y^2)r} (yz, -xz, 0)$$

where  $r = |\mathbf{x}|$ . Compute  $\nabla \times \mathbf{A}$ . Does this contradict the result of Question 4? Apply Stokes' theorem to  $\nabla \times \mathbf{A}$  on the open surface

$$S_\epsilon = \{\mathbf{x} : |\mathbf{x}| = 1, x^2 + y^2 \geq \epsilon^2\}$$

How does this help reconcile the existence of  $\mathbf{A}$  with the result of Question 4?

6. Use Gauss' flux method to find the electric field  $\mathbf{E} = \mathbf{E}(\mathbf{x})$  due to a spherically symmetric charge density

$$\rho(r) = \begin{cases} 0 & 0 \leq r \leq a \\ \rho_0 r/a & a < r < b \\ 0 & r \geq b \end{cases}$$

Now find the electric potential  $\phi = \phi(r)$  *directly* from Poisson's equation by writing down the general, spherically symmetric solution to Laplace's equation in each of the intervals  $0 < r < a$ ,  $a < r < b$  and  $r > b$ , and adding a particular integral where necessary. You should assume that  $\phi$  and  $\phi'$  are continuous at  $r = a$  and  $r = b$ . Check this solution gives rise to the same electric field using  $\mathbf{E} = -\nabla\phi$ .

7. The scalar field  $\psi(r)$  only depends on  $r = |\mathbf{x}|$ . Use Cartesian coordinates and suffix notation to show

$$\nabla\psi = \psi'(r)\frac{\mathbf{x}}{r} \quad \text{and} \quad \nabla^2\psi = \psi''(r) + \frac{2}{r}\psi'(r).$$

Verify this result using your expression for the Laplacian in spherical polar coordinates. Find a non-singular, spherically symmetric solution to the equation  $\nabla^2\psi = 1$  for  $r < R$  subject to the requirement that  $\psi(R) = 1$ .

8. Consider a complex valued function  $f = \phi(x, y) + i\psi(x, y)$  satisfying  $\partial f/\partial\bar{z} = 0$ , where  $\partial/\partial\bar{z} = \frac{1}{2}(\partial/\partial x + i\partial/\partial y)$ . Show that  $\nabla^2\phi = \nabla^2\psi = 0$ . Show also that a curve on which  $\phi$  is constant is orthogonal to a curve on which  $\psi$  is constant, at a point where they intersect. Find  $\phi$  and  $\psi$  when  $f = ze^z$ ,  $z = x + iy$ , and compare with Question 4 on Examples Sheet 2.

**9a.** Using Cartesian coordinates  $(x, y)$ , find all solutions of Laplace's equation  $\nabla^2\psi = 0$  in two dimensions of the form  $\psi(x, y) = f(x)e^{\alpha y}$ , with  $\alpha$  constant. Hence find a solution on the region  $0 < x < a$  and  $y > 0$  with boundary conditions:

$$\psi(0, y) = \psi(a, y) = 0 \quad \text{and} \quad \psi(x, 0) = \lambda \sin(\pi x/a)$$

and  $\psi(x, y) \rightarrow 0$  as  $y \rightarrow \infty$ .

**b.** Using the formula for the 2d Laplacian in plane polar coordinates  $(r, \theta)$ , verify that Laplace's equation in the plane has solutions of the form  $\psi(r, \theta) = Ar^\alpha \cos \beta\theta$ , if  $\alpha$  and  $\beta$  are related appropriately. Hence find solutions on the following regions, with the given boundary conditions ( $\lambda$  a constant):

(i)  $r < R$  with  $\psi(R, \theta) = \lambda \cos \theta$ ,

(ii)  $r > R$  with  $\psi(R, \theta) = \lambda \cos \theta$  and  $\psi(r, \theta) \rightarrow 0$  as  $r \rightarrow \infty$ ,

(iii)  $a < r < b$  with  $\mathbf{n} \cdot \nabla\psi(a, \theta) = 0$  and  $\psi(b, \theta) = \lambda \cos 2\theta$ .

**10.** Let  $\psi$  and  $\phi$  be scalar functions. Using an integral theorem, establish *Green's second identity*

$$\int_V (\phi \nabla^2 \psi - \psi \nabla^2 \phi) dV = \int_{\partial V} (\phi \nabla \psi - \psi \nabla \phi) \cdot d\mathbf{S}$$

**11.** Show that the solution to the following boundary value problem is unique on  $V$ :

$$-\nabla^2\psi + \psi = \rho(\mathbf{x})$$

with  $\mathbf{n} \cdot \nabla\psi = f(\mathbf{x})$  on  $\partial V$ .

**12.** Show that the Laplace equation  $\nabla^2\psi = 0$  has a unique solution on  $V$  when subjected to the boundary condition on  $\partial V$

$$(\mathbf{n} \cdot \nabla\psi)g(\mathbf{x}) + \psi = f(\mathbf{x})$$

where  $g(\mathbf{x}) \geq 0$  on  $\partial V$ . Find a non-zero solution to Laplace's equation on  $|\mathbf{x}| \leq 1$  which satisfies the boundary conditions above with  $f = 0$  and  $g = -1$  on  $|\mathbf{x}| = 1$ .

**13.** Let  $u$  be harmonic on  $V$  and  $v$  a smooth function that satisfies  $v = 0$  on  $\partial V$ . Show that

$$\int_V \nabla u \cdot \nabla v \, dV = 0.$$

Now if  $w$  is any function on  $V$  with  $w = u$  on  $\partial V$ , show, by considering  $v = w - u$ , that

$$\int_V |\nabla w|^2 \, dV \geq \int_V |\nabla u|^2 \, dV.$$

**14\*.** Show that a harmonic function  $\psi$  at the point  $\mathbf{a}$  is equal to the average of its values on the interior of the ball  $B_r(\mathbf{a}) = \{\mathbf{x} : |\mathbf{x} - \mathbf{a}| < r\}$ , for any  $r > 0$ . By considering  $\nabla\psi$  and the previous result for large  $r$ , or otherwise, prove that if  $\psi$  is bounded and harmonic on  $\mathbb{R}^3$  then it is constant.

**15\*.** For a volume  $V$  with smooth boundary  $S$ , establish the identity

$$\text{vol}(V) = \frac{1}{3} \int_S \mathbf{x} \cdot d\mathbf{S}$$

Suppose now that  $V = V(t)$ , and the velocity of a point  $\mathbf{x} \in V$  is  $\mathbf{v}(\mathbf{x})$ . Show that

$$\frac{d}{dt} \text{vol}(V) = \int_S \mathbf{v} \cdot d\mathbf{S}.$$

Using this result, or otherwise, obtain *Reynold's Transport Theorem* for a scalar function  $\rho(\mathbf{x}, t)$ :

$$\frac{d}{dt} \int_{V(t)} \rho \, dV = \int_{V(t)} \frac{\partial \rho}{\partial t} \, dV + \int_{S(t)} \rho(\mathbf{v} \cdot d\mathbf{S}).$$

Interpret this result.

[Hint: it is better to think physically about this problem rather than simply trying to manipulate equations.]